

Medication Administration Protocol: AUBAGIO

Document Control Information

Document ID:	MED-PROT-6992
Effective Date:	
Review Date:	2025-10-14
Therapeutic Class:	Adverse Event Report
Risk Level:	Critical

Standard Dosing Protocol

Current Standard Dose: Standard dosing per label

Monitoring Requirements: Standard warnings

Administration Guidelines

Route of Administration: IM

Frequency: Three times daily

Duration: 14 days

Patient Population: Patients similar to case: 2, age 48

Contraindications and Precautions

Contraindications: Enhanced screening for risk factors

Adverse Events: Muscle spasms, Condition aggravated, Pain

Detailed Clinical Considerations

Patient Monitoring Protocol

Regular monitoring is essential for safe administration of AUBAGIO. Current protocols require Standard warnings. Healthcare providers should assess patient response and adjust therapy accordingly. Baseline laboratory values should be obtained prior to initiation. The mechanism of action involves AUBAGIO mechanism with identified safety concern. Clinical efficacy has been demonstrated in Muscle spasms, Condition aggravated, Pain.

Special Populations

Elderly Patients: Dose adjustment may be required based on renal function and overall health status.

Pediatric Patients: Safety and efficacy not established in patients under 18 years.

Pregnancy: Use only if potential benefit justifies potential risk to fetus.

Nursing Mothers: Exercise caution when administering to nursing women.

Last revised: June 2024