

# Termodünaamika: molekulaarkinemaatika

Saskia, 28. september 2021

$$k = R/N_A$$

Eelmisest korrast:

$$pV = \nu RT \quad (1)$$

Üheaatomilistel gaasidel (He) on  $i=3$ , kaheaatomilistel ( $H_2$ )  $i=5$  ning mitmeaatomilistel ( $CH_4$ )  $i=6$  vabadusastet.

Tähised:

$p$  - gaasi rõhk,  $V$  - gaasi ruumala,  $T$  - temperatuur

$\nu$  - moolide arv,  $M$  - molaarmass

$i$  - vabadusaste

$U$  - siseenergia

$K_k$  - kineetiline energia

$R = 8,3145 \text{ J/mol}\cdot\text{K}$  ideaalgaasi konstant

$k = 1,3806 \text{ J/K}$  Boltzmanni konstant

$N_A = 6.022 \cdot 10^{23}$  molekul/mol Avogadro arv

$v_{rms}$  - osakese ruutkeskmise (root mean square) kiirus

Ruutkeskmise leidmine:  $x_{rms} = \frac{1}{n} \sum_{i=1}^n x_i^2$

## 1 Rõhk

Tõesta, et gaasi rõhu arvutamiseks kehtib valem

$$p = \frac{\nu M v_{rms}^2}{iV} \quad (2)$$

## 2 Gaasiosakese kineetiline energia

Tõesta, et molekuli kineetilise energia arvutamiseks kehtib valem

$$K_k = \frac{i}{2} kT \quad (3)$$

## 3 Siseenergia

Tõesta, et gaasi siseenergia arvutamiseks kehtib valem

$$U = \frac{i}{2} \nu RT \quad (4)$$

## 4 Disk in Gas (EuPhO 2018 Pr2)

Consider a thin flat disk of mass  $M$  and face area  $S$  at temperature  $T_1$  resting initially in weightlessness in a gas of mass density  $\rho$  at temperature  $T_0$  ( $T_1 = 1000T_0$ ).

One of the faces of the disk is covered with a thermally insulating layer, the other face has a very good thermal contact with the surrounding gas: gas molecules of mass  $m$  obtain the temperature of the disk during a single collision with the surface.

Estimate the initial acceleration  $a_0$  and maximal speed  $v_{max}$  of the disk during its subsequent motion.

Assume the heat capacity of the disk to be on the order of  $Nk_B$ , where  $N$  is the number of atoms in it, and  $k$  is the Boltzmann constant, and molar masses of the gas and the disk's material to be of the same order. The mean free path length of molecules is much larger than the size of the disk.

Neglect any edge effects occurring at the edge of the disk.

## 5 Physics Cup 2012 Pr3

Determine or estimate the net heat flux density  $P$  between two parallel plates at distance  $L$  from each other, which are at temperatures  $T_1$  and  $T_2$ , respectively. The space between the plates is filled with a monoatomic gas of molar density  $n$  and of molar mass  $M$ . You may use the following approximations:

1. The gas density is so low that the mean free path  $\lambda \gg L$
2.  $T_1 \gg T_2$
3. When gas molecules bounce from the plates, they obtain the temperature of the respective plates instantaneously