

Machine Learning

Homework 1

Due on March 29, 2018

1. Let

$$A_+ = \{(0, 0), (1, 1), (-1, 1), (1, -1), (-1, -1)\}$$

and

$$A_- = \{(1, 0), (-1, 0), (0, 1), (0, -1)\}$$

represent the *positive* and *negative* training instances respectively.

- (a) Plot the decision boundary for the 1-nearest neighbor algorithm.
 - (b) What is the training set accuracy for 3-nearest neighbor algorithm?
 - (c) What is the *confusion* matrix for 3-nearest neighbor algorithm on the training set?
2. Let the $A \in R^{n \times n}$ be a symmetric positive definite matrix. Show that all eigenvalues of matrix A are positive.
3. Let $Z = [X_1; X_2; X_3]$ be a random vector and Σ be a matrix with size 3×3 where $\Sigma_{ij} = Cov(X_i, X_j)$ and $\Sigma_{ii} = Var(X_i)$. Let the random variable $W = \mathbf{a}^\top Z = a_1 X_1 + a_2 X_2 + a_3 X_3$ where $\mathbf{a} = [a_1; a_2; a_3]$. *i.e.*, the random variable W is the *projection* of random vector Z onto the vector \mathbf{a} . Find the variance of W .

4. Let S be a set of 10,000 random numbers generated by the *uniform distribution*, $U[0;1]$. You have to estimate the *mean* of this *uniform distribution*.
- (a) Randomly select 10 number from the set S and then use the average of these 10 number as the estimation. Repeat this experiment 20 times. What is the *sample mean* and *sample standard deviation* of these 20 random experiments?
- (b) Randomly select 1,000 number from the set S and then use the average of these 1,000 number as the estimation. Repeat this experiment 50 times. What is the *sample mean* and *sample standard deviation* of these 50 random experiments?

5. Consider a linear system of equations as follows:

$$\begin{array}{rclcl} 2x_1 & + & 2x_2 & & = & 5 \\ x_1 & + & & x_3 & = & 2 \\ & & 2x_2 & + & 2x_3 & = & 3 \\ x_1 & + & & & x_3 & = & 4 \end{array}$$

Find the least squares approximation solution for it.

6. Generate a training dataset with size 1000 by yourself.
- (a) That is, $S = \{(\mathbf{x}^i, y_i) | \mathbf{x}^i = (\mathbf{x}_1^i, \mathbf{x}_2^i) \in R^2, \text{ and } y_i \in R, i = 1, \dots, 1000\}$, where \mathbf{x}_1 and \mathbf{x}_2 are generated by the *uniform distribution*, $U[-1;1]$. The observation value $y = 2\mathbf{x}_1^2 - 0.6\mathbf{x}_2^2 + 1.5\mathbf{x}_1\mathbf{x}_2 + \mathbf{x}_1 + 2\mathbf{x}_2 + \epsilon$ where ϵ is the random noise generated by $N(0, 1)$.
- (b) Find a *quadratic* function $f(\mathbf{x})$, that is fitted in the training dataset S .
- (c) Compute the *MAE*, mean of absolute error, and plot the function you get.