

Properties of Collinear

1 10th Maths - Chapter 7

This is Problem-2 from Exercise 7.3.2

1. In each of the following find the value of 'k', for which the points are collinear.
 1. $(7, -2), (5, 1), (3, k)$
 2. $(8, 1), (k, -4), (2, -5)$.

2 Solution for problem :1

$$\mathbf{D} = (\mathbf{A} - \mathbf{B}) = \left(\begin{pmatrix} 7 \\ -2 \end{pmatrix} - \begin{pmatrix} 5 \\ 1 \end{pmatrix} \right) = \begin{pmatrix} 2 \\ -3 \end{pmatrix} \quad (1)$$

$$\mathbf{E} = (\mathbf{A} - \mathbf{C}) = \left(\begin{pmatrix} 7 \\ -2 \end{pmatrix} - \begin{pmatrix} 3 \\ K \end{pmatrix} \right) = \begin{pmatrix} 4 \\ 12 - K \end{pmatrix} \quad (2)$$

If points on a line are collinear, rank of matrix is " 1 " then the vectors are linearly dependent. For 2×2 matrix Rank =1 means Determinant is 0.

Through pivoting, we obtain

$$\mathbf{F} = \begin{pmatrix} \mathbf{D} \\ \mathbf{E} \end{pmatrix} \quad (3)$$

$$\begin{pmatrix} 2 & -3 \\ 4 & -2 - k \end{pmatrix} \quad (4)$$

$$\begin{pmatrix} 2 & -3 \\ 4 & -2-k \end{pmatrix} \xrightarrow{R2=R2-2R1} \begin{pmatrix} -2-k & -4 \\ 4 & -2-k \end{pmatrix} \quad (5)$$

if the rank of the matrix is 1 means any one of the row must be zero. So, making the first element in the matrix to 0.

$$-2 - k - 2(-3) = 0 \quad (6)$$

$$-2 - k - 6 = 0 \quad (7)$$

$$k = 4 \quad (8)$$

Hence proved.




Figure 1

3 Solution for problem :2

$$\mathbf{A} = \begin{pmatrix} 8 \\ 1 \end{pmatrix} \mathbf{B} = \begin{pmatrix} k \\ -4 \end{pmatrix} \mathbf{C} = \begin{pmatrix} 2 \\ -5 \end{pmatrix} \quad (9)$$

$$\mathbf{D} = (\mathbf{A} - \mathbf{B}) = \left(\begin{pmatrix} 8 \\ 1 \end{pmatrix} - \begin{pmatrix} k \\ -4 \end{pmatrix} \right) = \begin{pmatrix} 8-k \\ 5 \end{pmatrix} \quad (10)$$

$$\mathbf{E} = (\mathbf{A} - \mathbf{C}) = \left(\begin{pmatrix} 8 \\ 1 \end{pmatrix} - \begin{pmatrix} 2 \\ -5 \end{pmatrix} \right) = \begin{pmatrix} 6 \\ 6 \end{pmatrix} \quad (11)$$

If points on a line are collinear, rank of matrix is " 1 " then the vectors are linearly dependent. For 2×2 matrix Rank = 1 means Determinant is 0.

Through pivoting, we obtain

$$\mathbf{F} = \begin{pmatrix} \mathbf{D} \\ \mathbf{E} \end{pmatrix} \quad (12)$$

$$\begin{pmatrix} 8-k & 5 \\ 6 & 6 \end{pmatrix} \quad (13)$$

$$\begin{pmatrix} 8-k & 5 \\ 6 & 6 \end{pmatrix} \xrightarrow{R1=3R1-R2} \begin{pmatrix} 24-3k & 15 \\ 6 & 6 \end{pmatrix} \quad (14)$$

if the rank of the matrix is 1 means any one of the row must be zero. So, making the first element in the matrix to 0.

$$3(8-k) - 3(5) = 0 \quad (15)$$

$$24 - 3k - 15 = 0 \quad (16)$$

$$k = 3 \quad (17)$$

Hence proved.




Figure 2