






## 4.3 Chi-Square Test of Independence: What Is It?


**MBC 638**

**Data Analysis and Decision Making**















## Chi-Square Test for Independence

- **$\chi^2$  test for independence:** a procedure used to determine if two variables are related or are statistically independent

## Chi-Square Test for Independence

- **$\chi^2$  test for independence:** a procedure used to determine if two variables are related or are statistically independent
- Convention:
  - $H_0$ :
  - $H_a$ :

## Chi-Square Test for Independence

- **$\chi^2$  test for independence:** a procedure used to determine if two variables are related or are statistically independent
- Convention:
  - $H_0$ :
  - $H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent (i.e., there *is* a relationship).

## Chi-Square Test for Independence

- **$\chi^2$  test for independence:** a procedure used to determine if two variables are related or are statistically independent
- Convention:
  - $H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent (i.e., there is *no* relationship).
  - $H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent (i.e., there *is* a relationship).

## Chi-Square Test for Independence

- **$\chi^2$  test for independence:** a procedure used to determine if two variables are related or are statistically independent
- Convention:
  - $H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent (i.e., there is *no* relationship).
  - $H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent (i.e., there *is* a relationship).
- How it works:

## Chi-Square Test for Independence

- **$\chi^2$  test for independence:** a procedure used to determine if two variables are related or are statistically independent
- Convention:
  - $H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent (i.e., there is *no* relationship).
  - $H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent (i.e., there *is* a relationship).
- How it works:
  - Compares "observed" counts and "expected" counts or frequencies



## Chi-Square Test for Independence

- **$\chi^2$  test for independence:** a procedure used to determine if two variables are related or are statistically independent
- Convention:
  - $H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent (i.e., there is *no* relationship).
  - $H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent (i.e., there *is* a relationship).
- How it works:
  - Compares "observed" counts and "expected" counts or frequencies
  - Does not give kind (positive/negative) or intensity of relationship



## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

		Variable 2								
Variable 1									Totals	
									Row total	
	Totals		Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Row total
										Grand total

## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

		Variable 2							
Variable 1									Totals
									Row total
								Row total	
								Row total	
								Row total	
Totals		Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Grand total

- Put your data in a two-way table.

## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

		Variable 2						
Variable 1								Row total
								Row total
Totals		Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Grand total

- Put your data in a two-way table.
- Depending on the variables, use two or more columns or rows.

## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

*Example: Does day of week affect car sales?*

		Variable 2						
Variable 1								Row total
								Row total
Totals		Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Grand total

- Put your data in a two-way table.
- Depending on the variables, use two or more columns or rows.

## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

Example: Does day  
of week affect car  
sales?

		Variable 2							Totals
Variable 1	Customer lead resulting in:	Sale of car							
		Not selling a car							Row total
									Row total
									Grand total
		Totals	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	

- Put your data in a two-way table.
- Depending on the variables, use two or more columns or rows.

## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

Example: Does day  
of week affect car  
sales?

		Variable 2							Totals
Variable 1	Customer lead resulting in:	Sale of car	MON	TUES	WED	THUR	FRI	SAT	
		Not selling a car							Row total
									Row total
									Grand total
		Totals	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	

- Put your data in a two-way table.
- Depending on the variables, use two or more columns or rows.



## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

Example: Does day of week affect car sales?

		Variable 2							Totals
		MON	TUES	WED	THUR	FRI	SAT	SUN	
Variable 1 Customer lead resulting in:	Sale of car	Count data	Count data	Count data	Count data	Count data	Count data	Count data	Row total
	Not selling a car	Count data	Count data	Count data	Count data	Count data	Count data	Count data	Row total
	Totals	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Grand total

- Put your data in a two-way table.
- Depending on the variables, use two or more columns or rows.

## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

Example: Does day of week affect car sales?

		Variable 2							Totals
		MON	TUES	WED	THUR	FRI	SAT	SUN	
Variable 1 Customer lead resulting in:	Sale of car	Count data	Count data	Count data	Count data	Count data	Count data	Count data	Row total
	Not selling a car	Count data	Count data	Count data	Count data	Count data	Count data	Count data	Row total
	Totals	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Grand total

- Put your data in a two-way table.
- Depending on the variables, use two or more columns or rows.
- In each cell, count the frequency of simultaneous occurrence.

## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

Example: Does day  
of week affect car  
sales?

		Variable 2							Totals
		MON	TUES	WED	THUR	FRI	SAT	SUN	
Variable 1 Customer lead resulting in:	Sale of car	Count data	Count data	Count data	Count data	Count data	Count data	Count data	Row total
	Not selling a car	Count data	Count data	Count data	Count data	Count data	Count data	Count data	Row total
	Totals	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Grand total

- Put your data in a two-way table.
- Depending on the variables, use two or more columns or rows.
- In each cell, count the frequency of simultaneous occurrence.
  - Discrete data: need a lot of data (> 5 counts per cell)

## Setup for Analysis of Two-Way Tables

$H_0$ : Categorical Variable 1 and Categorical Variable 2 are independent.

$H_a$ : Categorical Variable 1 and Categorical Variable 2 are not independent.

Example: Does day  
of week affect car  
sales?

		Variable 2							Totals
		MON	TUES	WED	THUR	FRI	SAT	SUN	
Variable 1 Customer lead resulting in:	Sale of car	Count data	Count data	Count data	Count data	Count data	Count data	Count data	Row total
	Not selling a car	Count data	Count data	Count data	Count data	Count data	Count data	Count data	Row total
	Totals	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Col. total	Grand total

- Put your data in a two-way table.
- Depending on the variables, use two or more columns or rows.
- In each cell, count the frequency of simultaneous occurrence.
  - Discrete data: need a lot of data (> 5 counts per cell)
  - No exact sample size specified