

5.4 Confidence Interval Example

MBC 638

Data Analysis and Decision Making

Example: Hank's Process

Example: Hank's Process

- Find 95% confidence interval about the mean

Example: Hank's Process

- Find 95% confidence interval about the mean
 - Need lower and upper limits

Example: Hank's Process

- Find 95% confidence interval about the mean
 - Need lower and upper limits
- Use data given by example:

Example: Hank's Process

- Find 95% confidence interval about the mean
 - Need lower and upper limits
- Use data given by example:
 - $n = 30$

Example: Hank's Process

- Find 95% confidence interval about the mean
 - Need lower and upper limits
- Use data given by example:
 - $n = 30$
 - $\bar{x} = 17.23$

Example: Hank's Process

- Find 95% confidence interval about the mean
 - Need lower and upper limits
- Use data given by example:
 - $n = 30$
 - $\bar{x} = 17.23$
 - $s = 4.52$

Example: Hank's Process

- Find 95% confidence interval about the mean
 - Need lower and upper limits
- Use data given by example:
 - $n = 30$
 - $\bar{x} = 17.23$
 - $s = 4.52$
- Assume $\alpha = 0.05$

Confidence Interval Formulas for Mean

Boundaries of interval (two sided)	When population standard deviation is <i>known</i> (not often)	When population standard deviation is <i>unknown</i> and sample size n is <i>large</i> (≥ 30)	When population standard deviation is <i>unknown</i> and sample size n is <i>small</i> (< 30)
Upper confidence limits for μ	$U = \bar{x} + z^* \frac{\sigma}{\sqrt{n}}$	$U = \bar{x} + z^* \frac{s}{\sqrt{n}}$	$U = \bar{x} + t \frac{s}{\sqrt{n}}$
Lower confidence limits for μ	$L = \bar{x} - z^* \frac{\sigma}{\sqrt{n}}$	$L = \bar{x} - z^* \frac{s}{\sqrt{n}}$	$L = \bar{x} - t \frac{s}{\sqrt{n}}$
			$df = n - 1$

Confidence Interval Formulas for Mean

Boundaries of interval (two sided)	When population standard deviation is <i>known</i> (not often)	When population standard deviation is <i>unknown</i> and sample size n is <i>large</i> (≥ 30)	When population standard deviation is <i>unknown</i> and sample size n is <i>small</i> (< 30)
Upper confidence limits for μ	$U = \bar{x} + z^* \frac{\sigma}{\sqrt{n}}$	$U = \bar{x} + z^* \frac{s}{\sqrt{n}}$	$U = \bar{x} + t \frac{s}{\sqrt{n}}$
Lower confidence limits for μ	$L = \bar{x} - z^* \frac{\sigma}{\sqrt{n}}$	$L = \bar{x} - z^* \frac{s}{\sqrt{n}}$	$L = \bar{x} - t \frac{s}{\sqrt{n}}$
			$df = n - 1$

Confidence Interval for Hank's Process

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}}$$

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23 \pm (1.96)$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23 \pm (1.96) \frac{4.52}{\sqrt{30}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23 \pm (1.96) \frac{4.52}{\sqrt{30}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23 \pm (1.96) \frac{4.52}{\sqrt{30}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23 \pm (1.96) \frac{4.52}{\sqrt{30}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).

Confidence Interval for Hank's Process

$$U \& L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23 \pm (1.96) \frac{4.52}{\sqrt{30}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).

Confidence Interval for Hank's Process

$$U \text{ \& } L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23 \pm (1.96) \frac{4.52}{\sqrt{30}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).

$$15.61 \leq \mu \leq 18.85$$

Confidence Interval for Hank's Process


$$U \text{ \& } L = \bar{x} \pm z^* \frac{s}{\sqrt{n}} = 17.23 \pm (1.96) \frac{4.52}{\sqrt{30}}$$

- Use data given by example: $n = 30$; $\bar{x} = 17.23$; $s = 4.52$.
- Plus/minus gives margin of error above and below the parameter estimate.
- In Table D, find z^* at 95% confidence ($p = 0.95$).


$$15.61 \leq \mu \leq 18.85$$

- We are 95% confident that the mean of the population (μ_{pop}) is between 15.61 and 18.85 days.

Other Ways to Interpret This CI




Other Ways to Interpret This CI




- The above distribution is a distribution of sample means, so the probability of the true population mean being within these bounds is 95%.

Other Ways to Interpret This CI



- The above distribution is a distribution of sample means, so the probability of the true population mean being within these bounds is 95%.
- In 95% of all xbars (calculated from different samples of this process each having a different xbar) the interval (the shaded region) will include the population mean.
- **We are 95% confident that the population mean lies between this upper and lower bound.**
- Confidence intervals are NOT used to determine an interval at which any one observation of the distribution lies.

Other Ways to Interpret This CI



- The above distribution is a distribution of sample means, so the probability of the true population mean being within these bounds is 95%.
- In 95% of all xbars (calculated from different samples of this process each having a different xbar) the interval (the shaded region) will include the population mean.
- **We are 95% confident that the population mean lies between this upper and lower bound.**
- Confidence intervals are NOT used to determine an interval at which any one observation of the distribution lies.

In Excel:

=CONFIDENCE.NORM(

In Excel:

=CONFIDENCE.NORM(alpha,

In Excel:

=CONFIDENCE.NORM(alpha, std dev,

Confidence Intervals

Confidence Interval = Parameter estimate \pm [(Confidence) x (Variability)]

\bar{x}

Sample mean is a
parameter
estimate

margin of error (E)

Add and
subtract this
value from the
parameter
estimate

Use
Excel to
calculate
"E"

=CONFIDENCE.NORM(alpha, std dev, sample size)

Hank's example:
Mean=17.23 days

=CONFIDENCE.NORM(0.05, 4.52 ,30) = 1.617

Confidence interval = $17.23 \pm 1.617 = (15.61, 18.85)$