

--	--	--	--	--	--	--	--	--	--

**Fifth Semester B.E. Degree Examination, June/July 2023**  
**Mathematics for Machine Learning**

Time: 3 hrs.

Max. Marks: 100

*Note: Answer any FIVE full questions, choosing ONE full question from each module.*

**Module-1**

- 1 a. Define the linear dependent and linear independent of the vector space  $V(F)$ . Also show that the set of vectors  $(1 \ 0 \ 1)$ ,  $(1 \ 1 \ 0)$ ,  $(-1, \ 0, \ -1)$  is linearly dependent in  $V_3(\mathbb{R})$ . (06 Marks)
- b. Solve the system of equations and also show that the solution is unique.  
 $x_1 + x_2 + x_3 = 3$   
 $x_1 - x_2 + 2x_3 = 2$   
 $2x_1 + 3x_3 = 1$ . (06 Marks)
- c. For the matrix  $A = \begin{bmatrix} 1 & -1 & 2 \\ 3 & 1 & 0 \end{bmatrix}$ . Determine the linear transformation  
 $T : V_3(\mathbb{R}) \rightarrow V_2(\mathbb{R})$  relative to the basis  $B_1$  and  $B_2$  of  $V_3(\mathbb{R})$  are  $V_2(\mathbb{R})$ .  
 i)  $B_1 = \{(1 \ 1 \ 1) \ (1 \ 2 \ 3) \ (1 \ 0 \ 0)\}$   
 ii)  $B_2 = \{(1, \ 1) \ (1, \ -1)\}$  (08 Marks)

**OR**

- 2 a. Define:  
 i) An inner product space  
 ii) Projection of two vectors  $u$  and  $v$   
 iii) Orthogonal vectors  
 iv) An orthogonal set. (08 Marks)
- b. Solve by using the Gaussian elimination method  
 $2x_1 + x_2 + 4x_3 = 12$   
 $4x_1 + 11x_2 - x_3 = 33$   
 $8x_1 - 3x_2 + 2x_3 = 20$  (06 Marks)
- c. Obtain the matrix of linear transformation  $T : V_2(\mathbb{R}) \rightarrow V_3(\mathbb{R})$ , defined by  $T(x, y) = (x + y, x, 3x - y)$  with respect to the basis  $B_1$  and  $B_2$  where  $B_1 = \{(1, 1), (3, 1)\}$  and  $B_2 = \{(1, 1, 1), (1, 1, 0), (1, 0, 0)\}$ . (06 Marks)

**Module-2**

- 3 a. Show that the given vector form an orthogonal basis for  $\mathbb{R}^3$  also express  $\vec{0}$  as a linear combination of the basis vector, write the coordinate vector  $[W]_B$  of  $\vec{W}$  with respect to the basis  $B = \{\vec{V}_1, \vec{V}_2, \vec{V}_3\}$  of  $\mathbb{R}^3$  where  $V_1 = \begin{bmatrix} -1 \\ 0 \\ -1 \end{bmatrix}$   $V_2 = \begin{bmatrix} 3 \\ 6 \\ 3 \end{bmatrix}$   $V_3 = \begin{bmatrix} 3 \\ -3 \\ 3 \end{bmatrix}$   $W = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$ . (08 Marks)



18A156

- b. Reduce the matrix to diagonal form

$$A = \begin{bmatrix} 11 & -4 & -7 \\ 7 & -2 & -5 \\ 10 & -4 & -6 \end{bmatrix}$$

(06 Marks)

- c. If  $v = \begin{bmatrix} 7 \\ 6 \end{bmatrix}$  and  $u = \begin{bmatrix} 4 \\ 2 \end{bmatrix}$  then find the orthogonal projection of  $v$  on to  $u$  and the orthogonal set.

(06 Marks)

OR

- 4 a. Find the singular value decomposition [SVD] of the matrix  $A = \begin{bmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ . (10 Marks)
- b. Show that the Eigen values of the following matrix are all equal, and also find the corresponding eigen vector.

$$A = \begin{bmatrix} -3 & -7 & -5 \\ 2 & 4 & 3 \\ 1 & 2 & 2 \end{bmatrix}$$

(10 Marks)

### Module-3

- 5 a. A particle moves along the curve  $\vec{r} = t^2 \hat{i} - t^3 \hat{j} + t^4 \hat{k}$ , where 't' is the time. Find the magnitude of tangential component of its acceleration  $t = 1$ . (06 Marks)
- b. If  $U = x + y + z$ ,  $V = x^2 + y^2 + z^2$ ,  $W = xy + yz + zx$ , then prove that  $\text{grad } u$ ,  $\text{grad } v$ ,  $\text{grad } w$  are coplanar. (06 Marks)
- c. If  $f(x) = \sqrt{x^2 + \exp(x^2)} + \cos(x^2 + \exp(x^2))$  find  $\frac{df}{dx}$ . Using the following computation graph and the intermediate variables a, b, c, d where  $a = x^2$ ,  $b = \exp a$ ,  $c = a + b$ ,  $d = \sqrt{c}$ ,  $e = \cos c$ ,  $f = d + e$ .




Fig.Q.5(c)

(08 Marks)

OR

- 6 a. If the directional derivative of  $\phi = axy^2 + byz + cz^2x^3$  at  $(-1, 1, 2)$  has a maximum magnitude of 32 units in the direction of parallel to y-axis find a, b, c. (08 Marks)
- b. Define gradient of a vector valued function consider the function  $h : \mathbb{R} \rightarrow \mathbb{R}^2$   $h(t) = (f \circ g)(t)$   $f : \mathbb{R}^2 \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}^2$ , if  $f(x) = \exp(x_1 x_2^2)$   $x = \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = g(t) = \begin{bmatrix} t \cos t \\ t \sin t \end{bmatrix}$  then compute gradient of  $h$  with respect to  $t$ . (12 Marks)



18AI56

#### Module-4

- 7 a. State and prove Baye's theorem on conditional probability. (08 Marks)  
 b. Let A and B be two events, which are not mutually exclusive and are connected with random experiment. Given that  $P(A) = 3/4$   $P(B) = 1/5$   $P(A \cap B) = 1/20$  then find: i)  $P(A \cup B)$  ii)  $P(A \cap \bar{B})$  iii)  $P(\bar{A} \cap B)$  iv)  $P(A/B)$  and  $P(B/A)$ . (06 Marks)  
 c. A random variable x has the following probability distribution:

x	0	1	2	3	4	5	6	7
P(x)	0	K	2K	2K	3K	K <sup>2</sup>	2K <sup>2</sup>	7K <sup>2</sup> + K

Find : i) Value of K ii)  $P(x < 6)$  iii)  $P(x \geq 6)$ . (06 Marks)

OR

- 8 a. Test whether the following function is a density function  $f(x) = \begin{cases} e^{-x}, & x \geq 0 \\ 0, & x < 0 \end{cases}$  if so determine the probability that the variate having its density function will fall in the interval (1, 2). (08 Marks)  
 b. The length of the telephone conversation in a booth has been an exponential distribution and found on an average to be 5 minutes. Find the probability that a random call made from this booth i) Ends in less than 5 minutes ii) Between 5 and 10 minutes. (06 Marks)  
 c. Define binomial distribution and find the binomial probability distribution which has mean 2 and variance 4/3. (06 Marks)

#### Module-5

- 9 a. By using gradient descent method (steepest method) for  $f(x_1, x_2) = x_1 - x_2 + 2x_1^2 + 2x_1x_2$  has the optimal solution starting from the point (0, 0) carry out four iterations. (12 Marks)  
 b. Use Lagrange's multiplier, find the dimension of the rectangular box, which is open at the top of maximum capacity whose volume is 32 cubic feet. (08 Marks)

OR

- 10 a. Given that  $x + y + z = a$  where 'a' is a constant, find the extreme value of the function  $f(x, y, z) = x^m y^n z^p$ . (08 Marks)  
 b. Define a convex and a concave function, test the nature of definiteness by checking its extreme values for the function  $f(x_1, x_2) = x_1^3 + x_2^3 + 2x_1^2 + 4x_2^2 + 6$ . (06 Marks)  
 c. Determine whether the function  $f(x) = x \log_2 x$  is convex or not for  $x > 0$ . (06 Marks)

\* \* \* \* \*