

Assignment 2

Exercises

P1 $a, b, c \in \mathbb{Q}$ and $a \neq 0$

$$ax = b \rightarrow x = ba^{-1}$$

$$axa^{-1} = ba^{-1}$$

$$aa^{-1}x = ba^{-1}$$

$$1x = ba^{-1}$$

$$x = ba^{-1}$$

property in slide ($xz = yz \rightarrow x = y, z \neq 0$)

commutativity

identity

In order to prove the uniqueness we will proceed by contradiction.

Assume there are two different solutions, x and y .

$$ax = b \text{ and } ay = b \text{ therefore } ax = ay$$

Now using the field axioms and properties we can prove it.

$$axa^{-1} = aya^{-1}$$

$$aa^{-1}x = aa^{-1}y$$

$$1x = 1y$$

$$x = y$$

property in slide ($xz = yz \rightarrow x = y, z \neq 0$)

commutativity

identity

We have just showed that the solution is unique.

P2 For all $x, y \in \mathbb{Q}$ if $0 < x < y$ then $x^2 < y^2$.

Knowing that $0 < x < y$ we can use the order axiom number 5 (O5).

Starting from $x < y$ we can multiply both members by x one time and a second one by y .

$$xx < yx \text{ and } xy < yy \text{ by O5 therefore}$$

$$xx < yx = xy < yy \text{ commutativity}$$

Since

$$x^2 = xx < yy = y^2$$

we proved

$$x^2 < y^2$$

P3 Base case $n=1$.

$$|x_1| \leq |x_1| \text{ is true.}$$

Let n be an arbitrary element of \mathbb{N} , assume $P(n)$ which is

$$|x_1 + x_2 + \dots + x_n| \leq |x_1| + |x_2| + \dots + |x_n|$$

and prove $P(n+1)$.

$P(n+1)$ is

$$|x_1 + x_2 + \dots + x_n + x_{n+1}| \leq |x_1| + |x_2| + \dots + |x_n| + |x_{n+1}|$$

We can write $P(n+1)$ as $|x_1 + x_2 + \dots + x_n + x_{n+1}| = |(x_1 + x_2 + \dots + x_n) + x_{n+1}|$

Now we are going to prove that $|x+y| \leq |x| + |y|$.

Since $x \leq |x|, y \leq |y|$ and $-x \leq |x|, -y \leq |y|$ we can say $x+y \leq |x| + |y|$ and

$-(x+y) = -x-y \leq |x|+|y|$ which means $|x+y| \leq |x|+|y|$.
Therefore if we call $x = (x_1 + x_2 + \dots + x_n)$ and $y = x_{n+1}$ then

$$|(x_1 + x_2 + \dots + x_n) + x_{n+1}| = |x+y|$$

therefore

$$|x+y| \leq |x|+|y|$$

which is

$$|(x_1 + x_2 + \dots + x_n) + x_{n+1}| \leq |(x_1 + x_2 + \dots + x_n)| + |x_{n+1}|$$

Using the induction hypothesis

$$|(x_1 + x_2 + \dots + x_n)| + |x_{n+1}| \leq |x_1| + |x_2| + \dots + |x_n| + |x_{n+1}|.$$

We have just showed that $\forall n \in \mathbb{N} P(n)$ holds.

P4 Bonus Exercise

Given $a, b, c \in \mathbb{R}$ which satisfy the equation

$$a+b=c.$$

Is also true that $\frac{1}{a} + \frac{1}{b} = \frac{1}{c}$?

$$a, b, c \neq 0$$

$$\frac{1}{a} + \frac{1}{b} = \frac{1}{c} \Rightarrow$$

$$\Rightarrow \frac{b+a}{ab} = \frac{1}{a+b} \Rightarrow$$

$$a+b=c$$

$$\Rightarrow \frac{(a+b)^2}{ab} = 1 \Rightarrow$$

$$a, b, c \neq 0$$

$$\Rightarrow (a+b)^2 = ab \text{ which is } (a+b)^2 - ab = 0 \Rightarrow$$

$$\Rightarrow a^2 + ab + b^2 = 0$$

$$\Delta = \sqrt{b^2 - 4b^2} = \sqrt{-3b^2}$$

knowing that $\forall x \in \mathbb{R}, x \geq 0$ and $x^2 = 0 \Leftrightarrow x = 0$

Since $\Delta = \sqrt{-3b^2}$ and $-3b^2$ is a negative number the equation has no solutions.