
A. Nucleotide Diversity Across Populations (Single-Copy Partition)
Nucleotide Diversity $\hat{\pi}$ (10^{-3} / bp)
0
2
4
6
8
10
Japonica
Indica
Indica_Canonical
Spontanea
Northern_Margin_Russia
East_Asia
South_Southeast_Asia


B. Tajima's D Neutrality Statistic Across Cohorts
Tajima's D
3
2
1
0
−1
−2
Significance Threshold (|D| > 1.8)
Japonica
Indica
Indica_Canonical
Spontanea
Northern_Margin_Russia
East_Asia
South_Southeast_Asia


C. Diversity Partitioned by Quadripartite Compartments (Canonical)
Population
Japonica
Indica_Canonical
All_Sativa_Canonical
Nucleotide Diversity $\hat{\pi}$ (10^{-3} / bp)
0.0
0.1
0.2
0.3
0.4
0.5
LSC
IRb
SSC
IRa
Quadripartite Domain


D. Functional Constraint: Protein-Coding (CDS) vs Intergenic (IGS)
Population
Japonica
Indica_Canonical
All_Sativa_Canonical
Nucleotide Diversity $\hat{\pi}$ (10^{-3} / bp)
0.0
0.2
0.4
0.6
0.8
1.0
Protein_Coding_CDS
Intergenic_Spacers_IGS
Functional Category