

A
Standard Mantel: r_M = 0.783 (95% CI [0.53, 0.94], exact p = 0.039)
Spearman Rank: ρ_M = 0.832 (exact p = 0.014)
Partial Mantel (| Genome Group): r_M = 0.778 (exact p = 0.061)
Ambon × Mauli
Ambon × Mas
Ambon × Kalimbang
Ambon × Awak
Ambon × Talas
Awak × Mas
Awak × Mauli
Kalimbang × Mauli
Awak × Kalimbang
Mauli × Talas
Mas × Talas
Kalimbang × Talas
Awak × Talas
Mas × Mauli
Kalimbang × Mas
Pairwise Morphological Euclidean Distance (Z-Standardized, VIF < 5)
Pairwise ITS Maximum Likelihood Genetic Distance (Substitutions / Site)
Parental Cross Architecture
Interspecific (Acuminata AA/AAA x Balbisiana ABB)
Intra-Acuminata Diploid Cross (AA x AA/AAB)
Intra-Acuminata Ploidy Bridge (Diploid AA x Triploid AAA)
B
Pedicel / Stalk Width (cm) r = +0.81 (p = 0.06)
Fruit Length (cm) r = +0.53 (p = 0.15)
Fruit Width (cm) r = +0.49 (p = 0.07)
Fruit Diameter With Peel (cm) r = +0.31 (p = 0.25)
Pulp Diameter Without Peel (cm) r = +0.24 (p = 0.44)
Fruits Per Hand Class (IPGRI) r = +0.17 (p = 0.43)
Pedicel Surface Hairiness (IPGRI) r = +0.07 (p = 0.50)
Longitudinal Fruit Curvature (IPGRI) r = -0.05 (p = 1.00)
Pedicel / Stalk Length (cm) r = -0.14 (p = 0.65)
Fruit Apex / Tip Architecture (IPGRI) r = -0.14 (p = 0.75)
Pericarp / Peel Thickness (cm) r = -0.17 (p = 0.55)
Fruit Rachis Posture (IPGRI) r = -0.35 (p = 0.40)
Single-Trait Mantel Correlation with ITS Genetic Distance (r_M)
Morphological Character Class
Continuous (Collinear, VIF >= 5)
Continuous (Retained, VIF < 5)
Qualitative IPGRI Descriptor (Gower)