

Week 2

Graph Theory And Application*

2.1 Connection in Graphs

2.1.1 Walk

A Walk in a graph is a sequence of edges such that each edge (except the first one) starts with a vertex where the previous edges ended. i.e The length of a walk is number of edges in it.

Path- A path is a walk where all edges are distinct(no repeated edges).

Simple Path- A simple path is a path where all vertices are distinct

Cycle- A cycle is a path in which first and the last vertices are same.

Trail- A trail is a walk with no repeated vertex.

2.1.2 Connected and Disconnected Graph

A graph is **connected** if there is path between any two pair of vertices in that graph. Otherwise it is **disconnected**.

Connected component of Graph G is maximal connected subgraphs of G. (in other words, those connected subgraphs which are not contained in larger connected subgraphs of G.)

Connected component of Graph G is maximal connected subgraphs of G.

*Lecturer: Anand Misra. Scribe: Satya Prakash Sharma.

2.1.3 Union of graphs

The union of graphs G_1, G_2, \dots, G_k written as $\cup_{i=1}^k G(i)$ is the graph with vertex set and edge set $\cup_{i=1}^k V(G_i)$

Proposition

- **Every graph with n vertices and k edges has at least $n-k$ components.** - Since k edges can connect maximum of $k+1$ nodes and form 1 component, so here we have now $n-(k+1)$ nodes remaining which form $n-(k+1)$ components, so in total we have $1+n-(k+1)$ i.e. $n-k$. Therefore, Every graph with n vertices and k edges has at least $n-k$ components

2.1.4 Cut Vertex and Cut Edge

Cut Vertex- A cut-Vertex of a graph is a vertex whose deletion increases the number of components.

Cut Edge- A cut-edge of a graph is an edge whose deletion increases the number of components.

2.1.5 Theorem

- **An edge is a cut-edge if and only if it belongs to no cycle.** Take any edge $e = u, v$. Remove this edge from our graph: if the graph is still connected, then there is some path from u to v not involving e ; consequently, if we add e to the end of this path, we get a cycle. Thus, if e is not a cut-edge, it's involved in a cycle.

Conversely: suppose that $e = u, v$ lies in a cycle. Let P be the path from u to v that doesn't use e (i.e. go the other way around the cycle.) Pick any x, y in G ; because G is connected, there's a path from x to y in G . Take this path, and edit it as follows: whenever the edge e shows up, replace this with the path P (or P traced backwards, as needed.) This then creates a walk from x to y ; by deleting cycles, this walk will always become a path, and thus G is connected. So if e is involved in a cycle, it's not a cut-edge.

- **Every closed odd walk contains an odd cycle.** Let l be closed odd walk, For $l=1$, obviously true. (self loop). *Suppose it is true for $l < L$, then we have to show that this is also true for L .*

Case1: If there is no repetition of vertex in walk, then a closed walk = a closed cycle.

Case2: If there is repetition in vertex in the walk, and let us suppose v is vertex which repeats. Break the walk into two v - v walks (say w_1 and w_2). Since $l = w_1 + w_2$ is odd, that means either w_1 or w_2 is odd walk. And surely they both are less than L . From the induction one of them (odd walk one) contains an odd cycle.

- **A closed even walk need not contain a cycle.** Example: If I go to jodhpur from my home and return back to my home from jodhpur with the same route and it will count as a closed even walk, but it is not a cycle because *cycle is a path in which first and the last vertices are same, and path has not repeated edges*

2.2 Bipartite Graph

A **bipartition** of a graph is a specification of two disjoint independent sets in G whose union is $V(G)$. The statement “ G is a bipartite graph with bipartition X and Y ” specifies one such partition.

2.2.1 Proposition

- **A simple path is a bipartite graph.-** We can build a bipartite graph by have first node in 1 set and next to first in other set and do same for next to next as well.
- **A C_n is bipartite iff n is even.-** If n is even then then We can build the bipartite graph using first logic(A simple path). Similary, If C_n is bipartite graph then it must have n ==even if n is odd then we can not put them in two set.

2.2.2 Theorem

- **A graph is bipartite iff it has no odd cycle.**

Necessary condition: Assume graph is bipartite and X and Y are two independent sets. To have a cycle, one has to traverse X to Y to X or Y to X to Y one or more time. Therefore, a bipartite graph can't have odd cycle.

Sufficient condition: If G has no odd cycle \Rightarrow it does not contain a cycle OR it contain even cycle.

- If does not contain a cycle : take one vertex in X and next vertex in Y .
- If it contain even cycle : Partition the graph such that each even length cycle is one subgraph. We know C_n is bipartite for even length cycle.

2.3 Eulerian Circuit

Eulerian Graph- A graph is Eulerian if it has a closed trail containing all the edges. We call a closed trail a circuit when we do not specify the first vertex but keep the list in cyclic order.

Eulerian Circuit- An Eulerian circuit in a graph is a circuit containing all the edges.

Lemma- If every vertex of a graph has degree at least 2, then G has cycle.

2.3.1 Theorem


- A graph G is Eulerian iff it has at most one non-trivial component and all its vertices have even degree.

Necessary condition: Assume a graph G is Eulerian.

- It has closed trail containing all the edges.
- There will be an incoming and outgoing edge for all the vertices.
- All its vertices have even degree. (It has to have one non-trivial component, as more than one components closed trail might not be possible).

Sufficient condition: Assume a graph G has at most one non-trivial component and all its vertices have even degree.

- If number of edges $m = 0$ then G is Eulerian.
- Assume this is true for all graphs having less than M edges and obviously above properties. Then
- Each vertex has at least two degree.
- G contains cycle (According to above Lemma). say C . Remove $E(C)$ from G to construct G' .
- G' has less than M edges and Each vertex of G' has even degree.
- It can have more than one component.
- By the induction hypo. Each component of G' contain Eulerian Cycle.
- We combine these Eulerian cycles with C to construct an Eulerian circuit as follows: Traverse C until a component of G' 's appear, then traverse Eulerian cycle of that component, come back to C and repeat this.



2.4 Vertex Degree and Counting

Degree of a vertex- Degree of a vertex v in a graph G written as $d(v)$ is number of edges incidents to v , except that each loop at v counts twice. The maximum and minimum number of $\delta(G)$ and $\Delta(G)$ degrees are denoted by respectively.

Order of Graph The order of a graph G , $n(G)$ is number of vertices in G .

Size of Graph The size of a graph G , $e(G)$ is number of edges in G .

- The order and size of complete graph K_n : nC_2
- The order and size of complete bipartite graph $K_{m,n}$: $m*n$

2.4.1 Handshaking Lemma

If G is a graph then

$$\sum_{v \in V(G)} d(v) = 2e(G)$$

Let $G = (V, E)$ be a graph and let C be a connected component of G . Place one coin on each node in C for each edge in E incident to it.

Notice that the number of coins on any node v is equal to $\deg(v)$.

We claim that there are an even total coins distributed across all the nodes of G .

Notice that each edge contributes two coins to the total, one for each of its endpoints. This means that there are $2m$ total coins distributed across the nodes of V , where m is the number of edges adjacent to nodes in C , and $2m$ is even. Since there are an even number of coins distributed across the nodes, our earlier theorem tells us that the number of nodes in G with an odd number of coins on them must be even. The number of coins on each node is the degree of that node, and therefore there must be an even number of nodes of odd degree.