

Solution of Non-homogeneous Differential equations

A linear ordinary differential equation of order n , is written as:

$$a_0 y^n + a_1 y^{n-1} + \dots + a_{n-1} y' + a_n y = r(x)$$

$$(a_0 D^n + a_1 D^{n-1} + \dots + a_{n-1} D' + a_n) y = r(x)$$

$$F(D)y = r(x)$$

$$y_p = \frac{1}{F(D)} r(x)$$

Case 1: $r(x) = e^{ax}$, For P.I. Put $D = a$

$$1. \quad y'' - 2y' - 3y = 3e^{2x}$$

$$2. \quad y''' - 2y'' - 5y' + 6y = 4e^{-x} - e^{2x}$$

$$3. \quad y'' + y' - 6y = 5e^{-3x}$$

$$y'' - 4y' + y = e^{\frac{x}{2}}$$

Case 2: $r(x) = \cos ax$ or $\sin ax$ For P.I. Put $D^2 = -a^2$

$$y'' + 4y = 6 \cos x$$

$$2y'' + y' - y = 16 \cos 2x$$

$$y'' - 5y' + 4y = 65 \sin 2x$$

$$y''' - y'' + 4y' - 4y = \sin 3x$$

$$y'' + y = 6 \sin x$$

Case 3: $r(x) = x^n$

$$y'' + 5y' + 4y = 3x + 2$$

$$y'' + 16y = 64x^2$$

$$y^{iv} + 3y'' = x^2$$

Case 4: $r(x) = e^{ax}h(x)$

$$y'' + 2y' + 5y = e^{-x} \cos 2x$$

$$y'' - y = 6xe^x$$

$$2y'' + 7y' - 4y = xe^{-4x}$$

Case 5: $r(x) = x^n \sin ax$ or $x^n \cos ax$

$$y' + y = x \cos x$$

$$y'' + 4y' + 3y = x \sin 2x$$

$$y'' + 3y' + 2y = x e^x \sin x$$

Solution of Euler-Cauchy equations

$$2x^2y'' + 3xy' - 3y = x^3$$

$$x^2y'' + 5xy' + 3y = \ln x$$

$$x^3y'' + 5x^2y' + 5xy' + y = x^2 + \ln x$$

$$x^2y'' + 5xy' - 5y = 24x \ln x$$

$$4x^2y'' + y = \log x, \quad y(1) = 0, y(e) = 5$$

$$x^2y'' - xy' + 2y = 6, y(1) = 1, y'(1) = 2$$

$$2x^2y'' + 3xy' - y = x, y(1) = 1, y(4) = \frac{41}{16}$$

$$x^2y'' - 3xy' + 3y = 5x^2 - x, y(1) = 1, y'(1) = \frac{3}{2}$$

Method of Undetermined Coefficients:

| | |
|----------------|------------------------------------|
|r(x)..... |Initial Guess for Y_p |
|----------------|------------------------------------|

| P(x) | Q(x) |
|---|---|
| $c e^{ax}$ | $d e^{ax}$ |
| $\alpha \cos(bx) \text{ or } \beta \sin(bx)$ | $A \cos(bx) \text{ or } B \sin(bx)$ |
| $P(x) \cos(bx) \text{ or } P(x) \sin(bx)$ | $Q(x) \cos(bx) \text{ or } R(x) \sin(bx)$ |
| $P(x) e^{ax} \cos(bx) \text{ or } P(x) e^{ax} \sin(bx)$ | $Q(x) e^{ax} \cos(bx) \text{ or } R(x) e^{ax} \sin(bx)$ |
| | |
| | |

Note: whenever the term(or terms) of C.F. and P.I. is (are) same in that case we have to multiply the P.I.'s term with x.

$$y'' - 3y' - 10y = x$$

$$2y'' - y' - 3y = x^2 + 1$$

$$4y'' - y = e^x$$

$$4y'' - y = e^x + e^{3x}$$

$$y'' - y = e^x$$

$$y'' + y = \sin 2x$$

$$y'' + y = \sin x$$

$$y'' + y = e^x \sin x$$

Method of variation of Parameters

Consider the second order non-homogeneous linear equation

$$a_0(x)y'' + a_1(x)y' + a_2(x)y = r(x), \quad a_0(x) \neq 0$$

$$C.F = y = C_1y_1(x) + C_2y_2(x)$$

$$y(x) = A(x)y_1(x) + B(x)y_2(x)$$

$$A(x) = - \int \frac{g(x)y_2(x)}{W(x)} dx + c_1$$

$$B(x) = \int \frac{g(x)y_1(x)}{W(x)} dx + c_2$$

$$g(x) = \frac{r(x)}{a_0(x)}$$

$$W = \begin{vmatrix} y_1 & y_2 \\ y_1' & y_2' \end{vmatrix}$$

$$y'' + 3y' + 2y = 2e^x$$

$$y'' + 16y = 32 \sec 2x$$

$$x^2y'' + xy' - y = x$$

$$y'' + 4y' + 4y = e^{-2x} \sin x$$

$$y'' + 2y' + 2y = e^{-x} \cos x$$

$$y'' + 6y' + 9y = \frac{e^{-3x}}{x}$$

$$x^2 y'' + xy' - 4y = x^2 \ln x, \quad y_1 = x^2, \quad y_2 = \frac{1}{x^2}$$

Solution of systems of equations

$$y_1' = 2y_1 + y_2, \quad y_2' = y_1 + 2y_2$$

$$y_1' = y_1 + y_2, \quad y_2' = 9y_1 + y_2$$

$$y_1' = y_2, \quad y_2' = -9y_1$$

$$y_1' = 2y_1 + y_2, \quad y_2' = -18y_1 - 7y_2$$

$$y_1' + y_2 = 4 \sin t, \quad y_2' + y_1 = 8 \cos t$$

$$y_1' + y_1 + 3y_2 = 4e^{-t}, \quad y_2' + 4y_1 - 3y_2 = 8t$$

$$y_1' + 3y_1 + y_2 = 6e^t, \quad y_2' - 5y_1 - 3y_2 = 3e^{-t}$$

$$y_1' + 4y_1 - 5y_2 = 16 \sin t, \quad y_2' + 5y_1 - 4y_2 = e^t$$

$$y_1' + y_1 - 3y_2 = 6e^{-t}, \quad y_2' + 2y_1 - 14y_2 = 12e^t$$

$$\begin{aligned}(D - 2)y_1 + (D + 3)y_2 &= t + 4, \\ (3D + 5)y_1 + (2D + 6)y_2 &= 5t + 3\end{aligned}$$