

Normal ODE, Dependent –Independent functions , Homogeneous Linear ODE:

- 1.) The differential equation $y'' + 3y' + \sqrt{x}y = \sin x$ is normal on every subinterval of
- a) $(-\infty, \infty)$ b) $[0, \infty)$ c) $(0, \infty)$ d) $(-\infty, 1), (-1, 1), (1, \infty)$
- 2.) The differential equation $(x^2 - 1)y'' + 2xy' + y = x \ln x$ is normal on every subinterval of
- a) $(-\infty, \infty)$ b) $(0, \infty)$ c) $(-\infty, 2)$ d) $(0, 1), (1, \infty)$
- 3.) Which of the following functions are linearly independent for $x \in (0, \infty)$?
- a) $1, x, x^2, 1+x$ b) $1, x(1-x), x^2, x$ c) $2x, 6x+3, 3x+2$ d) $1, x, x^2, x^2(1-x)$
- 4.) If $y_1(x)$ and $y_2(x)$ be the linearly independent solutions of the equation $y'' + a(x)y' + b(x)y = 0$ on an interval I then which of the following is true
- a) both $y_1(x)$ and $y_2(x)$ vanishes for some $x_0 \in I$
- b) both $y_1(x)$ and $y_2(x)$ take extreme values for some $x_0 \in I$
- c) both $y_1(x)$ and $y_2(x)$ can not vanishes for some $x_0 \in I$ d) None of these
- 5.) The general solution of the differential equation $y'' + 2\pi y' + \pi^2 y = 0$ is
- a) $(A + Bx)e^{-\pi x}$ b) $(A + Bx)e^{-x}$ c) $(A + B)e^{-\pi x}$ d) $(A + Bx)e^{\pi x}$
- 6.) The differential equation whose linearly independent solutions are e^{2x}, xe^{2x} is ?
- a) $y'' + 4y' + 4y = 0$ b) $y'' + 4y' - 4y = 0$ c) $y'' - 4y' + 4y = 0$ d) $y'' - 4y' - 4y = 0$
- 7.) The lowest possible order of homogeneous linear differential equation whose particular solution is $3\cos 2x + 5\sinh 3x$ is
- a) 2 b) 3 c) 5 d) 4
- 8.) The roots of characteristic equation of differential equation $y^{iv} + 8y'' - 9y = 0$
- a) $\pm 1, \pm 3i$ b) $\pm i, \pm 3i$ c) $\pm i, \pm 3$ d) $\pm 1, \pm 3$
- 9.) The general solution of differential equation $y''' - 2y'' + y' = 0$ is
- a) $Ae^x + (Bx + C)$ b) $A + (Bx + C)e^x$ c) $(Bx + C)e^x$ d) $A + (Bx + C)e^{-x}$
- 10.) The lowest possible order of homogeneous linear differential equation whose particular solution is $1 + x + e^x - 3e^{3x}$ is
- a) 4 b) 3 c) 2 d) 5

(Operator method) Exp(ax), cosh(ax) sinh(ax), h(x). exp(ax): : Polynomila , Sin(ax) , cos(ax):

1. The particular integral $\frac{1}{D+3} e^{2x}$ is
(a) $\frac{1}{5} e^{2x}$ (b) $\frac{1}{5}$ (c) $\frac{1}{3} e^{2x}$ (d) $\frac{1}{2x+3} e^{2x}$
2. The particular integral $\frac{1}{D^2-9} e^{3x}$ is
(a) $\frac{1}{6} e^{3x}$ (b) $\frac{x e^{3x}}{6}$ (c) $\frac{x}{3} e^{3x}$ (d) doesn't exist
3. The particular integral $\frac{1}{f(D)} e^{ax} g(x)$ is
(a) $e^{ax} \frac{1}{f(D)} g(x)$ (b) $g(x) \frac{1}{f(D)} e^{ax}$ (c) $e^{ax} \frac{1}{f(D+a)} g(x)$ (d) $\frac{1}{f(a)} e^{ax} g(x)$
4. The particular integral $\frac{1}{f(D^2)} e^{ax}$ is
(a) $\frac{1}{f(-a^2)} e^{ax}$ (c) $\frac{1}{f(a^2)} e^{ax}$, provided $f(a^2) = 0$
(b) $\frac{1}{f(a^2)} e^{ax}$, provided $f(a^2) \neq 0$ (d) $\frac{1}{f(a)} e^{ax}$, provided $f(a) \neq 0$
5. The particular integral $\frac{1}{D^3-D^2+4D-4} \sin 3x$ is
(a) $-\frac{1}{5} \sin 3x$ (c) $\frac{1}{9} x \cos 3x$
(b) $\frac{1}{50} (\sin 3x + x \cos 3x)$ (d) $\frac{1}{50} (\sin 3x + 3 \cos 3x)$
6. The particular integral of the differential equation $y'' + y = 6 \sin x$ is
(a) $6 \cos x$ (b) $3x \sin x$ (c) $-3x \cos x$ (d) $6 x \cos x$
7. The particular integral $\frac{1}{D+5} (2016)^x$ is
(a) $\frac{1}{2021} (2016)^x$ (b) $x (2016)^x$ (c) $\frac{1}{\ln 2016} (2016)^x$ (d) $\frac{1}{(\ln 2016)+5} (2016)^x$
8. The particular integral of $\frac{d^2 y}{dx^2} + \frac{dy}{dx} = x^2 + 2x + 4$ is
(a) $\frac{x^2}{3} + 4x$ (b) $\frac{x^3}{3} + 4x$ (c) $\frac{x^3}{3} + 4$ (d) $\frac{x^2}{3} + 4x^2$
9. The particular integral of $\frac{d^4 y}{dx^4} - 16 \frac{d^2 y}{dx^2} = (8x + 16)$ is
(a) $-\frac{x^2}{12} - \frac{x}{2}$ (b) $\frac{x^3}{6} + 2x$ (c) $\frac{x^3}{3} + 4$ (d) $\frac{x^2}{3} + 4x$
10. The particular integral $\frac{1}{(D+1)^3} (2x + 4)$ is
(a) $\frac{x^2}{3} + 4x$ (b) $2x + 4$ (c) $x - 2$ (d) $2x - 2$
11. The particular integral of the differential equation $y'' + 4y = \sin x \cos x$ is
(a) $6 \cos x$ (b) $3x \sin x \cos x$ (c) $-3x \cos x$ (d) $-\frac{x}{8} \cos 2x$
12. The particular integral $\frac{1}{D^2} \cos 2x$ is
(a) $-4 \cos 2x$ (b) $-4 \sin 2x$ (c) $\frac{1}{2} \sin 2x$ (d) $-\frac{1}{4} \cos 2x$
13. The particular integral $\frac{1}{D-1} \cos 2x$ is
(a) $-\frac{1}{5} (\cos 2x - 2 \sin 2x)$ (b) $\frac{1}{2} \cos 2x$ (c) $-\sin 2x$ (d) $-\frac{1}{5} (\sin 2x + 2 \cos 2x)$

Method of Variation of Parameter, Method of Undetermined co-efficient:

1 The value of parameter $A(x)$ for LDE $y'' - 2y' - 3y = e^x$ using method of variation of parameters when $y_1 = e^{-x}$ and $y_2 = e^{3x}$ is

- (a) $-\frac{e^{2x}}{8} + c$ (b) $\frac{3}{2}e^x + c$ (c) $\frac{2}{3}e^{3x} + c$ (d) $-\frac{e^{5x}}{18} + c$

2 Solving by variation of parameter for the equation $y'' + y = \sec x$, the value of Wronskian

- a. 1 b. 2 c. 3 d. 4

3 Solving by variation of parameters for the equation $y'' - 4y' + 3y = e^x$, $x \neq 0$ the value of Wronskian is

- (a) $2e^x$ (b) $3e^{4x}$ (c) $2e^{4x}$ (d) $4e^x$

4 If by the method of variation of parameter $y(x) = A(x)\sin x + B(x)\cos x$ is the general solution of $y'' + y = \sec x$ then $B(x)$ is

- (A) $\ln|\sin x| + c$, (B) $\ln|\cos x| + c$, (C) $X + c$, (D) $\ln|x| + c$

5 The choice of particular integral for the equation $y'' - 9y = 13e^{3x}$ is

- (a) ce^{3x} (b) cxe^{3x} (c) cx^2e^{3x} (d) none of these

6 By the method of undetermined coefficients the choice of particular integral of the ODE

$$y'' + 4y' + 4y = 12e^{-2x}$$

- (A) ax^2e^{2x} , (B) ax^2e^{-2x} , (C) $12ae^{2x}$, (D) $12axe^{-2x}$

7 By the method of undetermined coefficients the trial solution for y_p for the differential equation $y'' + 3y' + 2y = 12x^2$ is of the form

- (A) $a + bx + cx^2$ (B) $a + bx$ (C) $ax + bx^2 + cx^3$ (D) None of these

8 By the method of undetermined coefficients the trial solution for y_p for the differential equation $y'' + 2y' + y = 6e^{-x}$ is of the form

- (a) Ae^{-x} (b) Bxe^{-x} (c) Cx^2e^{-x} (d) None of these

Q19. By the method of variation of parameter if $A(x)\cos x + B(x)\sin x$ be the particular integral of the differential equation

$$y'' + y = \sec x \text{ then } A(x) \text{ is}$$

- (a) $-\log|\cos x|$ (b) $\log|\cos x|$ (c) $\log|\sin x|$ (d) $-\log|\sin x|$

Q20. By the method of variation of parameter if $A(x)\cos 2x + B(x)\sin 2x$ be the particular integral of the differential equation

$$y'' + 4y = \sec 2x \text{ then } B(x) \text{ is}$$

- (a) $3x/2$ (b) $x^2/2$ (c) $-x/2$ (d) $x/2$

Q21. By method of undetermined coefficients the assumed particular integral of the differential equation

$$y'' + 9y = \cos 3x \text{ is}$$

- (a) $a\cos 3x + b\sin 3x$ (b) $x(a\cos 3x + b\sin 3x)$ (c) $x^2(A\cos 3x + b\sin 3x)$ (d) None of these

Q22. By method of undetermined coefficients the assumed particular integral of the differential equation

$$y'' + y = \sin x \text{ is}$$

- (a) $a\cos x + b\sin x$ (b) $x(a\cos x)$ (c) $x^2(A\cos 3x + b\sin 3x)$ (d) $x(a\cos x + b\sin x)$

Q23. By method of undetermined coefficients the assumed particular integral of the differential equation

$$y'' - 2y' - 3y = 6e^{-x} - 8e^x \text{ is}$$

- (a) $a e^{-x} + b e^x$ (b) $a x e^{-x} + b e^x$ (c) $a e^{-x} + b x e^x$ (d) $a x^2 e^{-x} + b e^x$

Q25. By method of undetermined coefficients the assumed particular integral of the differential equation

$$y'' - y = \sin x \text{ is}$$

- (a) $a \cos x + b \sin x$ (b) $x(a \sin x)$ (c) $x^2(A \cos x + b \sin x)$ (d) $x(a \cos x + b \sin x)$

Q26. By method of undetermined coefficients the assumed particular integral of the differential equation

$$y'' - 4y' + 13y = 12e^{2x} \sin 3x \text{ is}$$

- (a) $a \cos 3x + b \sin 3x$ (b) $x e^{2x}(a \sin 3x + b \cos 3x)$
(c) $x^2(A \cos 3x + b \sin 3x)$ (d) $x^2 e^{2x}(a \cos x + b \sin x)$

Euler-Cauchy Equation , Simultaneous ODE

Q28. If $D = \frac{d}{dx}$, then $\frac{1}{x^2 D^2 + 2} 16x^3$ is equals to

- a) $\frac{1}{2}x^3$ b) $2x^3$ c) $\frac{1}{4}(\log x)^3$ d) $\frac{1}{4}x^3$

Q29. C.F of $(x^2 D^2 - xD)y = 0$ is

- a) $a + bx$ b) $ax + bx^2$ c) $a \log x + bx^2$ d) $a + bx^2$

Q30. To convert the Euler Cauchy equation into a linear equation with constant coefficient we assume

- a) $x = e^t$ b) $x = t$ c) $x = t^2$ d) None of these

Q31. If the system of equations is $(2D - 4)y_1 + (3D + 5)y_2 = 3t + 2$, $(D - 2)y_1 + (D + 1)y_2 = t$ then y_2 is

- a) $ae^{-3t} + \frac{1}{9}(3t + 5)$ b) ae^{-3t} c) $ae^{-2t} + \frac{1}{9}(3t + 5)$ d) None of these

Q27. Which of the following is Euler-Cauchy equation?

- a) $x^3 y'' + x^2 y' + y = 0$ b) $x^2 y'' + xy' + y = 0$ c) $x^2 y'' + xy = 0$ d) None of these

Q40. Particular integral of $x^2 y'' - 2y = 2x + 6$ is

- a) $-x - 3$ b) $x - 3$ c) $-x + 3$ d) None of these

Q38. The complimentary function of the differential equation $x^2 y'' - xy' + y = \log x$ is given by

- a) $ae^x + be^{2x}$ b) ax c) $(a + b \log x)x$ d) $ax \log x$

Q36. For the given system of linear differential equation $y_1' = 2y_1 + y_2$, $y_2' = y_1 + 2y_2$, then the second order linear differential equation satisfied by y_1 is

- a) $y_1'' + 4y_1' + 3y_1 = 0$ b) $y_1'' - 4y_1' + 3y_1 = 0$ c) $y_1'' - 4y_1' - 3y_1 = 0$ d) None of these

❖ **Q.5** For the differential equation $(x^3 D^3 - 3xD + 3)y = 0$ using transformation $x = e^t$ the roots of its operator notation are:

- a) 1, 1, 3 b) 1, -1, 3 c) 1, 1, -3 d) 1, 2, 3

❖ **Q.25** If $y_1(t), y_2(t)$ satisfy the equations $y_1' + 5y_2 = 0$, $y_2' + y_1 = 0$ and $y_2'' + by_2 = 0$ is the second order differential equation satisfied by y_1 then what is the value of b

- a) 5 b) -5 c) 3 d) -3

❖ **Q.29** If $D = \frac{d}{dx}$, then $\frac{1}{(x^2 D^2 + 2)} 16x^3$ is equal to

- a) $\frac{1}{2}x^3$ b) $2x^3$ c) $\frac{1}{4}(\log x)^3$ d) $\frac{1}{4}x^3$

❖ For a given system of linear differential equation $y_1' = 2y_1 + y_2$, $y_2' = y_1 + 2y_2$, the second order linear differential satisfied by the y_1 is

- (A) $y_1'' + 4y_1' + 3y_1 = 0$ (B) $y_1'' - 4y_1' + 3y_1 = 0$ (C) $y_1'' - 4y_1' - 3y_1 = 0$ (D) none

- ❖ The particular integral of differential equation ($x > 0$)
 $x^3 y''' + 5x^2 y'' + 5xy' + y = x^2$ Using the transformation $x = e^t$, we get (in operator notation) $[\theta^3 + 2\theta^2 + 2\theta + 1]y = e^{2t}$ is
- (A) $\frac{1}{21}e^{2t}$ (B) $\frac{1}{31}e^{-2t}$ (C) $-\frac{1}{51}e^{2t}$ (D) $\frac{1}{21}e^{7t}$