

Assignment-1(EE5600)

Satyam Singh
EE20MTECH14015

Abstract—This assignment deals with basic coordinate geometry.

Download tex file from

<https://github.com/satyam463/EE5600Ass1/blob/main/Ass1.tex>

Consider Fig.0

$$\mathbf{B} = \begin{pmatrix} x_1 \\ y_1 \end{pmatrix}, \mathbf{C} = \begin{pmatrix} x_2 \\ y_2 \end{pmatrix}, \mathbf{A} = \begin{pmatrix} x_3 \\ y_3 \end{pmatrix} \quad (2.0.1)$$

The line joining BC divided into the ratio l:k at point of division \mathbf{P} can be written as

$$l(\mathbf{CP}) = k(\mathbf{PB}) \quad (2.0.2)$$

$$l(\mathbf{P} - \mathbf{C}) = k(\mathbf{B} - \mathbf{P}) \quad (2.0.3)$$

$$(l + k)\mathbf{P} = l\mathbf{C} + k\mathbf{B} \quad (2.0.4)$$

$$(l + k)\mathbf{P} = \begin{pmatrix} \mathbf{C} & \mathbf{B} \end{pmatrix} \begin{pmatrix} l \\ k \end{pmatrix} \quad (2.0.5)$$

Now, the line joining PA divided into the ratio m:k+l at point of division \mathbf{Q} can be written as

$$(l + k)(\mathbf{PQ}) = m(\mathbf{QA}) \quad (2.0.6)$$

$$(l + k)(\mathbf{Q} - \mathbf{P}) = m(\mathbf{A} - \mathbf{Q}) \quad (2.0.7)$$

$$(l + k + m)\mathbf{Q} = (l + k)\mathbf{P} + m\mathbf{A} \quad (2.0.8)$$

From Eq.2.0.4 substitute $(l + k)\mathbf{P}$ in Eq.2.0.8

$$(l + k + m)\mathbf{Q} = l\mathbf{C} + k\mathbf{B} + m\mathbf{A} \quad (2.0.9)$$

$$\mathbf{Q} = \frac{1}{l + k + m} \begin{pmatrix} \mathbf{C} & \mathbf{B} & \mathbf{A} \end{pmatrix} \begin{pmatrix} l \\ k \\ m \end{pmatrix} \quad (2.0.10)$$

Hence, \mathbf{Q} is the required coordinate of the latter point of section.

1 PROBLEM STATEMENT

1.1 Vector2, Example-1, 22

The coordinates of vertices of a triangle are (x_1, y_1) , (x_2, y_2) , and (x_3, y_3) . The line joining the first two is divided into the ratio l:k, and line joining this point of division to the opposite angular point is then divided in the ratio m:k+l. Find the coordinates of the latter point of section.

2 SOLUTION




Fig. 0: Triangle ABC with vertices $\mathbf{A} \begin{pmatrix} 2 \\ 4 \end{pmatrix}$, $\mathbf{B} \begin{pmatrix} 0 \\ 0 \end{pmatrix}$, $\mathbf{C} \begin{pmatrix} 4 \\ 0 \end{pmatrix}$, and $\begin{pmatrix} l \\ m \end{pmatrix} = \begin{pmatrix} 0.5 \\ 0.5 \end{pmatrix}$ are used for python plot. The point of section $\mathbf{P} = \begin{pmatrix} 2 \\ 0 \end{pmatrix}$, $\mathbf{Q} = \begin{pmatrix} 2 \\ 1.33 \end{pmatrix}$.