

# Prediction of the Effect of Single Amino Acid Protein Variants Using Deep Mutational Scanning Data

University of Bologna — Master Thesis in Bioinformatics

Pierotti Saul


Internal Advisor: Prof. Pietro Di Lena

External Advisor: Prof. Arne Elofsson (Stockholm University)


July 19, 2021

# Deep Mutational Scanning

High-throughput technique for obtaining fitness information on a large number of mutations




# Some Notable Variant Effect Predictors




Among these predictors only Envision was trained on deep mutational scanning data


# Structure of the Project



# I Considered Only Single Amino Acid Variants




# I Used the Training Dataset of Envision (Gray et al., 2018)




# Agreement among Experimental Results

Two independent deep mutational scanning experiments on Ubiquitin are present in the training dataset. Their correlation is low.



# Interesting Patterns in Mutation Sensitivity







- ▶ Polar residues seem more tolerant to mutations than hydrophobic residues
- ▶ Proline (P) is the most disruptive residue
- ▶ Tryptophan (W) is hard to replace




# Exposure Explains the Mutability of Polar Residues

When filtering by Relative Solvent Accessibility (RSA) apolar residues are **not** more sensitive to mutations than polar residues




# Buried Residues Are More Conserved




# Features Used for the Predictions

All the features are derived from the sequences: I did **not** use structural information




# Single Protein Models with Naive Testing




- ▶ A different model trained for each protein
- ▶ Half of the mutations set aside for testing
- ▶ Hyperparameters optimized in the remaining half with cross-validation
- ▶ Good results but likely overfitting of the testing set

# Single Protein Models with Testing by Position



- Same as before but mutations in the same position segregated in the training or testing sets
- Performances are lower (and more realistic)

# Leave-One-Protein-Out (LOPO) Models



- ▶ Models trained on the whole dataset while leaving one protein out
- ▶ For the left-out protein, half of the mutations used for testing and half for validation
- ▶ Spearman correlation coefficient used for evaluation

# Discussion

Complex models do not improve much on linear regression

Unsupervised models perform similarly to supervised models

There is strong variability between datasets

How validation and testing are performed is crucial

Performances on par with other predictors can be reached without structural features

# Future Directions

Unsupervised models  
seem promising and may  
be worth exploring more

Training on more deep  
mutational scanning studies

Tuning the set  
of features

Using residue contacts in a graph  
convolutional neural network

Finding a better normalization strategy  
for the scores from different experiments



# Bibliography I


- Adzhubei, I. A., Schmidt, S., Peshkin, L., Ramensky, V. E., Gerasimova, A., Bork, P., Kondrashov, A. S. & Sunyaev, S. R. (2010). A method and server for predicting damaging missense mutations. *Nature Methods*, 7(4), 248–249. <https://doi.org/10.1038/nmeth0410-248>
- Eddy, S. R. (2011). Accelerated profile HMM searches (W. R. Pearson, Ed.). *PLoS Computational Biology*, 7(10), e1002195. <https://doi.org/10.1371/journal.pcbi.1002195>
- Gray, V. E., Hause, R. J., Luebeck, J., Shendure, J. & Fowler, D. M. (2018). Quantitative missense variant effect prediction using large-scale mutagenesis data. *Cell Systems*, 6(1), 116–124.e3. <https://doi.org/10.1016/j.cels.2017.11.003>
- Hecht, M., Bromberg, Y. & Rost, B. (2015). Better prediction of functional effects for sequence variants. *BMC Genomics*, 16(S8). <https://doi.org/10.1186/1471-2164-16-s8-s1>
- Hopf, T., Ingraham, J., Poelwijk, F., Schärfe, C., Springer, M., Sander, C. & Marks, D. (2017). Mutation effects predicted from sequence co-variation. *Nature Biotechnology*, 35(2), 128–135. <https://doi.org/10.1038/nbt.3769>

# Bibliography II

- Klausen, M. S., Jespersen, M. C., Nielsen, H., Jensen, K. K., Jurtz, V. I., Sønderby, C. K., Sommer, M. O. A., Winther, O., Nielsen, M., Petersen, B. & Marcatili, P. (2019). NetSurfP-2.0: Improved prediction of protein structural features by integrated deep learning. *Proteins: Structure, Function, and Bioinformatics*, 87(6), 520–527. <https://doi.org/10.1002/prot.25674>
- Riesselman, A. J., Ingraham, J. B. & Marks, D. S. (2018). Deep generative models of genetic variation capture the effects of mutations. *Nature Methods*, 15(10), 816–822. <https://doi.org/10.1038/s41592-018-0138-4>
- Sim, N.-I., Kumar, P., Hu, J., Henikoff, S., Schneider, G. & Ng, P. C. (2012). SIFT web server: Predicting effects of amino acid substitutions on proteins. *Nucleic Acids Research*, 40(W1), W452–W457. <https://doi.org/10.1093/nar/gks539>
- Yang, J., Anishchenko, I., Park, H., Peng, Z., Ovchinnikov, S. & Baker, D. (2020). Improved protein structure prediction using predicted interresidue orientations. *Proceedings of the National Academy of Sciences*, 117(3), 1496–1503. <https://doi.org/10.1073/pnas.1914677117>


# Supplementary Material

Secondary structure is of limited importance in the discrimination of damaging mutations



# Supplementary Material

The correlation among features follows predictable patterns



# Supplementary Material

## Precision of trRosetta (Yang et al., 2020) in predicting residue contacts

Dataset	Medium-range ( $s \geq 12$ )			Long-range ( $s \geq 24$ )		
	Top $L/5$	Top $L/2$	Top $L$	Top $L/5$	Top $L/2$	Top $L$
beta-lactamase	1.00	0.92	0.86	0.96	0.93	0.76
WW_domain	0.95	0.90	0.83	0.90	0.87	0.75
PSD95pdz3	0.96	0.92	0.80	0.92	0.81	0.70
kka2_1:2	1.00	1.00	0.96	1.00	1.00	0.89
hsp90	1.00	1.00	0.96	1.00	1.00	0.89
Ubiquitin	0.98	0.92	0.82	1.00	0.90	0.70
Pab1	0.80	0.72	0.67	0.87	0.74	0.60
E1_Ubiquitin	0.82	0.86	0.77	0.91	0.75	0.54
gb1	1.00	0.85	0.46	0.63	0.40	0.22

## Quality of the predicted structural features from NetsurfP-2 (Klausen et al., 2019)


Feature	Evaluation metric	Score
Relative solvent accessibility	Pearson $r$	0.79
Accessible surface area	Pearson $r$	0.80
$Q_3$ secondary structure	$Q_3$ accuracy	0.85
$Q_8$ secondary structure	$Q_8$ accuracy	0.72
$\phi$ torsion angle	Circular correlation	0.73
$\psi$ torsion angle	Circular correlation	0.87

## Relationship between the features used in the models and the fitness scores

Feature	Pearson $r$	Spearman $\rho$	Kendall $\tau$
PSSM mutation score	-0.29	-0.28	-0.19
Netsurf predicted RSA	0.34	0.37	0.25
Netsurf predicted ASA	0.32	0.35	0.24
Netsurf predicted disorder	0.06	0.18	0.12
EVcouplings epistatic model	0.46	0.50	0.34
EVcouplings independent model	0.44	0.44	0.30
EVcouplings frequency	0.19	0.35	0.24
EVcouplings conservation	-0.32	-0.33	-0.23
Closeness centrality (trRosetta predicted contacts)	-0.16	-0.17	-0.11
Betweenness centrality (trRosetta predicted contacts)	-0.20	-0.29	-0.19
Degree centrality (trRosetta predicted contacts)	-0.12	-0.13	-0.09
Load centrality (trRosetta predicted contacts)	-0.20	-0.29	-0.19
Harmonic centrality (trRosetta predicted contacts)	-0.19	-0.20	-0.14
Clustering coefficient (trRosetta predicted contacts)	0.23	0.25	0.17
Linear-circular correlation			
Netsurf predicted $\phi$ torsion angle			0.01
Netsurf predicted $\psi$ torsion angle			0.02
	Kruskal-Wallis $\chi^2$	$p$ -value	
Wild-type residue	1482.40	$< 2.20 \cdot 10^{-16}$	
Mutated residue	708.53	$< 2.20 \cdot 10^{-16}$	
Netsurf predicted $Q_3$ secondary structure	215.33	$< 2.20 \cdot 10^{-16}$	
Netsurf predicted $Q_8$ secondary structure	351.97	$< 2.20 \cdot 10^{-16}$	


# Supplementary Material

## Feature importances for the gradient boosted tree general models



# Supplementary Material


## Feature importances for the linear regression general models





# Supplementary Material

Comparison of the performances of single protein models and general models



# Supplementary Material

## Confidence intervals in prediction performances

Dataset name	Model	95 % C.I. (Pearson)	95 % C.I. (Spearman)
beta-lactamase	Naive	0.89 to 0.91	0.86 to 0.88
beta-lactamase	By position	0.79 to 0.83	0.75 to 0.78
beta-lactamase	LOPO	—	0.68 to 0.72
WW_domain	Naive	0.67 to 0.82	0.70 to 0.84
WW_domain	By position	0.57 to 0.73	0.52 to 0.72
WW_domain	LOPO	—	0.56 to 0.73
PSD95pdz3	Naive	0.74 to 0.83	0.73 to 0.80
PSD95pdz3	By position	0.55 to 0.67	0.54 to 0.65
PSD95pdz3	LOPO	—	0.53 to 0.63
kka2_1:2	Naive	0.74 to 0.78	0.72 to 0.76
kka2_1:2	By position	0.65 to 0.69	0.64 to 0.68
kka2_1:2	LOPO	—	0.60 to 0.64
hsp90	Naive	0.82 to 0.87	0.65 to 0.71
hsp90	By position	0.69 to 0.76	0.50 to 0.57
hsp90	LOPO	—	0.38 to 0.45
Ubiquitin	Naive	0.75 to 0.83	0.78 to 0.83
Ubiquitin	By position	0.49 to 0.60	0.46 to 0.57
Ubiquitin	LOPO	—	0.30 to 0.43
Pab1	Naive	0.80 to 0.87	0.75 to 0.82
Pab1	By position	0.65 to 0.76	0.64 to 0.73
Pab1	LOPO	—	0.60 to 0.70
E1_Ubiquitin	Naive	0.75 to 0.85	0.65 to 0.75
E1_Ubiquitin	By position	0.59 to 0.72	0.36 to 0.51
E1_Ubiquitin	LOPO	—	0.50 to 0.63
gb1	Naive	0.90 to 0.93	0.88 to 0.92
gb1	By position	0.31 to 0.46	0.29 to 0.45
gb1	LOPO	—	0.32 to 0.47

# Supplementary Material

Statistical significance of performance differences. Starred values are significant with Bonferroni correction.

$$\alpha = \frac{0.05}{27} = 0.00185185$$

Dataset name	Model 1	Model 2	p-value
beta-lactamase	Linear regression	Gradient boosted trees	$1 \cdot 10^{-4} *$
beta-lactamase	Linear regression	EVmutation	$1 \cdot 10^{-4} *$
beta-lactamase	EVmutation	Gradient boosted trees	$1 \cdot 10^{-4} *$
WW_domain	Linear regression	Gradient boosted trees	$1 \cdot 10^{-4} *$
WW_domain	Linear regression	EVmutation	$1 \cdot 10^{-4} *$
WW_domain	EVmutation	Gradient boosted trees	$1 \cdot 10^{-4} *$
PSD95pdz3	Linear regression	Gradient boosted trees	$1 \cdot 10^{-4} *$
PSD95pdz3	Linear regression	EVmutation	$1 \cdot 10^{-4} *$
PSD95pdz3	EVmutation	Gradient boosted trees	0.51
kka2_1:2	Linear regression	Gradient boosted trees	0.00
kka2_1:2	Linear regression	EVmutation	0.01
kka2_1:2	EVmutation	Gradient boosted trees	$1 \cdot 10^{-4} *$
hsp90	Linear regression	Gradient boosted trees	$1 \cdot 10^{-4} *$
hsp90	Linear regression	EVmutation	$1 \cdot 10^{-4} *$
hsp90	EVmutation	Gradient boosted trees	0.24

The table continues on the next slide

# Supplementary Material

Statistical significance of performance differences. Starred values are significant with Bonferroni correction.

$$\alpha = \frac{0.05}{27} = 0.00185185$$

The table continues from the previous slide

Dataset name	Model 1	Model 2	<i>p</i> -value
Ubiquitin	Linear regression	Gradient boosted trees	$1 \cdot 10^{-4} *$
Ubiquitin	Linear regression	EVmutation	$1 \cdot 10^{-4} *$
Ubiquitin	EVmutation	Gradient boosted trees	$1 \cdot 10^{-4} *$
Pab1	Linear regression	Gradient boosted trees	$1 \cdot 10^{-4} *$
Pab1	Linear regression	EVmutation	$1 \cdot 10^{-4} *$
Pab1	EVmutation	Gradient boosted trees	$1 \cdot 10^{-4} *$
E1_Ubiquitin	Linear regression	Gradient boosted trees	$1 \cdot 10^{-4} *$
E1_Ubiquitin	Linear regression	EVmutation	$1 \cdot 10^{-4} *$
E1_Ubiquitin	EVmutation	Gradient boosted trees	$1 \cdot 10^{-4} *$
gb1	Linear regression	Gradient boosted trees	$1 \cdot 10^{-4} *$
gb1	Linear regression	EVmutation	$1 \cdot 10^{-4} *$
gb1	EVmutation	Gradient boosted trees	$1 \cdot 10^{-4} *$