The Emissions–Temperature Relationship

> Part of the Functional Series. Warming tracks the CO_2 humans have put into the air over time. This document explains how the platform measures that relationship from observed data, how it compares with the IPCC's own figure, and how much weight the number can carry. The input series are described in Concentrations and Temperature.

1. Cumulative, not annual

A single year's emissions say little about that year's temperature: CO_2 stays in the atmosphere for a very long time, so today's warming reflects everything emitted so far. The climate-science term for the resulting relationship is the **transient climate response to cumulative CO_2 emissions (TCRE)**: roughly, how many degrees of warming follow from each additional trillion tonnes of CO_2.

The platform measures a simplified, data-driven analogue of it: a straight-line fit of the observed temperature anomaly against the observed cumulative CO_2 emitted since 1850.

Temperature (y)	Berkeley Earth global anomaly relative to 1850–1900, annual, 1850–2024
Emissions (x)	World cumulative CO_2 since 1850: fossil fuels and cement, **plus land-use change**, from OWID/Global Carbon Project, including international aviation and shipping
Fit	Ordinary least squares with an intercept; 175 annual points
Unit of the answer	°C per 1,000 $GtCO_2$ (a thousand billion tonnes)

2. The result

Quantity	Value
Slope, total anthropogenic CO_2	**0.485 °C per 1,000 $GtCO_2$**
95% interval (robust to autocorrelation)	0.441 – 0.529
R^2	0.89
Cumulative CO_2 since 1850, to 2024	about 2,750 $GtCO_2$ total; 1,849 $GtCO_2$ fossil and cement only
Same fit, fossil and cement only	0.749 (0.705 – 0.793)

Reading R^2 = 0.89. Cumulative emissions account for most of the long-run variation in the temperature record. That is a statement of association over 175 years in which both series trend upward; it is not, by itself, proof of cause. The cause is established by physics and other evidence, and the platform states plainly that this is "context, not proof".

3. Comparison with the IPCC range

The IPCC's Sixth Assessment Report gives a TCRE best estimate of 0.45 °C per 1,000 $GtCO_2$, with a **likely** range of 0.27–0.63 (Working Group I Summary for Policymakers, D.1.1; "likely" means at least a 66% probability). A *95% statistical interval* around a fitted slope and a *66% assessed range* from many lines of evidence are different kinds of interval, so "overlaps" below means the two are consistent, not that one validates the other.

Measure	Platform	Versus IPCC AR6
Total-CO_2 slope	0.485	Inside the range; 1.08 × the best estimate; the 95% interval overlaps the range
Fossil-and-cement-only slope	0.749	Above the range

The fossil-only figure sits higher because it leaves out land-use emissions: the same warming is being attributed to a smaller amount of CO_2, so each tonne appears to do more. The comparison is only meaningful for the total-CO_2 version, which matches the IPCC definition more closely, and even that is an analogue, not a restatement, because:

- the regression also absorbs warming from **non-CO_2 gases and aerosols** that rose and fell alongside CO_2;
- it uses **one observed climate history**, not an ensemble of Earth-system models;
- it depends on **land-use emission estimates** that are themselves uncertain (about ±2.6 $GtCO_2$ per year, one standard deviation, in recent decades).

4. How the uncertainty is handled

The 175 annual points are not independent: a warm year tends to follow a warm year. Ignoring that makes the fit look more certain than it is.

Check	What it shows
Residuals are autocorrelated	Lag-1 autocorrelation 0.59; Durbin–Watson 0.80 (a value near 2 would mean none)
Robust standard error	Newey–West (HAC) standard error 0.022, against 0.013 from the naive formula: about 1.7 × larger, so the naive interval would be too narrow
Lag-length sensitivity	The interval is 0.447–0.523, 0.441–0.529 and 0.435–0.535 for 4, 8 and 16 lags
Block bootstrap	Resampling the residuals in 10-year blocks, 2,000 times, gives 0.435–0.535, close to the robust interval; other block lengths from 5 to 30 years give intervals of similar width
Holdouts	The fit is also tested by fitting on early decades and predicting later ones, against a simple baseline

5. The relationship is not constant

Fitting only more recent periods gives larger slopes:

Fit period	Slope (°C per 1,000 $GtCO_2$)
1850–2024	0.485
1900–2024	0.527
1950–2024	0.581
1970–2024	0.628

If CO_2 alone set the temperature through a fixed proportionality, the slope would not drift this way. The drift is consistent with other influences changing over time (non-CO_2 gases, aerosols, the pace of ocean heat uptake), and it means the headline number is the best single summary of the full record, not a law. The platform uses the full-range value for its scenario translation and shows the shorter windows so a reader can see the sensitivity.

6. What this is not

- Not a climate model: no physics, no ocean, no feedbacks.
- Not an attribution of warming to any country. Country shares describe each country's part of *cumulative emissions*; no country series is regressed against temperature.
- Not a forecast on its own. Its use in the scenario translation is described in Scenarios.

7. Where this appears in the dashboard

Dashboard element	What it shows
Climate Correlation page	The relationship chart, the headline slope, and an IPCC AR6 comparison strip
"How this number was derived"	A seven-step explanation: what is regressed on what, the uncertainty, sensitivity, stability, the IPCC comparison, data and licences, and what the number is *not*
"Methodology & sources"	The sources table (coverage, vintage, licence), the baseline used by each view, the temperature offset and the two-totals note

See also

- Scenarios and Temperature Mapping
- Limitations and Uncertainty
- Technical: Multi-Source Data Pipelines: how the fit and its uncertainty are computed