Concentrations and Temperature: NOAA Mauna Loa, Law Dome and Berkeley Earth

> Part of the Functional Series. Emissions are only the first link of the chain. This document describes the two series that sit downstream of them: how much CO_2 has accumulated in the air, and how much the surface has warmed. Emissions are in the previous document.

1. Atmospheric CO_2: NOAA Mauna Loa and Law Dome

What it measures. The concentration of CO_2 in the global atmosphere, in parts per million (ppm). Unlike emissions, which are a *flow* (tonnes per year), concentration is a *stock*: what is left in the air after the oceans and land have taken up their share.

Two sources, spliced at 1959.

Period	Source	How obtained
1959 onward	NOAA Global Monitoring Laboratory, Mauna Loa Observatory (Hawaii)	Direct, continuous, in-situ measurement; 1958–1974 is the Scripps (Keeling) record that NOAA took over
Before 1959	Law Dome ice cores and firn (Antarctica)	Air trapped in ice, dated and smoothed into an annual spline (Etheridge et al.; MacFarling Meure et al.)

The two are joined in 1959 without adjustment. The platform stores how well they agree where they overlap (1959–2004): on average the ice-core series is 2.3 ppm below the instrument record, and the largest gap is 3.9 ppm. At the splice year itself the gap is 0.28 ppm.

Latest values. The 2025 annual mean is **427.35 ppm**, with 0.12 ppm measurement uncertainty. A monthly series (from March 1958) feeds the "latest reading" indicator on the dashboard; it shows the familiar seasonal cycle, and the platform also carries NOAA's deseasonalized value.

What to remember.

- **One station.** Mauna Loa is a single Northern Hemisphere site, and Law Dome reflects Southern Hemisphere air. Atmospheric CO_2 is well mixed, so the difference is small, but pre-1959 values are not directly comparable year to year with later ones.
- **Smoothed ice-core data.** The Law Dome spline attenuates variations shorter than about 20 years by half. It is a good record of the long rise, and a poor guide to individual years. No per-year uncertainty is published for it, so the platform's uncertainty column is empty before 1959.
- **A data gap.** The 2022 Mauna Loa eruption interrupted the record; NOAA fills December 2022 to July 2023 with measurements from Maunakea.
- **Licence.** NOAA data are public domain with a citation requested.

2. Global temperature: Berkeley Earth

What it measures. The annual change in global average surface temperature (land air temperature and sea surface temperature) as an **anomaly**: the difference from a reference period, not an absolute temperature.

The product. Berkeley Earth's High-Resolution Land/Ocean analysis, a 0.25° gridded field that combines a land-surface field with HadSST4 ocean data and extrapolates over sea ice from nearby land air temperature. Uncertainty is a 95% interval covering statistical and spatial undersampling and ocean biases. The platform holds 1850–2025.

Two reference periods.

Reference	Why it matters
1951–1980 (the file's native baseline)	Berkeley Earth's own convention
1850–1900 ("pre-industrial")	The reference used by the Paris Agreement and the IPCC. The platform *computes* the offset from the same dataset: the 1850–1900 mean is **–0.265 °C** relative to 1951–1980 (standard deviation 0.117 °C over 51 years), and subtracting it converts any anomaly to the Paris-relevant scale

Subtracting that offset gives the headline numbers: **2025 is 1.45 °C and 2024 is 1.55 °C above 1850–1900**, with a 95% statistical uncertainty of about ±0.04 °C. As a cross-check, the 2025 value agrees with Berkeley Earth's own published 1.44 ± 0.09 °C.

What to remember.

- **Preliminary.** Berkeley Earth labels this dataset "preliminary data — subject to change without notice — not yet peer reviewed". Values can be revised and no final citation exists yet. The platform replaced an older summary file in October 2026 after Berkeley Earth confirmed that the high-resolution product is its operational dataset.
- **"Pre-industrial" is a convention.** 1850–1900 is a stand-in for conditions before large-scale fossil fuel use. The offset comes from the dataset's earliest years, where uncertainty is largest. Other groups choose different baselines or reconstruct them differently, and that alone shifts the headline "°C above pre-industrial" by a few hundredths of a degree.
- **One year is not a trend.** Annual values are influenced by El Niño and La Niña; the Paris Agreement's temperature goals are about long-term averages, not single years. The Benchmarks section (in development) returns to this.
- **Licence.** CC BY-NC: non-commercial use and attribution to Berkeley Earth. The platform's use is non-commercial; any commercial use would need a licence from them.

3. How the three series connect

Link	Series	Unit	Nature
Emissions	OWID / GCP, PRIMAP-hist	Mt per year	Flow, inventory-based

Link	Series	Unit	Nature
Concentration	NOAA Mauna Loa + Law Dome	ppm	Stock, measured
Temperature	Berkeley Earth	°C above 1850–1900	Response, observed

The platform pairs them by calendar year and relates *cumulative* emissions to temperature, because warming tracks the total CO_2 added to the atmosphere over time, not a single year's emissions. The Methodology section (in development) covers how, and how much weight that relationship can bear.

See also

- Emissions Inventories
- Data Quality and Harmonization
- Technical: Multi-Source Data Pipelines — ingestion of these two series and the pairing of series by year