

# EE5609: Matrix Theory

## Assignment-4

Major Saurabh Joshi  
MTech Artificial Intelligence  
AI20MTECH13002

**Abstract**—This document contains solution to determine the conic representing the given equation.

Download the latex-tikz codes from latex-tikz codes from

<https://github.com/saurabh13002/EE5609/tree/master/Assignment4>

### 1 PROBLEM

What conic does the following equation represent.

$$y^2 - 2\sqrt{3}xy + 3x^2 + 6x - 4y + 5 = 0$$

Find the center and equation referred to centre.

### 2 SOLUTION

The general second degree equation can be expressed as follows,

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (2.0.1)$$

From the given second degree equation we get,

$$\mathbf{V} = \begin{pmatrix} 3 & -\sqrt{3} \\ -\sqrt{3} & 1 \end{pmatrix} \quad (2.0.2)$$

$$\mathbf{u} = \begin{pmatrix} 3 \\ -2 \end{pmatrix} \quad (2.0.3)$$

$$f = 5 \quad (2.0.4)$$

Expanding the determinant of  $\mathbf{V}$  we observe,

$$\begin{vmatrix} 3 & -\sqrt{3} \\ -\sqrt{3} & 1 \end{vmatrix} = 0 \quad (2.0.5)$$

Also

$$\begin{vmatrix} \mathbf{V} & \mathbf{u} \\ \mathbf{u}^T & f \end{vmatrix} = \begin{vmatrix} 3 & -\sqrt{3} & 3 \\ -\sqrt{3} & 1 & -2 \\ 3 & -2 & 5 \end{vmatrix} \neq 0 \quad (2.0.6)$$

Hence from (2.0.5) and (2.0.6) we conclude that given equation is a parabola. The characteristic equation of  $\mathbf{V}$  is given as follows,

$$|\mathbf{V} - \lambda \mathbf{I}| = \begin{vmatrix} 3 - \lambda & -\sqrt{3} \\ -\sqrt{3} & 1 - \lambda \end{vmatrix} = 0 \quad (2.0.7)$$

$$\implies \lambda^2 - 4\lambda = 0 \quad (2.0.8)$$

Hence the characteristic equation of  $\mathbf{V}$  is given by (2.0.8). The roots of (2.0.8) i.e the eigenvalues are given by

$$\lambda_1 = 0, \lambda_2 = 4 \quad (2.0.9)$$

The eigen vector  $\mathbf{p}$  is defined as,

$$\mathbf{V} \mathbf{p} = \lambda \mathbf{p} \quad (2.0.10)$$

$$\implies (\mathbf{V} - \lambda \mathbf{I}) \mathbf{p} = 0 \quad (2.0.11)$$

for  $\lambda_1 = 0$ ,

$$(\mathbf{V} - \lambda_1 \mathbf{I}) = \begin{pmatrix} 3 & -\sqrt{3} \\ -\sqrt{3} & 1 \end{pmatrix} \xrightarrow[R_1 = \frac{1}{\sqrt{3}} R_1]{R_2 = R_1 + R_2} \begin{pmatrix} \sqrt{3} & -1 \\ 0 & 0 \end{pmatrix} \quad (2.0.12)$$

Substituting equation 2.0.12 in equation 2.0.11 and upon normalizing we get we get

$$\implies \mathbf{p}_1 = \begin{pmatrix} 1/2 \\ \sqrt{3}/2 \end{pmatrix} \quad (2.0.13)$$

Again, for  $\lambda_2 = 4$ ,

$$(\mathbf{V} - \lambda_2 \mathbf{I}) = \begin{pmatrix} -1 & -\sqrt{3} \\ -\sqrt{3} & -3 \end{pmatrix} \xrightarrow[R_1 = -\sqrt{3} R_1]{R_2 = -\sqrt{3} R_1 + R_2} \begin{pmatrix} 1 & \sqrt{3} \\ 0 & 0 \end{pmatrix} \quad (2.0.14)$$

Substituting equation 2.0.14 in equation 2.0.11 and upon normalizing we get

$$\mathbf{p}_2 = \begin{pmatrix} -\sqrt{3}/2 \\ 1/2 \end{pmatrix} \quad (2.0.15)$$

The matrix P,

$$\mathbf{P} = (\mathbf{p}_1 \quad \mathbf{p}_2) = \begin{pmatrix} 1/2 & -\sqrt{3}/2 \\ \sqrt{3}/2 & 1/2 \end{pmatrix} \quad (2.0.16)$$

$$\mathbf{D} = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix} \quad (2.0.17)$$

$$\eta = 2\mathbf{p}_1^T \mathbf{u} = 3 - 2\sqrt{3} \quad (2.0.18)$$

The focal length of the parabola is given by:

$$\left| \frac{\eta}{\lambda_2} \right| = \left| \frac{3 - 2\sqrt{3}}{4} \right| = 0.116 \quad (2.0.19)$$

When  $|\mathbf{V}| = 0$ , (2.0.1) can be written as

$$\mathbf{y}^T \mathbf{D} \mathbf{y} = -\eta \begin{pmatrix} 1 & 0 \end{pmatrix} \mathbf{y} \quad (2.0.20)$$

And the vertex  $\mathbf{c}$  is given by

$$\begin{pmatrix} \mathbf{u}^T + \frac{\eta}{2} \mathbf{p}_1^T \\ \mathbf{V} \end{pmatrix} \mathbf{c} = \begin{pmatrix} -f \\ \frac{\eta}{2} \mathbf{p}_1 - \mathbf{u} \end{pmatrix} \quad (2.0.21)$$

Substituting the found values

$$\mathbf{u}^T + \frac{\eta}{2} \mathbf{p}_1^T = \begin{pmatrix} 3 & -2 \end{pmatrix} + \frac{3 - 2\sqrt{3}}{2} \begin{pmatrix} 1/2 & \sqrt{3}/2 \end{pmatrix} \quad (2.0.22)$$

$$\Rightarrow \mathbf{u}^T + \frac{\eta}{2} \mathbf{p}_1^T = \begin{pmatrix} \frac{15-2\sqrt{3}}{4} & \frac{-14-3\sqrt{3}}{4} \end{pmatrix} \quad (2.0.23)$$

$$\frac{\eta}{2} \mathbf{p}_1 - \mathbf{u} = \begin{pmatrix} \frac{-9-2\sqrt{3}}{4} \\ \frac{2+3\sqrt{3}}{4} \end{pmatrix} \quad (2.0.24)$$

using equations (2.0.3),(2.0.4),(2.0.13),(2.0.23),(2.0.24) and (2.0.13) in (2.0.21)

$$\begin{pmatrix} 2.88 & -2.20 \\ 3 & -\sqrt{3} \\ -\sqrt{3} & 3 \end{pmatrix} \mathbf{c} = \begin{pmatrix} -5 \\ -3.11 \\ 1.8 \end{pmatrix} \quad (2.0.25)$$

By performing row reductions on augmented matrix

$$\begin{pmatrix} 2.88 & -2.2 & -5 \\ 3 & -\sqrt{3} & -3.11 \\ -\sqrt{3} & 1 & 1.8 \end{pmatrix} \xleftrightarrow{R_2 \leftarrow R_2 / \sqrt{3}} \quad (2.0.26)$$

$$\begin{pmatrix} 2.88 & -2.2 & -5 \\ \sqrt{3} & -1 & -1.8 \\ -\sqrt{3} & 1 & 1.8 \end{pmatrix} \quad (2.0.27)$$

$$\xleftrightarrow{R_3 \leftarrow R_2 + R_3} \begin{pmatrix} 2.88 & -2.2 & -5 \\ \sqrt{3} & -1 & -1.8 \\ 0 & 0 & 0 \end{pmatrix} \quad (2.0.28)$$

Therefore,

$$\begin{pmatrix} 2.88 & -2.2 & -5 \\ \sqrt{3} & -1 & -1.8 \end{pmatrix} \xleftrightarrow[R_1 = \sqrt{3}R_1]{R_2 = (2.88)R_2} \begin{pmatrix} 4.99 & -3.81 & -8.66 \\ 4.99 & -2.88 & -5.191 \end{pmatrix} \quad (2.0.29)$$

$$\Rightarrow \begin{pmatrix} 4.99 & -3.81 & -8.66 \\ 4.99 & -2.88 & -5.191 \end{pmatrix} \xleftrightarrow{R_2 = R_2 - R_1} \begin{pmatrix} 4.99 & -3.81 & -8.66 \\ 0 & 0.93 & 3.46 \end{pmatrix} \quad (2.0.30)$$

Hence we get  $\mathbf{c} = \begin{pmatrix} 1.10 \\ 3.72 \end{pmatrix}$ . The vertex of parabola at (1.10, 3.72).




Fig. 1: Parabola with the center c