

EE5609: Matrix Theory

Assignment-8

Major Saurabh Joshi
MTech Artificial Intelligence
AI20MTECH13002

Abstract—This document uses properties of vector spaces and subspaces.

Download the latex code from

<https://github.com/saurabh13002/EE5609/tree/master/Assignment8>

1 PROBLEM

Let \mathbf{V} be the (real) vector space of all functions f from \mathbf{R} into \mathbf{R} . Is $f(0) = f(1)$ a subspace of \mathbf{V}

2 THEOREM

A non-empty subset \mathbf{W} of \mathbf{V} is a subspace of \mathbf{V} if and only if for each pair of vectors α, β in \mathbf{W} and each scalar c in \mathbf{R} the vector $c\alpha + \beta$ is again in \mathbf{W} .

3 SOLUTION

For each of the function to be a subspace, it must be closed with respect to addition and scalar multiplication in \mathbf{V} defined as, for $f, g \in \mathbf{W}$ and $c \in \mathbf{R}$

Then,

$$h = cf + g \quad (3.0.1)$$

$$h(0) = cf(0) + g(0) \quad (3.0.2)$$

$$= cf(1) + g(1) \quad (3.0.3)$$

$$= h(1) \quad (3.0.4)$$

Thus, $h(0) = h(1)$. Therefore, \mathbf{W} is a subset of \mathbf{V} and also a vector space. Therefore \mathbf{W} is a subspace of \mathbf{V} .

Hence, $f(0) = f(1)$ is a subspace of \mathbf{V} .