

This Provisional PDF corresponds to the article as it appeared upon acceptance. Copyedited and fully formatted PDF and full text (HTML) versions will be made available soon.

## **Telmisartan/hydrochlorothiazide versus valsartan/hydrochlorothiazide in obese hypertensive patients with type 2 diabetes: the SMOOTH study**

*Cardiovascular Diabetology* 2007, **6**:28 doi:10.1186/1475-2840-6-28

Arya M Sharma (sharma@ccc.mcmaster.ca)  
Jaime Davidson (j davidson1@tmo.blackberry.net)  
Stephen Koval (skoval@rdg.boehringer-ingelheim.com)  
Yves Lacourciere (yves.lacourciere@crchul.ulaval.ca)

**ISSN** 1475-2840

**Article type** Original investigation

**Submission date** 31 July 2007

**Acceptance date** 2 October 2007

**Publication date** 2 October 2007

**Article URL** <http://www.cardiab.com/content/6/1/28>

This peer-reviewed article was published immediately upon acceptance. It can be downloaded, printed and distributed freely for any purposes (see copyright notice below).

Articles in *Cardiovascular Diabetology* are listed in PubMed and archived at PubMed Central.

For information about publishing your research in *Cardiovascular Diabetology* or any BioMed Central journal, go to

<http://www.cardiab.com/info/instructions/>

For information about other BioMed Central publications go to

<http://www.biomedcentral.com/>

**Telmisartan/hydrochlorothiazide versus  
valsartan/hydrochlorothiazide in obese hypertensive patients  
with type 2 diabetes: the SMOOTH study**

**Arya M Sharma<sup>\*1</sup>, Jaime Davidson<sup>2</sup>, Stephen Koval<sup>3</sup>, Yves Lacourciere<sup>4</sup>**

<sup>1</sup>McMaster University, Hamilton, ON, Canada, <sup>2</sup>the Endocrine and Diabetes Associates of Texas, Dallas, TX, USA, <sup>3</sup>Boehringer Ingelheim Pharmaceuticals, Inc, Ridgefield, CT, USA and <sup>4</sup>Centre Hospitalier Universitaire de Québec - Pav CHUL, Sainte-Foy, Quebec, Canada.

\*Corresponding author: Dr A M Sharma, McMaster University, Michael G deGroote School of Medicine, Hamilton, ON, Canada; tel. +1 905 527 4322 ext 46439, fax +1 905 525 2260; e-mail: sharma@ccc.mcmaster.ca

## **Abstract**

**Background:** The Study of Micardis (telmisartan) in Overweight/Obese patients with Type 2 diabetes and Hypertension (SMOOTH) compared hydrochlorothiazide (HCTZ) plus telmisartan or valsartan fixed-dose combination therapies on early morning blood pressure (BP), using ambulatory BP monitoring (ABPM).

**Methods:** SMOOTH was a prospective, randomized, open-label, blinded-endpoint, multicentre trial. After a 2- to 4-week, single-blind, placebo run-in period, patients received once-daily telmisartan 80 mg or valsartan 160 mg for 4 weeks, with add-on HCTZ 12.5 mg for 6 weeks (T/HCTZ or V/HCTZ, respectively). At baseline and week 10, ambulatory blood pressure (ABP) was measured every 20 min and hourly means were calculated. The primary endpoint was change from baseline in mean ambulatory systolic and diastolic blood pressure (SBP; DBP) during the last 6 hours of the 24-hour dosing interval.

**Results:** In total, 840 patients were randomized. At week 10, T/HCTZ provided significantly greater reductions versus V/HCTZ in the last 6 hours mean ABP (differences in favour of T/HCTZ: SBP 3.9 mm Hg,  $p < 0.0001$ ; DBP 2.0 mm Hg,  $p = 0.0007$ ). T/HCTZ also produced significantly greater reductions than V/HCTZ in 24-hour mean ABP (differences in favour of T/HCTZ: SBP 3.0 mm Hg,  $p = 0.0002$ ; DBP 1.6 mm Hg,  $p = 0.0006$ ) and during the morning, daytime and night-time periods ( $p < 0.003$ ). Both treatments were well tolerated.

**Conclusions:** In high-risk, overweight/obese patients with hypertension and type 2 diabetes, T/HCTZ provides significantly greater BP lowering versus

V/HCTZ throughout the 24-hour dosing interval, particularly during the hazardous early morning hours.

## Background

Hypertension, obesity and type 2 diabetes are cardiovascular risk factors that commonly occur together. Insufficient suppression of the renin–angiotensin–aldosterone system (RAAS) has been implicated in the development of obesity-related high arterial pressure, and is linked with insulin resistance and type 2 diabetes [1,2]. RAAS blockade may, therefore, be particularly beneficial in the antihypertensive treatment of patients with type 2 diabetes, features of metabolic syndrome and obesity, particularly as this population is poorly controlled. In a cross-sectional prevalence study of 45,125 subjects from Germany, hypertension (blood pressure [BP]  $\geq 140/90$  mm Hg) was twice as common in obese as in non-obese patients (60.6 vs. 34.3%, respectively) [3]. Furthermore, BP control in diagnosed and treated obese hypertensive patients was extremely low (overall response [OR] = 0.8).

Adequate BP control is also overlooked during the morning hours. During this time there is typically a surge in BP, which is associated with a high incidence of cerebro- and cardiovascular events [4]. Consequently, the early morning hours are an important therapeutic target for antihypertensive treatment [5].

Telmisartan is a once-daily angiotensin II receptor blocker (ARB) with the longest plasma half-life of any ARB, providing 24-hour coverage of BP control from a single daily dose; the angiotensin type 1 ( $AT_1$ ) versus  $AT_2$  receptor affinity ratio for telmisartan is 3000-fold; however, it is higher for valsartan (about 20,000-fold) [6–8]. In two randomized studies of 1,279 hypertensive patients, telmisartan 80 mg significantly reduced the early morning systolic BP (SBP) surge compared with ramipril 10 mg [9]. Another

ARB, valsartan has been shown to improve obesity-related disorders, reduce body mass index (BMI) and lower BP [10]. Based on such findings, it is therefore relevant to compare telmisartan with valsartan. A previous pooled analysis of two studies in patients with uncomplicated hypertension showed that telmisartan 80 mg provided SBP reductions in the last 6 hours of the dosing interval and in the 24-hour mean that were superior to the equipotent valsartan 160 mg (by 2.7 and 2.0 mm Hg, respectively) [11]. In addition, two recent studies have shown that telmisartan 80 mg plus hydrochlorothiazide (HCTZ) 25 mg was superior to valsartan 160 mg plus HCTZ 25 mg [12,13]. However, there are few studies comparing telmisartan and valsartan when used in combination with low-dose HCTZ 12.5 mg, and few direct ARB comparisons in obese hypertensive patients with type 2 diabetes.

In this Study of Micardis in Overweight/Obese patients with Type 2 diabetes and Hypertension (SMOOTH), the effect on early morning BP of the fixed-dose combinations, telmisartan 80 mg plus HCTZ 12.5 mg (T/HCTZ) and valsartan 160 mg plus HCTZ 12.5 mg (V/HCTZ), were compared using ambulatory BP monitoring (ABPM). This is one of the largest ABPM studies performed in obese hypertensive patients.

## **Methods**

Men and women aged  $\geq 30$  years with mild-to-moderate hypertension, defined as mean seated cuff SBP 140–179 mm Hg and/or diastolic BP (DBP) 95–109 mm Hg were randomized. Patients were also required to have 24-hour mean ambulatory SBP  $\geq 130$  mm Hg and/or DBP  $\geq 85$  mm Hg, type 2 diabetes that had remained stable and controlled for  $\geq 3$  months, defined as glycolated

haemoglobin (HbA<sub>1c</sub>)  $\leq$  10% and a BMI  $\geq$  27 kg/m<sup>2</sup> in non-Asians and  $\geq$  24 kg/m<sup>2</sup> in Asians (patients' upper arm circumference was required to be  $\leq$  52 cm to maximize the accuracy of BP readings).

Patients were excluded from randomization if they had mean seated SBP  $\geq$  180 mm Hg or mean seated DBP  $\geq$  110 mm Hg during any visit of the placebo run-in period, or fasting serum glucose  $>$  17 mmol/l (300 mg/dl).

Premenopausal women who were nursing, pregnant or not using adequate contraception were excluded. In addition, patients with a history of coronary disease, congestive heart failure, or a recent acute cardiovascular event (previous 3 months) or stroke (previous 6 months) were excluded, as were those with secondary hypertension. Patients were not eligible for randomization if they had hepatic or renal impairment. Night-shift workers were also excluded.

The use of corticosteroids, cholestyramine and colestipol resins was prohibited throughout the SMOOTH study. In addition, medications known to affect BP were not allowed. These included vasopressors, vasodilators, beta-agonists and -antagonists, nitroglycerin spray or sublingual tablets, theophylline, dipyridamole, monoamine oxidase inhibitors, phenothiazine and tricyclic antidepressants, chronic use of oral, nasal or topical decongestants and antihypertensive medication other than the study drugs.

### ***Study design***

SMOOTH (clinical trial number: 502.399) used a prospective, randomized, open-label, blinded-endpoint (PROBE), multicentre design. A published meta-analysis has shown that blinded 24-h systolic and diastolic ambulatory BP

results from PROBE studies evaluating antihypertensive therapy were statistically equivalent to double-blind, placebo-controlled (DBPC) studies. [14]. These findings show that the PROBE study design is an attractive and validated alternative to the DBPC design, particularly in hypertension trials using ABPM. The PROBE design maintains the strict randomization procedure, incorporates clearly defined (and blinded) endpoints that help eliminate bias, and allows objective comparison of therapies by an independent body.

After a 2- to 4-week, single-blind, placebo run-in period in SMOOTH, patients who met inclusion and exclusion criteria were randomized on a 1:1 basis to 4 weeks' monotherapy with either telmisartan 80 mg or valsartan 160 mg, after which all subjects received add-on HCTZ 12.5 mg for the following 6 weeks. Randomization, stratified by study centre, was by an interactive voice response system and determined using ClinPro/LBL Version 5.2 (Clinical Systems, Inc. software, USA). Patients were instructed to take study medication once daily in the morning with water (and consistently with or without food) at approximately the same time each day (0900 hours  $\pm$  1 hour).

The SMOOTH study was conducted at 118 centres in Argentina, Australia, Canada, Mexico, New Zealand, South Korea, Taiwan and the USA, following approval by the relevant local Institutional Review Board or central Independent Ethics Committee, and according to the principles of Good Clinical Practices. SMOOTH is part of the PROTECTION Programme, which is supported by Boehringer Ingelheim.



### ***Efficacy and safety evaluations***

ABPM was conducted at baseline (end of placebo run-in) and week 10 using a Spacelab Model 90207 monitor (OSI Systems Company, Issaquah, WA, USA), which was applied before the patient took the scheduled study medication. ABP was automatically measured every 20 min during the 24-hour monitoring period following dosing, with hourly means calculated relative to both the dosing time and clock time. At each scheduled visit, clinic BP was recorded using a mercury sphygmomanometer, and was defined as the mean of three seated measurements collected 2 min apart, after the patient had been seated quietly for 5 min.

The primary endpoint in the SMOOTH study was the change from baseline in mean ambulatory SBP and DBP during the last 6 hours of the 24-hour dosing interval. Secondary endpoints were changes from baseline in mean 24-hour, morning (0600 hours to noon), daytime (noon to 2200 hours), and night-time (2200 hours to 0600 hours) ambulatory SBP and DBP; changes in trough seated clinic BP; and response rates based on both 24-hour mean ambulatory SBP (24-hour mean SBP < 130 mm Hg and/or reduction  $\geq 10$  mm Hg) and DBP (24-hour mean DBP < 80 mm Hg and/or reduction  $\geq 10$  mm Hg). The incidence and severity of adverse events were monitored throughout the study.

### ***Statistics***

For the primary endpoint, assuming standard deviations of 12 and 8 mm Hg for SBP and DBP, a sample size of 337 evaluable patients per treatment

group would have had approximately 90% power to detect a treatment difference of 3 mm Hg (SBP) and 2 mm Hg (DBP) at the 5% (two-sided) level of significance.

Statistical analysis was performed on the full analysis set, which included all patients who received a dose of study medication and who had successful ABPMs at baseline and at week 10. Treatment effects were compared by analysis of covariance, with treatment regimen and centre as main effects, and the last 6-hour mean ABP at baseline as covariate.

Testing of multiple endpoints (changes in the last 6-hour mean ambulatory SBP and DBP) utilized a pre-defined hierarchical testing procedure in the following order: (1) comparison of treatment groups based on SBP; and (2) if significant, comparison based on DBP. Thus, all significance testing utilized a two-sided 5% level of significance.

## **Results**

A total of 840 patients who met inclusion and exclusion criteria were randomized to either T/HCTZ (n = 428) or V/HCTZ (n = 412). Baseline characteristics of randomized patients were comparable for the two treatment groups and are shown in Table 1. About 10% of randomized patients in each treatment group did not complete the trial, most frequently as a result of adverse events (T/HCTZ, n = 10 [2.3%]; V/HCTZ, n = 17 [4.1%]) or withdrawal of consent (T/HCTZ, n = 11 [2.6%]; V/HCTZ, n = 6 [1.5%]). The full analysis set used for the efficacy analyses included 743 patients (T/HCTZ, n = 378; V/HCTZ, n = 365) who had successful ABPM measurements at both their baseline and final visits.

### ***Reduction of ambulatory blood pressure in the last 6 hours of the 24-hour dosing interval***

After 10 weeks' treatment, T/HCTZ provided significantly greater mean BP lowering compared with V/HCTZ in the last 6 hours of the 24-hour dosing interval (Figs. 1 and 2). Mean SBP/DBP changes were -18.4/-9.7 mm Hg in T/HCTZ-treated patients and -14.5/-7.7 mm Hg in V/HCTZ-treated patients. This equates to a significant adjusted mean difference in favour of T/HCTZ of 3.9 mm Hg (95% confidence interval [CI] -5.8, -2.1) in SBP ( $p < 0.0001$ ) and 2.0 mm Hg (95% CI -3.2, -0.8) in DBP ( $p = 0.0007$ ). Overall, the mean BP was 139.1/82.4 mm Hg (V/HCTZ) and 135.6/81.2 mm Hg (T/HCTZ).

Mean BP reductions in the last 6 hours were not significantly affected by age group ( $< 65$  years vs.  $\geq 65$  years) or race. Women had significantly greater BP reductions than men; 20.6/11.3 versus 16.8/8.4 mm Hg in patients taking T/HCTZ, and 16.1/9.4 versus 13.4/6.5 mm Hg in those taking V/HCTZ. However, the difference between treatments was consistent in both genders.

### ***Secondary endpoints***

The changes from baseline in hourly means for SBP and DBP after 10 weeks of treatment with either T/HCTZ or V/HCTZ are shown in Figure 1, and the additional reductions with T/HCTZ versus V/HCTZ during various periods are depicted in Figure 2. T/HCTZ produced significantly greater reductions in 24-hour mean BP than V/HCTZ during the 24-hour dosing interval (SBP 3.0 mm Hg,  $p = 0.002$ ; DBP 1.6 mm Hg,  $p = 0.0006$ ). In addition, during the morning, daytime and night-time periods, patients receiving T/HCTZ had significantly

greater reductions in mean ambulatory SBP and DBP compared with those taking V/HCTZ ( $p < 0.003$ ) (Fig. 2). Ambulatory SBP response rates were high in patients taking T/HCTZ (79.9%) versus V/HCTZ (75.6%). Ambulatory DBP response rates were similarly high in patients taking T/HCTZ (79.9%) versus V/HCTZ (77.3%). Clinic DBP and SBP response rates were also high in patients taking T/HCTZ (80.6% and 82.3%, respectively) versus V/HCTZ (75.5% and 75.0%). Clinic BP control ( $< 130/80$  mm Hg) was higher in patients taking T/HCTZ than V/HCTZ (26.4% vs. 24.0%, respectively). Clinic DBP control was also higher in the T/HCTZ group than V/HCTZ group (77.4% vs. 71.9%, respectively).

Reductions in trough seated clinic BP after 10 weeks' treatment were significantly greater with T/HCTZ compared with V/HCTZ (SBP 3.2 mm Hg,  $p = 0.0017$ ; DBP 1.2 mm Hg;  $p = 0.0446$ ). The advantage of telmisartan was already apparent after 4 weeks of treatment with monotherapy (SBP 2.5 mm Hg;  $p = 0.0106$ ; DBP 0.8 mm Hg;  $p = 0.1370$ ). The mean BP achieved with monotherapy was 146.6/86.3 mm Hg (valsartan) and 143.3/85.5 mm Hg (telmisartan). Clinic BP control ( $< 130/80$  mm Hg) was also higher following monotherapy with telmisartan compared with valsartan (13.6 vs. 10.6%, respectively) although these values were still low. Clinic DBP control was, however, much higher following monotherapy (64.5% [telmisartan] and 63.1% [valsartan]).

### ***Safety***

The overall incidence of adverse events was low with at least one adverse event reported by 158 T/HCTZ-randomized patients (36.9%) and 151 (36.7%)

V/HCTZ- randomized patients. With the exception of dizziness, which was reported in 14 (3.3%) T/HCTZ-randomized patients and 1 (0.2%) V/HCTZ-randomized patient, there were no differences in the incidence of specific adverse events between treatment groups.

Events were generally mild or moderate in intensity. Serious adverse events were reported by three (0.7%) patients in each treatment group during monotherapy, and four (1.0%) patients (T/HCTZ) and five (1.3%) patients (V/HCTZ) during combination therapy. Adverse events reported as serious were cardiac disorders (acute coronary syndrome, unstable angina, coronary artery disease and myocardial ischaemia), gastroesophageal reflux disease, general disorders (asthenia, chest pain, cellulites), injury (ankle fracture and joint dislocation), drug toxicity, increased blood creatine phosphokinase, inadequately controlled diabetes mellitus, arthralgia, back pain, nervous system disorders (carotid artery stenosis, cerebral haemorrhage, cerebral thrombosis, headache and intracranial aneurysm), bipolar I disorder and vascular disorders (hypertensive crisis and hypotension).

Treatment-related adverse events were infrequent in the T/HCTZ (27, 6.3%) and V/HCTZ (22, 5.3%) treatment arms; the most common were oedema (0 vs. 4 [1.0%]), dizziness (5 [1.2%] vs. 0), and headache (4 [0.9%] vs. 6 [1.5%] for T/HCTZ and V/HCTZ, respectively). Laboratory markers at baseline and following T/HCTZ or V/HCTZ treatment are summarized in Table 2. The addition of HCTZ did not result in any significant metabolic disturbances.

## **Discussion**

In a large, multicentre, PROBE study of overweight/obese patients with hypertension and type 2 diabetes, once-daily T/HCTZ was significantly more effective at lowering BP than V/HCTZ during the last 6 hours of the 24-hour dosing interval. T/HCTZ also produced significantly greater BP reductions than V/HCTZ over the entire 24 hours. Both ARBs were well tolerated.

Patients enrolled in the SMOOTH study were at very high risk of cardiovascular events and stroke because of the presence of additive cardiovascular risk factors. In patients with hypertension, antihypertensive agents are recommended with a BP target of 140/90 mm Hg in patients without diabetes and 130/80 mm Hg in those with diabetes [15,16]. However, there are no specific recommendations for patients with coexisting hypertension, obesity and type 2 diabetes.

Increased activity of the RAAS is associated with obesity-linked hypertension and also with the development of type 2 diabetes. Antihypertensive agents that block the RAAS may be beneficial in patients with type 2 diabetes and features of metabolic syndrome [2]. However, there are few studies focusing on angiotensin blockade in the management of the obese hypertensive patient. ARBs and angiotensin-converting enzyme inhibitors (ACEis) may have favourable metabolic effects in addition to their antihypertensive properties that could be beneficial in these obese patients [2]. This has already been shown with valsartan and irbesartan, both in combination with HCTZ [17,18]. In another study, the Candesartan Role on Obesity and on Sympathetic System (CROSS), candesartan was found to have comparable blood pressure lowering to HCTZ, but the ARB also had a significant benefit on insulin sensitivity [19]. The current study is novel, in that

it is one of the first to directly compare ARBs in such a large obese-hypertensive population, and to monitor antihypertensive effectiveness using 24-hour ABPM.

Furthermore, our study further confirms the superiority of telmisartan versus valsartan, when both are used in fixed-dose combination with HCTZ. Although physicians are likely to use an up-titration approach to patient management, there is now evidence to suggest that a fixed-dose combination is beneficial in obese hypertensive patients, and may help to overcome the barrier of poor patient compliance as a result of polypharmacy or lack of efficacy [20]. In the current study, the mean BP achieved with monotherapy was 146.6/86.3 mm Hg (valsartan) and 143.3/85.5 mm Hg (telmisartan), which was improved to 139.1/82.4 mm Hg (valsartan) and 135.6/81.2 mm Hg (telmisartan) when the patients were switched to low-dose combination with T/HCTZ. The percentage of patients achieving target BP ( $< 130/80$  mm Hg) doubled when patients were switched to combination therapy, from 13.6% (telmisartan) to 26.4% (T/HCTZ) and from 10.6% (valsartan) to 24.0% (V/HCTZ).

Acute cardiovascular events and stroke show a strong circadian pattern. The highest incidence of events occurs in the early morning hours, during which there is typically a surge in BP [9]. Unfortunately, this vulnerable time period coincides with a decline in efficacy of most antihypertensive agents because they are relatively short acting. Differential effects between antihypertensive agents on the magnitude of reduction in the early morning hours have been shown in this and previous studies. Telmisartan has shown superiority in the early morning hours over several other antihypertensives,

such as ramipril [21–23], losartan [24,25], and valsartan [9]. In two large studies, telmisartan 80 mg changed the overall mean systolic early morning surge by -1.5 mm Hg compared with a change of + 0.3 mm Hg by the ACEi ramipril ( $p = 0.0049$ ) [6]. In the ATHOS study, telmisartan/HCTZ reduced 24-hour SBP to a greater extent than amlodipine/HCTZ [26]. ATHOS showed that antihypertensive agents with a long half-life, such as telmisartan, can maintain efficacy throughout the 24-hour dosing period, thereby providing greater reductions in early morning BP than short-acting agents [6,27]. The improved early morning hypertension control with T/HCTZ in the SMOOTH study may, therefore, be explained by the 24-hour plasma elimination half-life of telmisartan [28]. In contrast, valsartan has a shorter half-life of about 16 hours [6].

BP is strongly correlated with vascular (and overall) mortality without any evidence of a threshold down to at least 115/75 mm Hg. This was demonstrated in a meta-analysis of data from 61 prospective observational studies of patients with vascular disease [29]. In the present study, T/HCTZ provided significant additional reductions versus V/HCTZ of 3.9 mm Hg in SBP and 2.0 in DBP during the last 6 hours of the 24-hour dosing interval. Reductions of this magnitude may be clinically meaningful. For example, in the above meta-analysis, it was calculated that a reduction in SBP of 2 mm Hg was associated with a 7% reduction in risk of mortality from ischaemic heart disease and a 10% reduction in risk of mortality from stroke [29]. In another study, reduction of early morning BP in patients with type 2 diabetes, hypertension, and nephropathy was associated with a significant slowing of the progression of renal damage [30].



In conclusion, in overweight/obese patients with hypertension and type 2 diabetes, once-daily T/HCTZ was significantly more effective than V/HCTZ during the last 6 hours of the 24-hour dosing interval, the time when most cardiovascular events occur [31]. These findings indicate significant potential treatment benefit of T/HCTZ for this at-risk patient population. With strong evidence available directly correlating BP reductions to cardiovascular and cerebrovascular risk reduction, the additional efficacy afforded by T/HCTZ over V/HCTZ may confer a clinical advantage.

## Acknowledgements

The SMOOTH study was supported by a grant from Boehringer Ingelheim. Previously published in abstract form [32]. The Authors would like to acknowledge the following SMOOTH Investigators for their participation in terms of patient enrollment and provision of data:

**Argentina:** Mario Bendersky, Alberto Caccavo, Carlos Cuneo, Alfredo Hirschson, Oscar Motaña, Ernesto Paolasso, Ramiro Sanchez.

**Australia:** Ajith Dissanayake, Garry Jennings, Jeff Karrasch, Andrew Lang, Russell Scott.

**Canada:** Carl Abbott, Ronnie Aronson, Ellen Burgess, Brian Carlson, Raja Cheyhab, Howard Conter, Geoff Davis, Jiri Frohlich, Daniel Gaudet, Arvinder Grover, Pavel Hamet, Evan Howlett, Ivan Jagas, Yves Lacourcière, Arun Nayar, Dennis O'Keefe, Michael O'Mahony, Akbar Panju, Robert Petrella, Calvin Powell, Robert Schlosser, Rizwan Somani, David Spence, Kim-Weng Tan, Ivor Teitelbaum, Hugh Tildesley, Robert Ting, Yaw Twum-Barima, Richard Tytus, Tom Wilson.

**Korea:** Dong-Seop Choi, Hyoung Woo Lee, Yang Wook Kim.

**Mexico:** Héctor Briseño, Ernesto Cardona Muñoz, Enrique Gómez Alvarez, Rogelio Heredia, Norberto Hernández, Jaime Illescas Díaz, Margarita López, Eduardo Mendiola, Rafael Olvera Ruiz, Eduardo Quintero.

**Taiwan:** Tien-Shang Huang.

**USA:** Dwayne Aboud, M Angeli Adamczyk, Luis Angles, Najam Awan, Michael Azorr, George Bakris, Kenneth Blaze, Samuel Blumenthal, David Calhoun, Vito Campese, DeAnna Cheung, Marc Comianos, Rafael Coplin, Steven Chrysant, Leo Egbujiobi, Brian Ellis, John Ervin, James Gavin III,

Richard Gerber, Larry Gilderman, Mark Gotfried, Alan Graff, Mark Graves,  
James Gray, James Gutmann, Gordy Hall, David Hassman, James Herron,  
Barry Horowitz, Timothy Howard, Joseph Izzo Jr, Anna Jackson, Spencer  
Jones, H Kennedy Cathcart, Lawrence Koehler, Andrew Lewin, Thomas  
Littlejohn III, Martha Logan, Robert Medalie, Scott Meyers, John Murray, Joel  
Neutel, Alan Niederman, David Pate, Andres Patron, Gary Plundo, Rakesh  
Prasad, Henry Punzi, Amin Radparvar, Marina Raikhel, Michael Rubenstein,  
Mercedes Samson, Sherwyn Schwartz, Gregory T Serfer, Shaukat Shah, Lok  
Shandilya, Stanley Sharp, Kevin Sheehan, Marc Shepard, William Smith,  
Ronald Sockolov, Dave E Webster, Nathaniel Winer.

## References

1. Rahmouni K, Correia ML, Haynes WG, Mark AL: **Obesity-associated hypertension: new insights into mechanisms.** *Hypertension* 2005, **45**:9-14.
2. Sharma AM: **Is there a rationale for angiotensin blockade in the management of obesity hypertension?** *Hypertension* 2004, **44**:12-19.
3. Bramlage P, Pittrow D, Wittchen HU, Kirch W, Boehler S, Lehnert H, Hoefler M, Unger T, Sharma AM: **Hypertension in overweight and obese primary care patients is highly prevalent and poorly controlled.** *Am J Hypertens* 2004, **17**:904-910.
4. Marler JR, Price TR, Clark GL, Muller JE, Robertson T, Mohr JP, Hier DB, Wolf PA, Caplan LR, Foulkes MA: **Morning increase in onset of ischemic stroke.** *Stroke* 1989, **20**:473-476.
5. Kario K: **Time for focus on morning hypertension: pitfall of current antihypertensive medication.** *Am J Hypertens* 2005, **18**:149-151.
6. Burnier M, Maillard M: **The comparative pharmacology of angiotensin II receptor antagonists.** *Blood Press* 2001, **10**(Suppl 1):6-11.
7. Kakuta H, Sudoh K, Sasmata M, Yamagishi S: **Telmisartan has the strongest binding affinity to angiotensin II type 1 receptor: comparison with other angiotensin II type 1 receptor blockers.** *Int J Clin Pharm Res* 2005, **25**:41-46..
8. Wienen W, Entzeroth M, van Meel JCA, Stangier J, Busch U, Ebner T, Schmid J, Lehmann H, Matzek K, Kempthorne-Rawson J, Gladigau V, Haezel HH: **A review on telmisartan: a novel, long-acting angiotensin II-receptor antagonist.** *Cardiovasc Drug Rev*, 2002, **18**:127-156.
9. Gosse P, Neutel JM, Schumacher H, Lacourciere Y, Williams B, Davidai G: **The effect of telmisartan and ramipril on early morning blood pressure surge: a pooled analysis of two randomized clinical trials.** *Blood Press Monit* 2007, **12**:141-147.
10. Fogari R, Derosa G, Zoppi A, Rinaldi A, Lazzari P, Fogari E, Mugellini A, Preti P: **Comparison of the effects of valsartan and felodipine on plasma leptin and insulin sensitivity in hypertensive obese patients.** *Hypertens Res* 2005, **28**:209-214.
11. Lacourcière Y, Krzesinski JM, White WB, Davidai G, Schumacher H: **Sustained antihypertensive activity of telmisartan compared with valsartan.** *Blood Press Monit* 2004, **9**:203-210.
12. White WB, Punzi HA, Murwin D, Koval SE, Davidai G, Neutel JM: **Effects of the angiotensin II receptor blockers telmisartan vs**

- valsartan in combination with hydrochlorothiazide 25 mg once daily for the treatment of hypertension.** *J Clin Hypertens* 2006, **8**:626-633.
13. White WB, Guthrie R, Chrysant SG, Murwin D, Davidai G, Koval S: **A second large trial of Telmisartan and HCTZ (80/25 mg) shows a larger antihypertensive effect than Valsartan and HCTZ (160/25 mg).** *J Clin Hypertens* 2007, **9**(Suppl. A):A44 (Abstract P-94).
  14. Smith DH, Neutel JM, Lacourciere Y, Kempthorne-Rawson J: **Prospective, randomized, open-label, blinded-endpoint (PROBE) designed trials yield the same results as double-blind, placebo-controlled trials with respect to ABPM measurements.** *J Hypertens* 2003, **21**:1291-1298.
  15. American Diabetes Association: **Treatment of hypertension in adults with diabetes.** *Diabet Care* 2002, **25**:199-201.
  16. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jr., Jones DW, Materson BJ, Oparil S, Wright JT, Jr., Roccella EJ: **The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure.** *JAMA* 2003, **289**:2560-2572.
  17. Sharma AM, Bramlage P, Kirch W: **Antihypertensive effect of irbesartan and predictors of response in obesity-associated hypertension: a prospective, open-label study.** *Clin Drug Investig* 2005, **25**:765-776.
  18. Jordan J, Engeli S, Boschmann M, Weidinger G, Luft FC, Sharma AM, Kreuzberg U: **Hemodynamic and metabolic responses to valsartan and atenolol in obese hypertensive patients.** *J Hypertens* 2005, **23**:2313-2318.
  19. Grassi G, Seravalle G, Dell'Oro R, Trevano FQ, Bombelli M, Scopelliti F, Facchini A, Mancia G: **Comparative effects of candesartan and hydrochlorothiazide on blood pressure, insulin sensitivity, and sympathetic drive in obese hypertensive individuals: results from the CROSS study.** *J Hypertens* 2003, **21**:1761-1769.
  20. Bangalore S, Shahane A, Parkar S, Messerli FH: **Compliance and fixed-dose combination therapy.** *Curr Hypertens Rep* 2007, **9**:184-189.
  21. Lacourcière Y, Neutel JM, Davidai G, Koval S: **A multicenter, 14-week study of telmisartan and ramipril in patients with mild-to-moderate hypertension using ambulatory blood pressure monitoring.** *Am J Hypertens* 2006, **19**:104-112.
  22. Poirier L, de Champlain J, Larochelle P, Lamarre-Cliche M, Lacourciere Y: **A comparison of the efficacy and duration of action of**

- telmisartan, amlodipine and ramipril in patients with confirmed ambulatory hypertension.** *Blood Press Monit* 2004, **9**:231-236.
23. Williams B, Gosse P, Lowe L, Harper R, on behalf of the PRISMA I Study Group: **The prospective, randomized investigation of the safety and efficacy of telmisartan versus ramipril using ambulatory blood pressure monitoring (PRISMA I).** *J Hypertens* 2006, **24**:193-200.
  24. Lacourcière Y, Neutel JM, Schumacher H: **Comparison of fixed-dose combinations of telmisartan/hydrochlorothiazide 40/12.5 mg and 80/12.5 mg and a fixed-dose combination of losartan/hydrochlorothiazide 50/12.5 mg in mild to moderate essential hypertension: Pooled analysis of two multicenter, prospective, randomized, open-label, blinded-end point (PROBE) trials.** *Clin Ther* 2005, **27**:1795-1805.
  25. Smith DH, Cramer MJ, Neutel JM, Hettiarachchi R, Koval S: **Comparison of telmisartan versus losartan: meta-analysis of titration-to-response studies.** *Blood Press Monit* 2003, **8**:111-117.
  26. Neldam S, Edwards C; ATHOS Study Group: **Telmisartan plus HCTZ vs. amlodipine plus HCTZ in older patients with systolic hypertension: results from a large ambulatory blood pressure monitoring study.** *Am J Geriatr Cardiol* 2006, **15**:151-160.
  27. Elliott HL, Meredith PA: **Trough:peak ratio: clinically useful or practically irrelevant?** *J Hypertens* 1995, **13**:279-283.
  28. Stangier J, Su CAPF, Roth W: **Pharmacokinetics of orally and intravenously administered telmisartan in healthy young and elderly volunteers and in hypertensive patients.** *J Int Med Res* 2000, **28**:149-167.
  29. Lewington S, Clarke R, Qizilbash N, Peto R, Collins R: **Age-specific relevance of usual blood pressure to vascular mortality: a meta-analysis of individual data for one million adults in 61 prospective studies.** *Lancet* 2002, **360**:1903-1913.
  30. Suzuki H, Kanno Y, Nakamoto H, Okada H, Sugahara S: **Decline of renal function is associated with proteinuria and systolic blood pressure in the morning in diabetic nephropathy.** *Clin Exp Hypertens* 2005, **27**:129-138.
  31. Elliott WJ: **Cyclic and circadian variations in cardiovascular events.** *Am J Hypertens* 2001, **14**:291S-295S.
  32. Sharma AM, Davidson JA, Gavin III JR, DeSousa NJ: **Comparison of the antihypertensive efficacy of telmisartan/hydrochlorothiazide vs valsartan/hydrochlorothiazide in high-risk overweight/obese**

**patients with hypertension and type 2 diabetes.** *Hypertension* 2005, 46:898-899.

## Figure legends

**Figure 1:** Changes from baseline in hourly means for (A) SBP and (B) DBP after treatment with T/HCTZ or V/HCTZ at 10 weeks. Squares: T/HCTZ; triangles: V/HCTZ.

**Figure 2:** Additional reductions with T/HCTZ compared with V/HCTZ in mean ambulatory SBP and DBP during various periods of the 24-hour dosing interval after treatment for 10 weeks. Black: SBP; white: DBP.



**Table 1: Patient demographics and baseline characteristics of randomized patients.**

|                                    | <b>T/HCTZ</b><br><b>(n = 428)</b> | <b>V/HCTZ</b><br><b>(n = 412)</b> |
|------------------------------------|-----------------------------------|-----------------------------------|
| Gender (n [%])                     |                                   |                                   |
| Men                                | 230 (53.7)                        | 239 (58.0)                        |
| Women                              | 198 (46.3)                        | 173 (42.0)                        |
| Age                                |                                   |                                   |
| Mean $\pm$ SD (years)              | 58.5 $\pm$ 9.7                    | 59.2 $\pm$ 9.7                    |
| $\geq 65$ years (n [%])            | 122 (28.5)                        | 119 (28.9)                        |
| Race (n [%])                       |                                   |                                   |
| White                              | 292 (68.2)                        | 292 (70.9)                        |
| Black                              | 61 (14.3)                         | 59 (14.3)                         |
| Asian                              | 75 (17.5)                         | 61 (14.8)                         |
| BMI                                |                                   |                                   |
| Mean $\pm$ SD (kg/m <sup>2</sup> ) | 33.6 $\pm$ 5.7                    | 33.4 $\pm$ 5.3                    |
| Hypertension duration              |                                   |                                   |
| Mean $\pm$ SD (years)              | 9.2 $\pm$ 9.5                     | 8.9 $\pm$ 9.2                     |
| 24-hour ABP                        |                                   |                                   |
| Mean SBP $\pm$ SD (mm Hg)          | 148.0 $\pm$ 11.9                  | 147.8 $\pm$ 12.4                  |
| Mean DBP $\pm$ SD (mm Hg)          | 83.1 $\pm$ 9.2                    | 83.3 $\pm$ 10.2                   |
| Trough seated BP                   |                                   |                                   |
| Mean SBP $\pm$ SD (mm Hg)          | 156.8 $\pm$ 12.5                  | 157.1 $\pm$ 12.3                  |
| Mean DBP $\pm$ SD (mm Hg)          | 92.1 $\pm$ 9.3                    | 91.7 $\pm$ 9.8                    |
| HbA1c (%)                          |                                   |                                   |
| Mean $\pm$ SD                      | 7.0 $\pm$ 1.1                     | 7.2 $\pm$ 1.2                     |

ABP: ambulatory blood pressure; DBP: diastolic blood pressure; T/HCTZ: telmisartan 80 mg plus HCTZ 12.5 mg; V/HCTZ: valsartan 160 mg plus HCTZ 12.5 mg; SBP: systolic blood pressure.

**Table 2: Summary of metabolic markers at baseline and on the last day of combination therapy.**

| Endpoint, mean $\pm$ SD    | T/HCTZ           |                  | V/HCTZ           |                  | Difference*<br>(T/HCTZ - V/HCTZ) |
|----------------------------|------------------|------------------|------------------|------------------|----------------------------------|
|                            | Baseline         | Last day         | Baseline         | Last day         | p value                          |
| Serum triglyceride (mg/dL) | 201.7 $\pm$ 247  | 211.8 $\pm$ 172  | 196.3 $\pm$ 151  | 210.5 $\pm$ 177  | 0.91                             |
| LDL-C (mg/dL)              | 165.6 $\pm$ 35.4 | 169.8 $\pm$ 38.1 | 162.8 $\pm$ 33.9 | 165.3 $\pm$ 36.0 | 0.17                             |
| HDL-C (mg/dL)              | 37.1 $\pm$ 6.5   | 36.6 $\pm$ 6.3   | 37.6 $\pm$ 7.2   | 37.1 $\pm$ 7.0   | 0.64                             |
| Total cholesterol (mg/dL)  | 195.7 $\pm$ 23.3 | 198.5 $\pm$ 21.7 | 194.5 $\pm$ 19.8 | 196.3 $\pm$ 20.9 | 0.20                             |
| Potassium (mEq/L)          | 4.4 $\pm$ 0.4    | 4.5 $\pm$ 0.4    | 4.4 $\pm$ 0.4    | 4.4 $\pm$ 0.4    | 0.03                             |
| Glucose (mg/dL)            | 128.8 $\pm$ 44.1 | 138.8 $\pm$ 53.1 | 134.1 $\pm$ 49.2 | 140.4 $\pm$ 55.2 | 0.58                             |
| HbA <sub>1c</sub> (%)      | 6.5 $\pm$ 0.3    | 6.6 $\pm$ 0.3    | 6.6 $\pm$ 0.3    | 6.7 $\pm$ 0.4    | 0.93                             |
| Albumin:creatinine ratio   | 97.6 $\pm$ 330   | 65.1 $\pm$ 266   | 66.7 $\pm$ 239   | 45.1 $\pm$ 177   | 0.97                             |


T/HCTZ: telmisartan 80 mg plus HCTZ 12.5 mg; V/HCTZ: valsartan 160 mg plus HCTZ 12.5

mg; LDL-C: low-density lipoprotein cholesterol; HDL-C: high-density lipoprotein cholesterol;

HbA<sub>1c</sub>: glycolated haemoglobin.

\*Analysis of change from baseline

A



B




Figure 1




Figure 2 \*p < 0.05; †p < 0.001; ‡p < 0.0001 for T/HCTZ vs. V/HCTZ