

```
In [1]: import numpy as np
import pandas as pd
import matplotlib.pyplot as plt
import seaborn as sns
import scipy
from scipy import stats
import statsmodels.api as sm
import pylab
import scipy.stats as stats

In [2]: cutlets = pd.read_csv ('/Users/SAURABH/Saurabh patil/DATA SCIENCE/Hypothesis/cutlets.csv')
```

```
In [3]: cutlets
```

Out[3]:		Unit A	Unit B
	0	6.8090	6.7703
	1	6.4376	7.5093
	2	6.9157	6.7300
	3	7.3012	6.7878
	4	7.4488	7.1522
	5	7.3871	6.8110
	6	6.8755	7.2212
	7	7.0621	6.6606
	8	6.6840	7.2402
	9	6.8236	7.0503
	10	7.3930	6.8810
	11	7.5169	7.4059
	12	6.9246	6.7652
	13	6.9256	6.0380
	14	6.5797	7.1581
	15	6.8394	7.0240
	16	6.5970	6.6672
	17	7.2705	7.4314
	18	7.2828	7.3070
	19	7.3495	6.7478
	20	6.9438	6.8889
	21	7.1560	7.4220
	22	6.5341	6.5217
	23	7.2854	7.1688
	24	6.9952	6.7594
	25	6.8568	6.9399
	26	7.2163	7.0133
	27	6.6801	6.9182
	28	6.9431	6.3346
	29	7.0852	7.5459
	30	6.7794	7.0992
	31	7.2783	7.1180
	32	7.1561	6.6965
	33	7.3943	6.5780
	34	6.9405	7.3875

```
In [4]: cutlets.shape
```

Out[4]: (35, 2)

```
In [5]: #Since the Unit A and Unit B are two different samples put together in a single dataset, hence we'll split them
#into two separate datasets
```

```
In [6]: unitA = cutlets['Unit A']
```

```
In [7]: unitB = cutlets['Unit B']
```

```
In [8]: unitA.describe()
```


Out[8]:	count	35.000000
	mean	7.019091
	std	0.288408
	min	6.437600
	25%	6.831500
	50%	6.943800
	75%	7.280550
	max	7.516900
	Name: Unit A, dtype: float64	

```
In [9]: unitB.describe()
```


Out[9]:	count	35.000000
	mean	6.964297
	std	0.343401
	min	6.038000
	25%	6.753600
	50%	6.939900
	75%	7.195000
	max	7.545900
	Name: Unit B, dtype: float64	

```
In [10]: #Checking if the samples are normally distributed
```

```
In [11]: measurements = np.random.normal(loc = 7.019091, scale =0.288408 , size=35)
stats.probplot(measurements, dist="norm", plot=pylab)
pylab.show()
```



```
In [12]: measurements = np.random.normal(loc = 6.964297, scale = 0.343401 , size=35)
stats.probplot(measurements, dist="norm", plot=pylab)
pylab.show()
```



```
In [13]: #By looking at their qqplots, it is evident that the samples are approximately normally distributed
```

```
In [14]: #Hence, now we'll proceed for sample t- test is equality of Means test. Sample Mean will tell us which program is better.
#Ho= Averages of diameters of Unit A is equal to Averages of diameters of unit B
#Ha= Averages of diameters of Unit A is not equal to Averages of diameters of unit B
```

```
In [15]: stats.ttest_ind (unitA, unitB)
```

Out[15]: Ttest\_indResult(statistic=0.7228688704678061, pvalue=0.4722394724599501)

```
In [16]: #Since pvalue(=0.47) > alpha(=0.05), hence we can't reject the null hypothesis
```

```
In [17]: #Conclusion: there is no significant difference in the diameters of Unit A and Unit B at 5% significance level
```

```
In [ ]:
```