
Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.125
0.100
0.075
0.050
0.025
0.000
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.125
0.100
0.075
0.050
0.025
0.000
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.25
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
10.0
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
3
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
5
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
3
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
3
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
5
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
3
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

505


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

542


5.050
5.025
5.000
4.975
4.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


1
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


0.6
0.4
0.2
0.0
Mean root−to−tip−distance (read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


0.75
0.50
0.25
0.00
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads in different subraphs
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

550


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
8
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1

557


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

572


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
10.0
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
3
2
1

603


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance (read−weighted)
0.5
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
4
3
2
1

622


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
10.0
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
3
2
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

720


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
9
6
3
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
3
2
1

792


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

804


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance (read−weighted)
0.100
0.075
0.050
0.025
0.000
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

883


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

902


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance (read−weighted)
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

928


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

940


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1

944


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

985


Tip or read count
5.050
5.025
5.000
4.975
4.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph

Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1