
Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
8
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
0
1
2
3
4
5
6
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0
3
6
9
Proportion of reads in different subraphs
0.00
0.25
0.50
0.75
1.00
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.2
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
5
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
3
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.25
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


2
1
0
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


15
10
5
0
Mean root−to−tip−distance
(read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


20
15
10
5
0
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads
in different subraphs
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


6
4
2
0
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


1.5
1.0
0.5
0.0
Mean root−to−tip−distance
(read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


2.5
2.0
1.5
1.0
0.5
0.0
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads
in different subraphs
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
10
8
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


1
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


2
1
0
Mean root−to−tip−distance
(read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


4
3
2
1
0
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads
in different subraphs
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


1
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


0.5
0.4
0.3
0.2
0.1
0.0
Mean root−to−tip−distance (read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


0.6
0.4
0.2
0.0
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads in different subraphs
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


1
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


1.0
0.5
0.0
Mean root−to−tip−distance
(read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.5
1.0
0.5
0.0
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads
in different subraphs
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
5
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
4
3
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
15
10
5
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
15
10
5
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


1
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


0.06
0.04
0.02
0.00
Mean root−to−tip−distance
(read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


0.075
0.050
0.025
0.000
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads
in different subraphs
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
10.0
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
12
8
4
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
10
8
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
10
5
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


5
4
3
2
1
0
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


2.0
1.5
1.0
0.5
0.0
Mean root−to−tip−distance
(read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


3
2
1
0
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads
in different subraphs
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

355


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
8
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.075
0.050
0.025
0.000
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.09
0.06
0.03
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.12
0.09
0.06
0.03
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

502


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

550


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
5
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

552


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.5
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

625


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
8
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

652


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
9
6
3
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
8
6
4
2
0
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
Tip set
All
Largest subgraph
Mean pairwise patristic distance
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1

705


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

730


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.04
0.03
0.02
0.01
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.04
0.03
0.02
0.01
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


1
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


1.5
1.0
0.5
0.0
Mean root−to−tip−distance
(read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.5
1.0
0.5
0.0
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads
in different subraphs
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

828


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

892


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
5
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1

895


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1

901


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.5
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1

962


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.5
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1

974


10.050
10.025
10.000
9.975
9.950
Tip or read count
3000
6000
9000
Window centre
Variable
Reads
Tips


1
Subgraph or clade count
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


0.4
0.3
0.2
0.1
0.0
Mean root−to−tip−distance
(read−weighted)
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


0.6
0.4
0.2
0.0
Mean pairwise patristic distance
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


1.00
0.75
0.50
0.25
0.00
Proportion of reads
in different subraphs
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1