
Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance
(read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.2
0.8
0.4
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.2
0.8
0.4
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
5
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
8
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.2
0.8
0.4
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1

202


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
10.0
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
8
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.09
0.06
0.03
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.08
0.06
0.04
0.02
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.075
0.050
0.025
0.000
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
8
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
10
5
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.100
0.075
0.050
0.025
0.000
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
12
8
4
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
2.0
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.075
0.050
0.025
0.000
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.09
0.06
0.03
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.8
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.20
0.15
0.10
0.05
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
4
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.6
0.4
0.2
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
1.2
0.9
0.6
0.3
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
4
3
2
1
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
8
6
4
2
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance (read−weighted)
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.5
1.0
0.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0.4
0.3
0.2
0.1
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips
Subgraph or clade count
3
2
1
0
3000
6000
9000
Window centre
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
6
4
2
0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Mean pairwise patristic distance
7.5
5.0
2.5
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
3000
6000
9000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
3000
6000
9000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance (read−weighted)
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
1.2
0.8
0.4
0.0
3000
6000
9000
Window centre
Tip set
All
Largest subgraph


Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
3000
6000
9000
Window centre
Subgraph rank (by tip count)
1