
Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
6
4
2
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
0
1
2
Tip set
All
Largest subgraph
Mean pairwise patristic distance
0
1
2
3
4
Proportion of reads in different subraphs
0.00
0.25
0.50
0.75
1.00
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
4
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
6
4
2
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
1

532


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
4
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
6
4
2
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips
Subgraph or clade count
1
Variable
Clades
Subgraphs
Mean root−to−tip−distance (read−weighted)
2.0
1.5
1.0
0.5
0.0
Tip set
All
Largest subgraph
Mean pairwise patristic distance
3
2
1
0
Proportion of reads in different subraphs
1.00
0.75
0.50
0.25
0.00
Subgraph rank (by tip count)
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
2
1
0
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
10
5
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
12
9
6
3
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
4
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
12
9
6
3
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
15
10
5
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
4
3
2
1

711


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
6
5
4
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
6
4
2
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
10.0
7.5
5.0
2.5
0.0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
3
2
1

783


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
8
6
4
2
0
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
0.3
0.2
0.1
0.0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
0.20
0.15
0.10
0.05
0.00
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
2
1

862


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
3.0
2.5
2.0
1.5
1.0
0.5
0.0
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
12
8
4
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
15
10
5
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
3
2
1


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
1
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
4
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
1

971


Tip or read count
10.050
10.025
10.000
9.975
9.950
1000
1250
1500
1750
2000
Window centre
Variable
Reads
Tips


Subgraph or clade count
4
3
2
1
0
1000
1250
1500
1750
2000
Window centre
Variable
Clades
Subgraphs


Mean root−to−tip−distance
(read−weighted)
10.0
7.5
5.0
2.5
0.0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Mean pairwise patristic distance
15
10
5
0
1000
1250
1500
1750
2000
Window centre
Tip set
All
Largest subgraph


Proportion of reads
in different subraphs
1.00
0.75
0.50
0.25
0.00
1000
1250
1500
1750
2000
Window centre
Subgraph rank
(by tip count)
4
3
2
1