

# Line Search Methods

Saurav Samantaray

Department of Mathematics

Indian Institute of Technology Madras

January 26, 2024



# Line Search Method

- In each iteration of a line search method a search direction  $p_k$  is computed, and
- then its decided how far to move along that direction.
- An iteration is given by

$$x_{k+1} = x_k + \alpha_k p_k \quad (1)$$

where  $\alpha_k > 0$  (scalar) called the step length.

The success of a line search method depends on effective choices of both:

- 1 the direction  $p_k$
- 2 the step length  $\alpha_k$

# The Steepest Descent Direction

- The steepest descent direction  $-\nabla f_k$  is the most obvious choice for search direction for a line search method.
- Among all Directions one could form from  $x_k$ , along  $-\nabla f_k$ ,  $f$  decreases most rapidly.

## Justification

- Consider any search direction  $p$  and step-length  $\alpha$ , we have

$$f(x_k + \alpha p) = f(x_k) + \alpha p^T \nabla f_k + \frac{1}{2} \alpha^2 p^T \nabla^2 f(x_k + tp) p \text{ for some } t \in (0, 1)$$

- Let  $\alpha \ll 1$  (small) and we consider the first-order approximation of  $f$  at  $x_k + \alpha p$  around  $x_k$  as:

$$f(x_k + \alpha p) \approx f(x_k) + \alpha p^T \nabla f_k$$

- Change in  $f$  moving from  $x_k$  to  $x_k + \alpha p$  is

$$f(x_k + \alpha p) - f(x_k)$$

# The Steepest Descent Direction

- As the distance moved in the direction is  $\alpha$ , therefore, the rate of change of  $f$  along the direction  $p$  at  $x_k$  is

$$\frac{f(x_k + \alpha p) - f(x_k)}{\alpha}$$

- which is coefficient of  $\alpha$ , i.e.

$$p^T \nabla f_k$$

- This implies smaller the above value is, more descent can be achieved.
- Hence, the unit direction  $p$  of most rapid decrease is the solution to the problem

$$\min_p p^T \nabla f_k, \quad \text{subject to } \|p\| = 1.$$

# The Steepest Descent Direction



$$p^T \nabla f_k = \|p\| \|\nabla f_k\| \cos \theta = \|\nabla f_k\| \cos \theta$$

where  $\theta$  is the angle between  $p$  and  $\nabla f_k$ .

- The minimiser is attained when  $\cos \theta = -1$  and

$$p = -\frac{\nabla f_k}{\|\nabla f_k\|}$$




Figure illustrates this direction is orthogonal to the contours of the function

# The Steepest Descent Direction

- At every step (iteration) in the steepest descent method the search direction is chosen along

$$p = -\nabla f_k$$

- $\alpha_k$  can be chosen in a variety of ways.
- One advantage of this method is it requires only the calculation of gradient ( $\nabla f_k$ ), but not second derivatives.
- Line search methods may use search directions other than the steepest descent direction.

# Descent Direction

## Descent Direction

Any direction that makes an **angle of strictly less than  $\frac{\pi}{2}$**  radians with  $\nabla f_k$  is guaranteed to produce a decrease in  $f$ , provided the step length is sufficiently small and is called a **descent direction**.

- Now consider

$$f(x_k + \epsilon p_k) = f(x_k) + \epsilon p_k^T \nabla f_k + \mathcal{O}(\epsilon^2).$$

- When  $p_k$  is a downhill (descent) direction, the angle  $\theta_k$  between  $p_k$  and  $\nabla f_k$  has  $\cos \theta_k < 0$ , so that

$$\begin{aligned} p_k^T \nabla f_k &= \|p_k\| \|\nabla f_k\| \cos \theta_k < 0 \\ \implies f(x_k + \epsilon p_k) &< f(x_k) \end{aligned}$$

- Most line search algorithms require  $p_k$  to be descent direction, because this property guarantees that the function  $f$  can be reduced along this direction.

# Newton Search Direction

- This direction is derived from the second-order Taylor series approximation to  $f(x_k + p)$ , which is

$$f(x_{k+p}) \approx f_k + p \nabla f_k + \frac{1}{2} p \nabla^2 f_k p \stackrel{\text{def}}{=} m_k(p). \quad (2)$$

- Assuming for the moment that  $\nabla^2 f_k$  is positive definite, we obtain the Newton direction by finding the vector  $p$  that minimizes  $m_k(p)$ .
- By simply setting the derivative of  $m_k(p)$  to zero, we obtain the following explicit formula:

$$p_k^N = -(\nabla^2 f_k)^{-1} \nabla f_k. \quad (3)$$

- The Newton direction is reliable when the difference between the true function  $f(x_k + p)$  and its quadratic model  $m_k(p)$  is not too large.



# Newton Search Direction

- If  $\nabla^2 f$  is sufficiently smooth, this difference introduces a perturbation of only  $\mathcal{O}(\|p\|^3)$ .
- Therefore, when  $\|p\|$  is small

$$f(x_k + p) \approx m_k(p) \quad \text{quite accurately}$$

- The Newton direction can be used in a line search method when  $\nabla^2 f_k$  is positive definite, as in this case we have

$$\begin{aligned} \nabla f_k^T p_k^N &= -p_k^{N^T} \nabla^2 f_k p_k^N \\ &< -\sigma_k \|p_k^N\|^2 \end{aligned}$$

for some  $\sigma_k > 0$  (+ve definiteness of  $\nabla^2 f_k$ )

- Unless the gradient  $\nabla f_k$  (and therefore the step  $p_k^N$ ) is zero, we have

$$\nabla f^T p_k^N < 0$$

- so the Newton direction is a descent direction.

# Newton Search Direction

- Unlike the steepest descent direction, there is a “natural” step length of 1 associated with the Newton direction.
- Adjust  $\alpha$  only when it does not produce a satisfactory reduction in the value of  $f$ .
- Note that when  $\nabla_k^f$  is not positive definite the Newton direction may not exist, since  $(\nabla^2 f_k)^{-1}$  may not exist.
- Even when it is defined, it may not satisfy the descent property, and therefore is unsuitable.
- Methods that use Newton direction have fast rate of local convergence (more on this later).
- After a neighbourhood of the solution is reached, convergence to high accuracy often occurs in just a few iterations.

# Newton Search Direction

- The main drawback is the need to calculate the Hessian  $\nabla^2 f(x)$ .
- Explicit computation of this matrix of second derivatives can sometimes be a cumbersome, error-prone, and expensive process.
- Finite-difference and automatic differentiation techniques come useful in avoiding the need to calculate second derivatives by hand.