

SBML Model Report

Model name: “Waugh2006 - Diabetic Wound Healing - TGF-B Dynamics”



May 17, 2018

1 General Overview

This is a document in SBML Level 2 Version 4 format. This model was created by the following two authors: Matthew Grant Roberts¹ and Catherine Lloyd² at June 25th 2010 at 12:01 a. m. and last time modified at March first 2018 at 12:58 a. m. Table 1 gives an overview of the quantities of all components of this model.

Table 1: Number of components in this model, which are described in the following sections.

Element	Quantity	Element	Quantity
compartment types	0	compartments	1
species types	0	species	4
events	0	constraints	0
reactions	0	function definitions	0
global parameters	11	unit definitions	9
rules	4	initial assignments	0

Model Notes

This a model from the article:

Macrophage dynamics in diabetic wound dealing.

Waugh HV, Sherratt JA. Bull Math Biol 2006 Jan;68(1):197-207 [16794927](#) ,

Abstract:

¹EMBL-EBI, mroberts@ebi.ac.uk

²University of Auckland, c.lloyd@auckland.ac.nz

Wound healing in diabetes is a complex process, characterised by a chronic inflammation phase. The exact mechanism by which this occurs is not fully understood, and whilst several treatments for healing diabetic wounds exist, very little research has been conducted towards the causes of the extended inflammation phase. We describe a mathematical model which offers a possible explanation for diabetic wound healing in terms of the distribution of macrophage phenotypes being altered in the diabetic patient compared to normal wound repair. As a consequence of this, we put forward a suggestion for treatment based on rectifying the macrophage phenotype imbalance.

This model was taken from the [CellML repository](#) and automatically converted to SBML.

The original model was: [Waugh HV, Sherratt JA. \(2006\) - version=1.0](#)

The original CellML model was created by:

Catherine Lloyd

c.lloyd@auckland.ac.nz

The University of Auckland

This model originates from BioModels Database: A Database of Annotated Published Models (<http://www.ebi.ac.uk/biomodels/>). It is copyright (c) 2005-2011 The BioModels.net Team.

To the extent possible under law, all copyright and related or neighbouring rights to this encoded model have been dedicated to the public domain worldwide. Please refer to [CC0 Public Domain Dedication](#) for more information.

In summary, you are entitled to use this encoded model in absolutely any manner you deem suitable, verbatim, or with modification, alone or embedded it in a larger context, redistribute it, commercially or not, in a restricted way or not..

To cite BioModels Database, please use: [Li C, Donizelli M, Rodriguez N, Dharuri H, Endler L, Chelliah V, Li L, He E, Henry A, Stefan MI, Snoep JL, Hucka M, Le Novre N, Laibe C \(2010\) BioModels Database: An enhanced, curated and annotated resource for published quantitative kinetic models. BMC Syst Biol., 4:92.](#)

2 Unit Definitions

This is an overview of 13 unit definitions of which four are predefined by SBML and not mentioned in the model.

2.1 Unit time

Name time

Definition 86400 s

2.2 Unit unit_0

Name $8.64\text{e-}11\text{m}\hat{6}/(\text{g}\cdot\text{s})$

Definition $(8.64 \cdot 10^{-11} \text{ m})^6 \cdot \text{g}^{-3} \cdot \text{s}^{-1}$

2.3 Unit `unit_1`

Name $0.0864 \cdot \text{mm}/(\text{g} \cdot \text{s})$

Definition $(0.0864 \text{ mm})^3 \cdot \text{g}^{-2} \cdot \text{s}^{-1}$

2.4 Unit `unit_2`

Name $1/(11.5741 \cdot \text{Mg} \cdot \text{s})$

Definition $(11.5741 \text{ Mg})^{-1} \cdot \text{s}^{-1}$

2.5 Unit `unit_3`

Name $1/(0.0115741 \cdot \text{m} \cdot \text{s})$

Definition $(0.0115741 \text{ m})^{-3} \cdot \text{s}^{-1}$

2.6 Unit `unit_4`

Name 1

Definition dimensionless⁰

2.7 Unit `unit_5`

Name $1/(0.0115741 \cdot \text{ms})$

Definition $(0.0115741 \text{ ms})^{-1}$

2.8 Unit `unit_6`

Name $0.001 \cdot \text{m}$

Definition $(0.0010 \text{ m})^3$

2.9 Unit `unit_7`

Name $0.0864 \cdot \text{g}/\text{s}$

Definition $(0.0864 \cdot 10^{-6} \text{ dimensionless})^0 \cdot 0.0864 \text{ g} \cdot \text{s}^{-1}$

2.10 Unit `substance`

Notes Mole is the predefined SBML unit for substance.

Definition mol

2.11 Unit volume

Notes Litre is the predefined SBML unit for volume.

Definition 1

2.12 Unit area

Notes Square metre is the predefined SBML unit for area since SBML Level 2 Version 1.

Definition m^2

2.13 Unit length

Notes Metre is the predefined SBML unit for length since SBML Level 2 Version 1.

Definition m

3 Compartment

This model contains one compartment.

Table 2: Properties of all compartments.

Id	Name	SBO	Spatial Dimensions	Size	Unit	Constant	Outside
COMpartment	Wound		3	1	litre	<input checked="" type="checkbox"/>	

3.1 Compartment COMpartment

This is a three dimensional compartment with a constant size of one litre.

Name Wound

4 Species

This model contains four species. The boundary condition of four of these species is set to `true` so that these species' amount cannot be changed by any reaction. Section 7 provides further details and the derived rates of change of each species.

Table 3: Properties of each species.

Id	Name	Compartment	Derived Unit	Constant	Boundary Condi- tion
K_T	K_T	COMpartment	$\text{mol} \cdot \text{l}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
phi_I	phi_I	COMpartment	$\text{mol} \cdot \text{l}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
phi_R	phi_R	COMpartment	$\text{mol} \cdot \text{l}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>
T	T	COMpartment	$\text{mol} \cdot \text{l}^{-1}$	<input type="checkbox"/>	<input checked="" type="checkbox"/>

5 Parameters

This model contains eleven global parameters.

Table 4: Properties of each parameter.

Id	Name	SBO	Value	Unit	Constant
tau1	tau1		-2.470	$(8.64 \cdot 10^{-11} \text{ m})^6 \cdot \text{g}^{-3} \cdot \text{s}^{-1}$	✓
tau2	tau2		21.940	$(0.0864 \text{ mm})^3 \cdot \text{g}^{-2} \cdot \text{s}^{-1}$	✓
tau3	tau3		6.410	$(11.5741 \text{ Mg})^{-1} \cdot \text{s}^{-1}$	✓
tau4	tau4		1.750	$(0.0115741 \text{ m})^{-3} \cdot \text{s}^{-1}$	✓
alpha	alpha		0.800	dimensionless ⁰	✓
k1	k1		0.050	dimensionless ⁰	✓
k2	k2		0.693	$(0.0115741 \text{ ms})^{-1}$	✓
k3	k3		0.002	$(0.0010 \text{ m})^3$	✓
k4	k4		0.070	$(0.0864 \cdot 10^{-6} \text{ dimensionless})^0 \cdot 0.0864 \text{ g} \cdot \text{s}^{-1}$	✓
d1	d1		0.200	$(0.0115741 \text{ ms})^{-1}$	✓
d2	d2		9.100	$(0.0115741 \text{ ms})^{-1}$	✓

6 Rules

This is an overview of four rules.

6.1 Rule K_T

Rule K_T is an assignment rule for species K_T:

$$\text{K_T} = \text{tau1} \cdot [\text{T}]^3 + \text{tau2} \cdot [\text{T}]^2 + \text{tau3} \cdot [\text{T}] + \text{tau4} \quad (1)$$

Derived unit $(8.64 \cdot 10^{-11} \text{ m})^6 \cdot \text{g}^{-3} \cdot \text{s}^{-1} \cdot \text{mol}^3 \cdot \text{l}^{-3}$

6.2 Rule phi_I

Rule phi_I is a rate rule for species phi_I:

$$\frac{d}{dt} \text{phi_I} = \text{alpha} \cdot [\text{K_T}] + \text{k1} \cdot \text{k2} \cdot [\text{phi_I}] \cdot (1 - \text{k3} \cdot ([\text{phi_I}] + [\text{phi_R}]))) - \text{d1} \cdot [\text{phi_I}] \quad (2)$$

6.3 Rule ϕ_R

Rule ϕ_R is a rate rule for species ϕ_R :

$$\frac{d}{dt}\phi_R = (1 - \alpha) \cdot [K_T] + k_1 \cdot k_2 \cdot [\phi_R] \cdot (1 - k_3 \cdot ([\phi_I] + [\phi_R])) - d_1 \cdot [\phi_R] \quad (3)$$

6.4 Rule T

Rule T is a rate rule for species T:

$$\frac{d}{dt}T = k_4 \cdot [\phi_I] - d_2 \cdot [T] \quad (4)$$

Derived unit $0.0864 \text{ g} \cdot \text{s}^{-1} \cdot \text{mol} \cdot \text{l}^{-1}$

7 Derived Rate Equations

When interpreted as an ordinary differential equation framework, this model implies the following set of equations for the rates of change of each species.

7.1 Species K_T

Name K_T

Initial concentration $296.53 \text{ mol} \cdot \text{l}^{-1}$

Involved in rule K_T

One rule determines the species' quantity.

7.2 Species ϕ_I

Name ϕ_I

Initial concentration $200 \text{ mol} \cdot \text{l}^{-1}$

Involved in rule ϕ_I

One rule determines the species' quantity.

7.3 Species ϕ_R

Name ϕ_R

Initial concentration $200 \text{ mol} \cdot \text{l}^{-1}$

Involved in rule ϕ_R

One rule determines the species' quantity.

7.4 Species T

Name T

Initial concentration $6 \text{ mol} \cdot \text{l}^{-1}$

Involved in rule T

One rule determines the species' quantity.

SBML²TeX was developed by Andreas Dräger^a, Hannes Planatscher^a, Dieudonné M Wouamba^a, Adrian Schröder^a, Michael Hucka^b, Lukas Endler^c, Martin Golebiewski^d and Andreas Zell^a. Please see <http://www.ra.cs.uni-tuebingen.de/software/SBML2LaTeX> for more information.

^aCenter for Bioinformatics Tübingen (ZBIT), Germany

^bCalifornia Institute of Technology, Beckman Institute BNMC, Pasadena, United States

^cEuropean Bioinformatics Institute, Wellcome Trust Genome Campus, Hinxton, United Kingdom

^dEML Research gGmbH, Heidelberg, Germany