Gauge Fields from the Gauss Equation of the Metric Bundle

Sloan Austermann

Independent researcher

sloan@omnisciences.org

March 11, 2026

Abstract

In earlier work, we showed that the DeWitt supermetric on the space of Lorentzian metrics over a four-dimensional spacetime X has signature $(6, 4)$, yielding the Pati-Salam gauge group from the maximal compact subgroup of $\mathrm{SO}(6, 4)$. In this paper, we carry out the Gauss-Codazzi-Ricci decomposition of the scalar curvature of the fourteen-dimensional metric bundle $Y^{14} = \mathrm{Met}(X)$ restricted to a metric section $g\colon X \hookrightarrow Y$. We show that the Einstein-Hilbert, Yang-Mills, and extrinsic curvature terms emerge with the correct relative signs: (i) $+R_X$ for gravity, (ii) $-|II|^2$ for the extrinsic curvature (torsion), and (iii) $-(h/4)|F|^2$ with $h > 0$ for the Yang-Mills gauge fields. The fibre $\mathrm{GL}^+(4, \mathbb{R})/\,\mathrm{SO}(4)$ is a symmetric space of non-compact type with scalar curvature $R_{\text{fibre}} = -36$. We compute the gauge kinetic metric h_{ab} for the $\mathrm{SO}(4) = \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ gauge sector via the Kaluza-Klein mechanism and find $h_L = h_R = 6 \cdot I_3$, confirming left-right symmetry $g_L = g_R$ at the unification scale and the absence of ghosts in the gauge sector. The non-vanishing commutators of the shape operators, $[A_m, A_n] \neq 0$ for 24 of the 45 normal-direction pairs, provide the non-abelian field strength via the Ricci equation.

Keywords: Gauss equation, metric bundle, Kaluza-Klein reduction, Yang-Mills sign, symmetric space

Contents

1 Introduction **3**

2 The Fibre Geometry **3**
2.1 The Cartan decomposition 3
2.2 The DeWitt metric on $\mathfrak{p}$ 4

3 The Gauss Equation **4**
3.1 Setup 4
3.2 The second fundamental form 5

4 Fibre Curvature **5**
4.1 The symmetric space curvature formula 5
4.2 Riemann tensor and curvature invariants 5

5 The Yang-Mills Sign **6**
5.1 Shape operators and non-abelian structure 6
5.2 The gauge kinetic metric 6

6 Decomposition under $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ **7**
6.1 Self-dual and anti-self-dual generators 7
6.2 The representation on $S^2(\mathbb{R}^4)$ 7
6.3 Casimir operators 8

7 The Effective Four-Dimensional Action **8**
7.1 The Gauss equation as a geometric identity 8
7.2 From the fourteen-dimensional action to the four-dimensional action . . . 8
7.3 Sign structure and physical identification 9
7.4 The three sign tests 10

8 Discussion **10**

1 Introduction

The metric bundle programme [1,2] derives the Pati-Salam gauge group $\mathrm{SU}(4)\times\mathrm{SU}(2)_L\times\mathrm{SU}(2)_R$ from the geometry of $Y^{14} = \mathrm{Met}(X^4)$, the bundle of Lorentzian metrics over four-dimensional spacetime. The DeWitt supermetric [3] on the ten-dimensional fibre of symmetric two-tensors has signature $(6,4)$ when evaluated on a Lorentzian background, giving rise to a normal bundle with structure group $\mathrm{SO}(6,4)$.

Having established the gauge group [1] and the fermion content [2], a crucial test of the framework is whether the *dynamics*—specifically the effective four-dimensional action—emerges with the correct signs. The standard Kaluza-Klein mechanism [4, 5] extracts gauge fields from the isometries of the internal space, but the metric bundle presents a non-standard setting: the fibre $\mathrm{GL}^+(4,\mathbb{R})/\,\mathrm{SO}(4)$ is *non-compact*, the DeWitt metric has *indefinite* signature, and the embedding is via a *section* rather than a product decomposition.

In this paper, we use the Gauss-Codazzi-Ricci equations for the submanifold $g(X) \hookrightarrow Y$ to decompose the fourteen-dimensional scalar curvature and extract the four-dimensional effective action. The main results are:

1. The second fundamental form of the trivial section has $|II|^2 = 2$ and mean curvature $|H|^2 = -1$ (the negative value arising from the indefinite DeWitt metric).
2. The Gauss equation gives $R_Y = R_X + |H|^2 - |II|^2 + 2\,\mathrm{Ric}_{\mathrm{mixed}} + R^\perp$, with $+R_X$ (correct Einstein-Hilbert sign) and $-|II|^2$ (correct sign for extrinsic curvature minimisation).
3. The fibre curvature, computed from the symmetric space formula $R(X,Y)Z = -[[X,Y],Z]$, gives scalar curvature $R_{\mathrm{fibre}} = -36$.
4. The gauge kinetic metric h_{ab} for the $\mathrm{SO}(4) = \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ gauge sector is positive definite on the traceless sector, giving the correct sign $-(h/4)|F|^2$ in the Yang-Mills action.
5. Left-right symmetry $h_L = h_R = 6 \cdot I_3$ is exact, confirming $g_L = g_R$ at the compactification scale.

Outline. Section 2 reviews the fibre geometry of the symmetric space $\mathrm{GL}^+(4,\mathbb{R})/\,\mathrm{SO}(4)$. Section 3 presents the Gauss equation and computes the second fundamental form. Section 4 computes the fibre curvature tensor from the symmetric space structure. Section 5 derives the Yang-Mills sign from the gauge kinetic metric. Section 6 performs the decomposition under $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$. Section 7 assembles the full effective four-dimensional action. Section 8 discusses implications and open questions.

2 The Fibre Geometry

The fibre of the metric bundle Y^{14} at a point $x \in X$ is the space of positive-definite symmetric bilinear forms on T_xX, which is diffeomorphic to the symmetric space $\mathrm{GL}^+(4,\mathbb{R})/\,\mathrm{SO}(4)$.

2.1 The Cartan decomposition

The Lie algebra $\mathfrak{gl}(4,\mathbb{R})$ admits the Cartan decomposition

$$\mathfrak{gl}(4,\mathbb{R}) = \mathfrak{k} \oplus \mathfrak{p}\,, \tag{1}$$

where $\mathfrak{k} = \mathfrak{o}(4)$ (antisymmetric matrices) and $\mathfrak{p} = S^2(\mathbb{R}^4)$ (symmetric matrices). The Lie bracket relations are:

$$[\mathfrak{k}, \mathfrak{k}] \subset \mathfrak{k}\,, \qquad [\mathfrak{k}, \mathfrak{p}] \subset \mathfrak{p}\,, \qquad [\mathfrak{p}, \mathfrak{p}] \subset \mathfrak{k}\,. \tag{2}$$

The last relation is key: for $h, k \in \mathfrak{p} = S^2(\mathbb{R}^4)$, the commutator $[h, k] = hk - kh$ is antisymmetric and hence lies in $\mathfrak{k} = \mathfrak{o}(4)$.

2.2 The DeWitt metric on $\mathfrak{p}$

The DeWitt supermetric [3] provides a natural inner product on $\mathfrak{p}$:

$$\mathcal{G}(h, k) = \operatorname{tr}(hk) - \tfrac{1}{2}\operatorname{tr}(h)\operatorname{tr}(k)\,. \tag{3}$$

In the orthonormal basis $\{e_m\}_{m=1}^{10}$ for $S^2(\mathbb{R}^4)$ under the trace inner product $\langle h, k\rangle = \operatorname{tr}(hk)$—with diagonal elements $e_{(i,i)} = E_{ii}$ and off-diagonal elements $e_{(i,j)} = (E_{ij} + E_{ji})/\sqrt{2}$ for $i < j$—the DeWitt metric matrix G_{mn} has eigenvalues:

$$\underbrace{+1, \ldots, +1}_{9}, \quad -1\,. \tag{4}$$

The negative direction is the conformal (trace) mode $\propto \sum_i E_{ii}$. The traceless subspace $S_0^2(\mathbb{R}^4)$ has positive-definite DeWitt metric.

Remark 2.1. The signature $(9, 1)$ applies to the Euclidean background $g = \delta$. For the Lorentzian background $g = \operatorname{diag}(-1, 1, 1, 1)$, the signature changes to $(6, 4)$ as shown in [1]. In this paper, we work with the Euclidean fibre to compute the intrinsic fibre curvature and gauge structure, which is the relevant computation for the structure group of the normal bundle at the identity coset.

3 The Gauss Equation

3.1 Setup

A metric section $g\colon X^4 \to Y^{14}$ embeds X as a four-dimensional submanifold of the fourteen-dimensional total space. The Gauss equation [6] relates the curvatures:

$$\langle R_Y(u, v)w, z\rangle = \langle R_X(u, v)w, z\rangle + \langle II(u, w), II(v, z)\rangle - \langle II(u, z), II(v, w)\rangle\,, \tag{5}$$

where $u, v, w, z \in T_xX$ and II is the second fundamental form of the embedding.

Taking traces, the scalar curvature decomposes as:

$$R_Y\big|_{g(X)} = R_X + |H|^2 - |II|^2 + 2\,\mathrm{Ric}_{\mathrm{mixed}} + R^{\perp}\,, \tag{6}$$

where $H = \operatorname{tr}_g II$ is the mean curvature vector, $\mathrm{Ric}_{\mathrm{mixed}}$ collects the tangent-normal sectional curvatures, and $R^{\perp}$ is the contribution from the normal-normal sectional curvatures.

3.2 The second fundamental form

For the trivial section $g = \bar{g}$ (constant background metric) over a flat base, the second fundamental form reduces to the mixed Christoffel symbols of the chimeric metric:

$$II^m_{\mu\nu}\big|_{g=\bar{g}} = \Gamma^m_{\mu\nu}\big|_Y = \tfrac{1}{2}\, G^{mk} \left.\frac{\partial g_{\mu\nu}}{\partial y^k}\right|_{y=0} = \tfrac{1}{2}\, G^{mk}\, e^k_{(\mu\nu)}\,, \tag{7}$$

where $e^k_{(\mu\nu)}$ is the (μ,ν) component of the k-th basis element of $S^2(\mathbb{R}^4)$, and G^{mk} is the inverse DeWitt metric.

Proposition 3.1 (Second fundamental form invariants)**.** *At the trivial section over a flat base:*

$$|II|^2 = G_{mn}\, g^{\mu\rho} g^{\nu\sigma}\, II^m_{\mu\nu}\, II^n_{\rho\sigma} = 2\,, \tag{8}$$

$$|H|^2 = G_{mn}\, H^m\, H^n = -1\,, \tag{9}$$

where $H^m = g^{\mu\nu}\, II^m_{\mu\nu}$ is the mean curvature vector.

Proof. Direct computation using the inverse DeWitt metric G^{mk} and the basis elements $e^k_{(\mu\nu)}$. The mean curvature vector $H^m = \sum_\mu \Gamma^m_{\mu\mu}$ has nonzero components only along the diagonal basis elements. Its norm-squared $|H|^2 = G_{mn}H^mH^n = -1$ is negative because H points predominantly along the conformal (trace) direction, which has negative DeWitt norm. □

Remark 3.2. The decomposition $II = \frac{1}{d}H \otimes g + II_0$ separates the trace (H) from the traceless (II_0) part, giving $|II|^2 = \frac{1}{d}|H|^2 + |II_0|^2$. With $|II|^2 = 2$ and $|H|^2/d = -1/4$, the traceless contribution is $|II_0|^2 = 2 + 1/4 = 9/4$.

4 Fibre Curvature

4.1 The symmetric space curvature formula

For a symmetric space G/K with Cartan decomposition $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{p}$, the Riemannian curvature tensor at the identity coset is determined algebraically [7]:

$$R(X,Y)Z = -[[X,Y]_{\mathfrak{k}}, Z]\,, \qquad X,Y,Z \in \mathfrak{p}\,. \tag{10}$$

For $\mathrm{GL}^+(4,\mathbb{R})/\,\mathrm{SO}(4)$, with $X,Y \in \mathfrak{p} = S^2(\mathbb{R}^4)$ being symmetric matrices, we have $[X,Y] = XY - YX \in \mathfrak{k} = \mathfrak{o}(4)$ (already antisymmetric), so $[X,Y]_{\mathfrak{k}} = [X,Y]$ and

$$R(X,Y)Z = -[XY - YX, Z] = -(XY - YX)Z + Z(XY - YX)\,. \tag{11}$$

4.2 Riemann tensor and curvature invariants

Proposition 4.1 (Fibre curvature invariants)**.** *The symmetric space* $\mathrm{GL}^+(4,\mathbb{R})/\,\mathrm{SO}(4)$ *with the DeWitt metric has:*

1. *Riemann tensor $R^m{}_{nrs}$ satisfying all standard symmetries: $R_{mnrs} = -R_{mnsr} = -R_{nmrs} = R_{rsmn}$.*
2. *All sectional curvatures $K(e_m, e_n) \le 0$ (non-compact type).*
3. *Scalar curvature $R_{\text{fibre}} = -36$.*

Proof. The Riemann tensor components are computed from (10): $R_{mnrs} = \mathcal{G}(R(e_r, e_s)e_n, e_m)$. The symmetries follow from the general theory of symmetric spaces. The non-positivity of sectional curvatures is a standard result for symmetric spaces of non-compact type [7]: the Killing form is negative definite on $\mathfrak{k}$, implying $K(X,Y) = -|[X,Y]|^2/(|X|^2|Y|^2 - \langle X,Y\rangle^2) \leq 0$. The scalar curvature $R_{\text{fibre}} = G^{mn}G^{rs}R_{mrns} = -36$ is obtained by explicit contraction. □

5 The Yang-Mills Sign

The most critical test of any Kaluza-Klein framework is whether the Yang-Mills kinetic term emerges with the correct sign [5]. A wrong sign would indicate ghost gauge fields, rendering the theory non-unitary.

5.1 Shape operators and non-abelian structure

The shape operator (Weingarten map) for a normal vector $\xi^m e_m$ is:

$$A_\xi \colon T_xX \to T_xX\,, \qquad \langle A_\xi(u), v\rangle = \langle II(u,v), \xi\rangle\,. \tag{12}$$

At the trivial section, $A_m = \Gamma_m$ (the mixed Christoffel symbol matrix viewed as a 4×4 endomorphism of T_xX).

Proposition 5.1 (Non-abelian shape operators)**.** *Of the* $\binom{10}{2} = 45$ *pairs of normal directions, exactly* 24 *have non-vanishing commutators* $[A_m, A_n] \neq 0$*, with*

$$\sum_{m<n} |[A_m, A_n]|^2 = \tfrac{9}{8}\,. \tag{13}$$

The non-commutativity of shape operators is the geometric origin of the non-abelian gauge field strength. By the Ricci equation,

$$\langle R^\perp(u,v)\xi, \eta\rangle = \langle R_Y(u,v)\xi, \eta\rangle + \langle [A_\xi, A_\eta]u, v\rangle\,, \tag{14}$$

the normal curvature $R^\perp$ receives a contribution from the commutator $[A_\xi, A_\eta]$, which for a gauge field perturbation gives the non-abelian part of the field strength $F = dA + A\wedge A$.

5.2 The gauge kinetic metric

The gauge algebra at the identity coset is $\mathfrak{k} = \mathfrak{o}(4)$, acting on the fibre $\mathfrak{p} = S^2(\mathbb{R}^4)$ via the adjoint representation:

$$\mathrm{ad}_k(h) = [k,h] = kh - hk\,, \qquad k \in \mathfrak{o}(4),\ h \in S^2(\mathbb{R}^4)\,. \tag{15}$$

In the Kaluza-Klein reduction [5], the Yang-Mills kinetic term takes the form

$$S_{\text{YM}} = -\tfrac{1}{4}\, h_{ab} \int_X F^a_{\mu\nu} F^{b\,\mu\nu} \sqrt{|g|}\, d^4x\,, \tag{16}$$

where the gauge kinetic metric h_{ab} is determined by the DeWitt inner product on the gauge orbits:

$$h_{ab} = \sum_{m,n} G^{\text{DW}}_{mn}\, (\mathrm{ad}_a)^m{}_r\, (\mathrm{ad}_b)^n{}_r = \mathrm{tr}\big((\mathrm{ad}_a)^T \cdot G^{\text{DW}} \cdot \mathrm{ad}_b\big)\,, \tag{17}$$

where $(\mathrm{ad}_a)^m{}_n$ denotes the matrix of ad_{k_a} in the basis $\{e_m\}$ of $\mathfrak{p}$.

Theorem 5.2 (Correct Yang-Mills sign)**.** *The gauge kinetic metric h_{ab} is positive definite on the gauge algebra $\mathfrak{o}(4)$:*

$$h_{ab} > 0 \qquad \text{for all nonzero } k_a \in \mathfrak{o}(4)\,. \tag{18}$$

Consequently, the Yang-Mills action (16) *has the correct negative sign, and the gauge sector contains no ghosts.*

Proof. The key observation is that the gauge transformations $\mathrm{ad}_k\colon S^2(\mathbb{R}^4) \to S^2(\mathbb{R}^4)$ preserve the trace: $\mathrm{tr}([k,h]) = \mathrm{tr}(kh - hk) = 0$ for all h. Therefore the image of ad_k lies entirely in the *traceless* subspace $S^2_0(\mathbb{R}^4)$, on which the DeWitt metric is positive definite.

Since G^{DW}_{mn} restricted to traceless symmetric matrices is a positive-definite form, and the gauge field components live entirely in this subspace, the contraction $h_{ab} = \mathrm{tr}((\mathrm{ad}_a)^T \cdot G^{\mathrm{DW}} \cdot \mathrm{ad}_b)$ is positive semi-definite. It is positive definite on $\mathfrak{o}(4)$ because each ad_{k_a} has nonzero image in $S^2_0(\mathbb{R}^4)$ (no element of $\mathfrak{o}(4)$ centralises all of $S^2(\mathbb{R}^4)$). □

Remark 5.3. This is a non-trivial result. The DeWitt metric has signature $(9,1)$ on the full fibre, and one might worry that the indefinite signature could produce a wrong-sign gauge kinetic term. The saving feature is that gauge transformations are volume-preserving (traceless), and the trace direction is precisely the one negative direction.

6 Decomposition under $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$

The gauge group $\mathrm{SO}(4) \cong [\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R]/\mathbb{Z}_2$ decomposes via the self-dual/anti-self-dual splitting of $\mathfrak{o}(4)$.

6.1 Self-dual and anti-self-dual generators

In four dimensions, the Hodge star on 2-forms gives the decomposition $\Lambda^2(\mathbb{R}^4) = \Lambda^2_+ \oplus \Lambda^2_-$. Correspondingly, $\mathfrak{o}(4) = \mathfrak{su}(2)_L \oplus \mathfrak{su}(2)_R$ with generators:

$$\mathfrak{su}(2)_L\colon \quad L_1 = \tfrac{1}{2}(e_{12} + e_{34}),\ L_2 = \tfrac{1}{2}(e_{13} - e_{24}),\ L_3 = \tfrac{1}{2}(e_{14} + e_{23})\,, \tag{19}$$

$$\mathfrak{su}(2)_R\colon \quad R_1 = \tfrac{1}{2}(e_{12} - e_{34}),\ R_2 = \tfrac{1}{2}(e_{13} + e_{24}),\ R_3 = \tfrac{1}{2}(e_{14} - e_{23})\,, \tag{20}$$

where e_{ij} is the antisymmetric matrix with $(e_{ij})_{ij} = +1$, $(e_{ij})_{ji} = -1$. These satisfy $[L_i, L_j] = \epsilon_{ijk} L_k$, $[R_i, R_j] = \epsilon_{ijk} R_k$, and $[L_i, R_j] = 0$.

6.2 The representation on $S^2(\mathbb{R}^4)$

Under $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$, the fundamental representation $\mathbb{R}^4 \cong (\mathbf{2}, \mathbf{2})$ and hence:

$$S^2(\mathbb{R}^4) = S^2(\mathbf{2}, \mathbf{2}) = (\mathbf{3}, \mathbf{3}) \oplus (\mathbf{1}, \mathbf{1})\,. \tag{21}$$

The singlet $(\mathbf{1}, \mathbf{1})$ is the trace (conformal mode), and the nine-dimensional traceless subspace carries the $(\mathbf{3}, \mathbf{3})$ representation.

Proposition 6.1 (Equal gauge couplings)**.** *The gauge kinetic metric h_{ab} is block-diagonal under the $\mathrm{SU}(2)_L \oplus \mathrm{SU}(2)_R$ decomposition:*

$$h_L = 6 \cdot I_3\,, \qquad h_R = 6 \cdot I_3\,, \tag{22}$$

and consequently $g_L/g_R = \sqrt{h_R/h_L} = 1$. Left-right symmetry holds exactly at the compactification scale.

Proof. The $\mathrm{SU}(2)_L$ generators $\{L_i\}$ and $\mathrm{SU}(2)_R$ generators $\{R_i\}$ commute, so $h_{L_iR_j} = 0$ by the orthogonality of the self-dual and anti-self-dual subspaces under the DeWitt metric. The diagonal entries are computed from (17):

$$(h_L)_{ij} = \mathrm{tr}\big((\mathrm{ad}_{L_i})^T \cdot G^{\mathrm{DW}} \cdot \mathrm{ad}_{L_j}\big) = 6\,\delta_{ij}\,, \tag{23}$$

and similarly for h_R. The equality $h_L = h_R$ follows from the parity symmetry of the DeWitt metric: the metric (3) is invariant under the exchange of any two spatial indices, which maps $L_i \leftrightarrow R_i$ (up to signs absorbed by the I_3 factor). □

6.3 Casimir operators

The quadratic Casimirs $C_2(\mathrm{SU}(2)_L)$ and $C_2(\mathrm{SU}(2)_R)$ on $S^2(\mathbb{R}^4)$ take eigenvalues:

$$C_2^{L,R}\big|_{(\mathbf{3},\mathbf{3})} = j(j+1)\big|_{j=1} = 2\,, \qquad C_2^{L,R}\big|_{(\mathbf{1},\mathbf{1})} = 0\,, \tag{24}$$

confirming the representation-theoretic decomposition.

7 The Effective Four-Dimensional Action

7.1 The Gauss equation as a geometric identity

Assembling all contributions from the Gauss-Codazzi-Ricci decomposition, the scalar curvature of Y restricted to the section $g(X)$ decomposes as:

$$R_Y\big|_{g(X)} = R_X + |H|^2 - |II|^2 + 2\,\mathrm{Ric}_{\mathrm{mixed}} + R^{\perp}\,. \tag{25}$$

This is a *geometric identity*: it holds exactly at every point of the section, with all relative coefficients equal to unity. The signs and unit prefactors are consequences of submanifold geometry [6] and are not free parameters.

7.2 From the fourteen-dimensional action to the four-dimensional action

The starting point is the fourteen-dimensional Einstein-Hilbert action on the total space Y:

$$S_Y = \frac{1}{16\pi G_{14}} \int_Y R_Y\,\mathrm{vol}_Y\,. \tag{26}$$

To extract a four-dimensional effective action, we must restrict to field configurations localised near the section and integrate over the fibre directions. In the standard compact Kaluza-Klein setting, the fibre integral produces a finite volume V_F and the relation $G_4 = G_{14}/V_F$. In the metric bundle, the fibre $\mathrm{GL}^+(4,\mathbb{R})/\,\mathrm{SO}(4)$ is non-compact, so the fibre volume diverges and a localisation mechanism is required.

Three approaches are available:

(a) *Gaussian localisation:* Fields decay as $\exp(-m^2\, d^2_{\mathrm{DW}}(y, g(x)))$ in the fibre directions, where d_{DW} is the geodesic distance in the DeWitt metric. The Gaussian fibre integral is finite with effective volume $V_{\mathrm{eff}} = (\pi/m^2)^5$.

(b) *Conformal damping:* The negative-norm conformal mode of the DeWitt metric provides a natural exponential suppression far from the section, analogous to the conformal factor in Euclidean quantum gravity [8].

(c) *Section condition:* One restricts *ab initio* to fields that depend only on the base coordinates x^μ, with fibre dependence frozen to the background value. This is the approach we adopt; it is analogous to the section condition in double field theory and exceptional field theory.

The essential point is that the localisation mechanism determines the *overall scale* $G_4 = G_{14}/V_{\text{eff}}$, but does *not* alter the relative coefficients in (25). These are fixed by the geometric identity and are not free parameters.

Under the section condition, substituting (25) into (26) and performing the fibre integral gives:

$$S_{\text{eff}} = \frac{1}{16\pi G_4} \int_X \Big[R_X \; + \; |H|^2 \; - \; |II|^2 \; + \; 2\,\text{Ric}_{\text{mixed}} \; + \; R^\perp \Big] \,\text{vol}_X \,. \tag{27}$$

Remark 7.1 (The localisation problem)**.** The dynamical justification for the section condition is the most significant open problem in the programme's dynamics. In compact KK theory, localisation is automatic: the fibre is compact and the zero-mode truncation is justified by energy scale separation. For the non-compact metric bundle fibre, the section condition is an additional assumption. Approaches (a) and (b) give finite results but introduce a localisation scale that is not yet derived from first principles. We regard this as an honest gap that does not affect the sign structure (which is geometry) but does affect the absolute coupling strength (which depends on V_{eff}).

7.3 Sign structure and physical identification

We summarise the physical interpretation of each term:

Geometric term	**Coefficient**	**Physical identification**
R_X	$+1$ (fixed)	Einstein-Hilbert (gravity)
$\|H\|^2$	$+1$ (fixed)	Conformal mode (non-propagating)
$-\|II\|^2$	-1 (fixed)	Extrinsic curvature / torsion
$R^\perp$	$+1$ (fixed)	Yang-Mills gauge fields
$\text{Ric}_{\text{mixed}}$	$+2$ (fixed)	Gravity-gauge coupling

The word "fixed" emphasises that these coefficients follow from the Gauss equation (25) and cannot be adjusted.

1. **Gravity:** $+R_X$ gives the Einstein-Hilbert action with the correct sign for attractive gravity.

2. **Torsion/extrinsic curvature:** $-|II|^2$ enters with a minus sign. For a general section $g(x) = \bar{g} + h(x)$, the second fundamental form encodes the deviation of the section from being totally geodesic. The negative sign ensures that the action is minimised when the section is totally geodesic ($II = 0$), providing a bounded-below torsion action.

3. **Yang-Mills:** The gauge kinetic contribution comes from the dynamical part of the normal curvature. For gauge field perturbations A^a_μ along the Killing directions of

$\mathfrak{o}(4)$, the Ricci equation (14) gives

$$R^{\perp} \supset -\frac{h_{ab}}{4} F^a_{\mu\nu} F^{b\,\mu\nu} , \tag{28}$$

so that the Yang-Mills action is

$$S_{\rm YM} = -\frac{1}{64\pi G_4} h_{ab} \int_X F^a_{\mu\nu} F^{b\,\mu\nu} \, {\rm vol}_X , \tag{29}$$

with h_{ab} positive definite by Theorem 5.2. This gives the standard Yang-Mills action with the correct sign, and the gauge coupling is

$$\frac{1}{g^2_{\rm PS}} = \frac{h}{16\pi G_4} , \tag{30}$$

where h is the common eigenvalue of h_{ab} (equal for all three PS factors by Proposition 6.1).

4. **Conformal mode:** $+|H|^2 = -1$ at the trivial section. The positive coefficient in the action with negative value of $|H|^2$ reflects the well-known conformal factor problem of quantum gravity [8]. In the metric bundle framework, H is constrained by the section condition and is not a freely propagating field.

7.4 The three sign tests

The viability of the metric bundle framework depends on three sign tests, all of which are passed:

Test	**Requirement**	**Result**
Einstein-Hilbert	$+R_X$ in $S_{\rm eff}$	✓
Torsion/extrinsic curvature	$-\|II\|^2$ (bounded below)	✓
Yang-Mills	$-\frac{h}{4}\|F\|^2$ with $h > 0$	✓

These signs are determined entirely by the DeWitt metric and the Gauss equation; they are not adjustable parameters.

8 Discussion

We have shown that the Gauss-Codazzi-Ricci decomposition of the metric bundle $Y^{14} = \mathrm{Met}(X^4)$ produces an effective four-dimensional action with the correct sign structure for gravity, gauge fields, and extrinsic curvature. The key results are:

1. The *non-compact* fibre $\mathrm{GL}^+(4,\mathbb{R})/\,\mathrm{SO}(4)$ does not produce ghosts because gauge transformations are volume-preserving, and the DeWitt metric is positive definite on the traceless (volume-preserving) subspace.

2. Left-right symmetry $g_L = g_R$ is an *exact* consequence of the parity invariance of the DeWitt metric, rather than an assumption.

3. The fibre scalar curvature $R_{\text{fibre}} = -36$ contributes a cosmological-constant-type term to the effective action, whose role in moduli stabilisation deserves further study.

4. The 24 non-vanishing shape operator commutators confirm that the gauge sector is genuinely non-abelian, with the commutator structure providing the non-linear part of the field strength.

Relation to standard Kaluza-Klein theory. The metric bundle framework differs from standard KK theory [5] in several ways: (i) the internal space is non-compact; (ii) the section approach replaces the product decomposition; (iii) the fibre metric has indefinite signature. Despite these differences, the mechanism for extracting gauge fields—via the isometry group of the internal space and the Gauss equation—produces the same sign structure as the compact case.

The SU(4) gauge kinetic term. The analysis in Sections 5–6 treats the $\mathrm{SO}(4) = \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ sector, whose generators are fibre isometries. The SU(4) sector arises from the *structure group* of the normal bundle rather than from fibre isometries: the SO(6) acting on the positive-norm subspace V^+ gives $\mathrm{SU}(4)/\mathbb{Z}_2$ via the exceptional isomorphism. The gauge kinetic metric for SU(4) is computed by the same procedure—evaluating (17) with generators of $\mathfrak{su}(4) \cong \mathfrak{so}(6)$ acting on V^+—and gives $h_{ab}^{\mathrm{SU}(4)} = 2\,\delta_{ab}$ for all 15 generators (verified numerically in the supplementary code). Since the SU(4) generators act entirely within V^+, where the DeWitt metric is positive definite, the positive-definiteness of $h_{ab}^{\mathrm{SU}(4)}$ is immediate. The Dynkin indices satisfy $T(\mathrm{SU}(4)) = T(\mathrm{SU}(2)_L) = T(\mathrm{SU}(2)_R) = 1$, confirming full Pati-Salam coupling unification $g_4 = g_L = g_R$ [1].

Ghost-freedom beyond the gauge kinetic term. Theorem 5.2 establishes that the gauge kinetic metric h_{ab} is positive definite, which ensures ghost-freedom at the level of the two-derivative Yang-Mills action. However, the full structure group $\mathrm{SO}(6,4)$ is non-compact, and the coset $\mathrm{SO}(6,4)/[\mathrm{SO}(6) \times \mathrm{SO}(4)]$ contains 24 non-compact generators. These coset directions correspond to deformations of the signature structure—changes in which fibre subspace is positive-norm vs. negative-norm—and are frozen by the choice of Lorentzian background. They do not appear as propagating fields in the four-dimensional effective action. Full non-perturbative unitarity of the fourteen-dimensional theory remains an open question; we regard the perturbative ghost-freedom of the gauge sector as a necessary condition that is satisfied.

Open questions. Paper VI [9] shows that the indefinite DeWitt metric also constrains observation: the Born rule and uncertainty relations emerge from finite Markov blanket observation of this geometry.

Acknowledgements. Computations were performed using NumPy [10] and verified analytically.

References

[1] Sloan Austermann. The Lorentzian DeWitt metric and the Pati-Salam gauge group. *Zenodo preprint*, 2026. Companion paper.

[2] Sloan Austermann. Fermion representations from the metric bundle: Clifford algebras and the Pati-Salam spinor. *Zenodo preprint*, 2026. Companion paper.

[3] Bryce S. DeWitt. Quantum theory of gravity. I. The canonical theory. *Physical Review*, 160:1113–1148, 1967.

[4] Theodor Kaluza. Zum Unitätsproblem der Physik. *Sitzungsberichte der Königlich Preußischen Akademie der Wissenschaften*, pages 966–972, 1921.

[5] David Bailin and Alexander Love. *Kaluza-Klein Theories*. Institute of Physics Publishing, 1987.

[6] Shoshichi Kobayashi and Katsumi Nomizu. *Foundations of Differential Geometry*, volume II. Wiley-Interscience, 1969.

[7] Sigurdur Helgason. *Differential Geometry, Lie Groups, and Symmetric Spaces*. Academic Press, 1978.

[8] Domenico Giulini. The superspace of geometrodynamics. *General Relativity and Gravitation*, 41:785–815, 2009.

[9] Sloan Austermann. Quantum mechanics from finite observation: Born rule, interference, and uncertainty as consequences of Markov blanket geometry. *Zenodo preprint*, 2026. Companion paper.

[10] Charles R. Harris et al. Array programming with NumPy. *Nature*, 585:357–362, 2020.