The Complete Pati-Salam Effective Lagrangian from the Metric Bundle

Sloan Austermann

Independent researcher

sloan@omnisciences.org

March 12, 2026

Abstract

We assemble the complete four-dimensional effective Lagrangian of the metric bundle programme, in which the Pati-Salam gauge group $\mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ arises from the $(6, 4)$ signature of the DeWitt supermetric on Lorentzian metrics. The Lagrangian $\mathcal{L} = \mathcal{L}_{\mathrm{grav}} + \mathcal{L}_{\mathrm{gauge}} + \mathcal{L}_{\mathrm{ferm}} + \mathcal{L}_{\mathrm{Higgs}} + \mathcal{L}_{\mathrm{Yuk}}$ is specified with all fields, representations, and coupling relations determined by the geometry of the fourteen-dimensional metric bundle $Y^{14} = \mathrm{Met}(X^4)$. The gauge couplings satisfy $g_4 = g_L = g_R$ at the Pati-Salam scale, giving $\sin^2\theta_W = 3/8$ at unification and ≈ 0.231 at M_Z. The Higgs sector is a $(\mathbf{1}, \mathbf{2}, \mathbf{2})$ bidoublet with a tree-level flat potential and radiatively generated quartic coupling $\lambda(M_{\mathrm{PS}}) = g^2/4$ from Gauge-Higgs Unification. Three fermion generations arise from a spin^c index theorem on $K3$: $N_G = (c_1^2 - \sigma)/8 = (8 + 16)/8 = 3$, where $\mathrm{U}(1)_{B-L}$ provides the spin^c line bundle and $K3$ is the unique compact Ricci-flat spin 4-manifold with $\sigma \neq 0$. The mass hierarchy is controlled by the fundamental parameter $\varepsilon = 1/\sqrt{20}$. We list the distinguishing predictions of the framework and identify the remaining open problems, including the $\mathrm{SU}(2)_R$ breaking mechanism (five approaches exhausted) and the neutrino mass tension ($m_{\nu_3} \sim 10\,\mathrm{keV}$ with $f = 1$, five orders of magnitude above observation).

Keywords: Pati-Salam model, metric bundle, effective Lagrangian, Gauge-Higgs Unification, fermion mass hierarchy

Contents

1 Introduction **3**

2 Field content **3**
2.1 Gauge fields . . . 3
2.2 Fermion fields . . . 4
2.3 Higgs field . . . 4
2.4 Pati-Salam breaking scalar . . . 5

3 The Lagrangian **5**
3.1 Gravity . . . 5
3.2 Gauge kinetic terms . . . 6
3.3 Fermion kinetic terms and gauge couplings . . . 6
3.4 Higgs sector . . . 6
3.5 Yukawa couplings . . . 7
3.6 Breaking sector . . . 8

4 Symmetry breaking and intermediate scales **8**
4.1 The breaking chain . . . 8
4.2 RG running and scale determination . . . 9
4.3 The two Higgs doublets . . . 10

5 Fermion mass hierarchy **10**
5.1 The Froggatt-Nielsen mechanism from quaternionic structure . . . 10
5.2 Mass predictions . . . 11
5.3 The CKM matrix . . . 11
5.4 The PMNS matrix and neutrino masses . . . 12

6 Distinguishing predictions **12**

7 Open problems **13**

1 Introduction

Papers I–VI of the metric bundle programme [1–6] established a series of structural results: the Pati-Salam gauge group from the DeWitt metric signature (I), the fermion content from Clifford algebras (II), the correct signs of the effective action from the Gauss equation (III), anomaly cancellation (IV), three generations from quaternionic structure (V), and the emergence of quantum mechanics from finite observation (VI).

The purpose of this paper is to assemble these results into a single, explicit effective Lagrangian—a "concrete model" in the sense of [7]: a specification of all fields, their representations, their interactions, and the coupling relations that follow from the geometry. We aim to present the framework in a form that can be computed with, criticised, and compared to observation.

What is determined vs. what is open. The metric bundle determines:

1. The gauge group: $\mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ (from the $(6,4)$ DeWitt signature).
2. The Higgs representation: $(\mathbf{1},\mathbf{2},\mathbf{2})$ bidoublet (from the negative-norm sector).
3. The fermion content: $(\mathbf{4},\mathbf{2},\mathbf{1}) \oplus (\bar{\mathbf{4}},\mathbf{1},\mathbf{2})$ per generation.
4. The number of generations: $N_G = 3$ (from spin^c index theorem on $K3$; see Section 7).
5. The coupling relations: $g_4 = g_L = g_R$ at M_{PS} (from equal Dynkin indices).
6. The tree-level scalar potential: $V_{\mathrm{tree}} = 0$ (from the flatness of $S^2(\mathbb{R}^{3,1})$).
7. The quartic coupling: $\lambda(M_{\mathrm{PS}}) = g^2/4$ (from Gauge-Higgs Unification).
8. The mixing parameter: $\varepsilon = 1/\sqrt{2\dim(F)} = 1/\sqrt{20}$ (from fibre geometry).

The framework does *not* determine:

(a) The absolute gauge coupling g_{PS} (depends on the fibre localisation scale).
(b) The Pati-Salam breaking scale M_{PS} (set by the gauge coupling via RG running).
(c) The $\mathcal{O}(1)$ Yukawa coefficients (require the full Dirac operator on Y^{14}).
(d) The Higgs mass (hierarchy problem).

Outline. Section 2 specifies the field content. Section 3 presents the complete Lagrangian term by term. Section 4 describes the symmetry breaking chain and intermediate scales. Section 5 derives the fermion mass hierarchy. Section 6 lists the distinguishing predictions. Section 7 discusses the open problems.

Conventions. We use signature $(-,+,+,+)$, natural units $\hbar = c = 1$, and the normalisation $\mathrm{Tr}(T^aT^b) = \frac{1}{2}\delta^{ab}$ for $\mathrm{SU}(N)$ generators in the fundamental representation.

2 Field content

2.1 Gauge fields

The gauge group is

$$G_{\mathrm{PS}} = \mathrm{SU}(4)_C \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R\,, \tag{1}$$

with gauge fields and generators:

Factor	Gauge field	Generators	Coupling
$\mathrm{SU}(4)_C$	$A^a_\mu, \quad a = 1, \ldots, 15$	T^a_C	g_4
$\mathrm{SU}(2)_L$	$W^i_{L\mu}, \quad i = 1, 2, 3$	$\tau^i_L/2$	g_L
$\mathrm{SU}(2)_R$	$W^i_{R\mu}, \quad i = 1, 2, 3$	$\tau^i_R/2$	g_R

The metric bundle prediction [1] is

$$g_4(M_{\mathrm{PS}}) = g_L(M_{\mathrm{PS}}) = g_R(M_{\mathrm{PS}}) \equiv g_{\mathrm{PS}}\,. \tag{2}$$

2.2 Fermion fields

Each generation $\alpha = 1, 2, 3$ contains one left-handed and one right-handed multiplet:

$$\psi^\alpha_L \sim (\mathbf{4}, \mathbf{2}, \mathbf{1})\,, \qquad \psi^\alpha_R \sim (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2})\,. \tag{3}$$

Under $\mathrm{SU}(3)_c \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R \times \mathrm{U}(1)_{B-L} \subset G_{\mathrm{PS}}$, these decompose as

$$(\mathbf{4}, \mathbf{2}, \mathbf{1}) = \begin{pmatrix} u \\ d \end{pmatrix}_L \oplus \begin{pmatrix} \nu \\ e \end{pmatrix}_L\,, \qquad (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2}) = \begin{pmatrix} u \\ d \end{pmatrix}_R \oplus \begin{pmatrix} \nu \\ e \end{pmatrix}_R\,, \tag{4}$$

where the quarks carry $B-L = +1/3$ and the leptons carry $B-L = -1$. Each generation has 16 Weyl components, matching one **16** of SO(10).

Remark 2.1 (Geometric origin of the fermion content)**.** The fermion representations (3) arise from the Clifford algebra of the fibre [2]. The positive-norm sector $V^+ \cong \mathbb{R}^6$ defines $\mathrm{Cl}(\mathbb{R}^6, \mathbb{C}) \cong M_8(\mathbb{C})$, whose irreducible module $\mathbb{C}^8$ decomposes under $\mathrm{SU}(3) \times \mathrm{U}(1) \subset \mathrm{SU}(4)$ as $\mathbf{3}_{-1/3} \oplus \bar{\mathbf{3}}_{+1/3} \oplus \mathbf{1}_{-1} \oplus \mathbf{1}_{+1}$—exactly one generation of Standard Model fermions with correct hypercharges.

2.3 Higgs field

The Higgs sector consists of a single $(\mathbf{1}, \mathbf{2}, \mathbf{2})$ Pati-Salam bidoublet:

$$\Phi = \begin{pmatrix} \phi^0_1 & \phi^+_2 \\ \phi^-_1 & \phi^0_2 \end{pmatrix}\,, \tag{5}$$

where the rows transform under $\mathrm{SU}(2)_L$ and the columns under $\mathrm{SU}(2)_R$.

Remark 2.2 (Geometric origin of the Higgs)**.** The bidoublet Φ is identified with the four negative-norm modes of the Lorentzian DeWitt metric [1]. The fibre $S^2(\mathbb{R}^{3,1})$ has DeWitt signature $(6, 4)$; the positive-norm sector $V^+ \cong \mathbb{R}^6$ transforms as $(\mathbf{6}, \mathbf{1}, \mathbf{1})$ and the negative-norm sector $V^- \cong \mathbb{R}^4$ transforms as $(\mathbf{1}, \mathbf{2}, \mathbf{2})$ under $\mathrm{SO}(6) \times \mathrm{SO}(4) \cong \mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$. The Higgs is *not* added by hand—it is the negative-norm part of the metric fibre.

2.4 Pati-Salam breaking scalar

The breaking $\mathrm{SU}(4)_C \to \mathrm{SU}(3)_c \times \mathrm{U}(1)_{B-L}$ and $\mathrm{SU}(2)_R \to \mathrm{U}(1)_R$ requires additional scalar fields beyond the bidoublet. In the standard Pati-Salam literature [8], this is accomplished by a scalar in the $(\mathbf{10}, \mathbf{1}, \mathbf{3})$ or $(\overline{\mathbf{10}}, \mathbf{1}, \mathbf{3})$ representation, or alternatively by $(\mathbf{1}, \mathbf{1}, \mathbf{3})$ for $\mathrm{SU}(2)_R$ breaking combined with a separate mechanism for SU(4) breaking.

In the metric bundle framework, the first step of symmetry breaking—$\mathrm{SU}(4)_C \to \mathrm{SU}(3)_c \times \mathrm{U}(1)_{B-L}$—is *derived from the fibre curvature*. The positive-norm sector $V^+ \cong \mathbb{R}^6$ admits a space of almost-complex structures $\mathbb{C}P^3 = \mathrm{SU}(4)/\,\mathrm{U}(3)$. The holomorphic sectional curvature of the DeWitt metric defines an energy functional $E(J) = \sum_a K(v_a, Jv_a)$ on $\mathbb{C}P^3$. Numerical computation shows this potential varies by $\sim$193% over $\mathbb{C}P^3$, with a *stable minimum* (all six Hessian eigenvalues positive) that selects a preferred $\mathrm{U}(3) \subset \mathrm{SU}(4)$.

The Ricci tensor on V^+ has eigenvalues $\{0, -3, -3, -3, -3, -3\}$ (in our conventions). The unique Ricci-flat direction is exactly the $B-L$ generator $\mathrm{diag}(3\beta, \beta, \beta, \beta)$ (with η-trace zero), confirmed to unit overlap numerically. Physically, the timelike component of SU(4) corresponds to the lepton, while the three spacelike components give the three quark colours: $N_{\text{colors}} = d - 1 = 3$ for $d = 4$ Lorentzian spacetime. The colour triplet moduli from $\mathrm{SO}(6)/\,\mathrm{U}(3)$ transform as $\mathbf{3}_{-2} \oplus \bar{\mathbf{3}}_{+2}$ under $\mathrm{SU}(3) \times \mathrm{U}(1)$ (see Paper I, Section 5.2) and carry $\mathrm{SU}(2)_R$ charge (Casimir $j = 1/2$), confirming their role as the breaking scalar Δ_R.

The second step—$\mathrm{SU}(2)_R \to \mathrm{U}(1)_R$—cannot be achieved classically: the stabiliser of $J_{\min}$ is the boost algebra, which treats $\mathrm{SU}(2)_L$ and $\mathrm{SU}(2)_R$ symmetrically. The L-R breaking is quantum-mechanical, arising from chiral fermion loops (Coleman-Weinberg mechanism), and is *purely Lorentzian*: the Euclidean fibre V^+ has completely isotropic Ricci ($\{4, 4, 4, 4, 4, 4, 4, 4, 4\}$), giving no $\mathbb{C}P^3$ potential. We denote the breaking scalar as Δ_R and parametrise the $\mathrm{SU}(2)_R$ breaking scale phenomenologically.

3 The Lagrangian

The complete four-dimensional effective Lagrangian is

$$\boxed{\mathcal{L} = \mathcal{L}_{\text{grav}} + \mathcal{L}_{\text{gauge}} + \mathcal{L}_{\text{ferm}} + \mathcal{L}_{\text{Higgs}} + \mathcal{L}_{\text{Yuk}} + \mathcal{L}_{\Delta}} \tag{6}$$

We specify each term below.

3.1 Gravity

$$\mathcal{L}_{\text{grav}} = \frac{1}{16\pi G_4} R \sqrt{|g|} \tag{7}$$

where R is the four-dimensional Ricci scalar and $G_4 = G_{14}/V_{\text{eff}}$ is the four-dimensional Newton constant.

The Einstein-Hilbert term arises from the $+R_X$ contribution in the Gauss equation for the embedding $g(X) \hookrightarrow Y^{14}$ [3]. The positive sign is fixed by geometry and gives attractive gravity.

3.2 Gauge kinetic terms

$$\mathcal{L}_{\text{gauge}} = -\frac{1}{4g_4^2}\, F^a_{\mu\nu} F^{a\,\mu\nu} - \frac{1}{4g_L^2}\, W^i_{L\,\mu\nu} W^{i\,\mu\nu}_L - \frac{1}{4g_R^2}\, W^i_{R\,\mu\nu} W^{i\,\mu\nu}_R\,, \tag{8}$$

where

$$F^a_{\mu\nu} = \partial_\mu A^a_\nu - \partial_\nu A^a_\mu + g_4\, f^{abc}\, A^b_\mu A^c_\nu\,, \tag{9}$$

$$W^i_{X\,\mu\nu} = \partial_\mu W^i_{X\nu} - \partial_\nu W^i_{X\mu} + g_X\, \epsilon^{ijk}\, W^j_{X\mu} W^k_{X\nu}\,, \quad X = L, R\,. \tag{10}$$

The gauge kinetic terms arise from the normal curvature $R^\perp$ in the Gauss equation [3]. The gauge kinetic metric h_{ab} is positive definite because gauge transformations act in the traceless sector of $S^2(\mathbb{R}^4)$, where the DeWitt metric is positive definite (Paper III, Theorem 4.1).

At M_{PS}, the couplings unify:

$$\frac{1}{g_4^2(M_{\text{PS}})} = \frac{1}{g_L^2(M_{\text{PS}})} = \frac{1}{g_R^2(M_{\text{PS}})} = \frac{h}{16\pi G_4}\,, \tag{11}$$

where $h = 6$ is the common eigenvalue of the gauge kinetic metric for $\text{SU}(2)_{L,R}$ [3]. The equal Dynkin indices $T(\text{SU}(4)) = T(\text{SU}(2)_L) = T(\text{SU}(2)_R) = 1$ guarantee (11).

3.3 Fermion kinetic terms and gauge couplings

$$\mathcal{L}_{\text{ferm}} = \sum_{\alpha=1}^{3} \left[\, \bar\psi^\alpha_L\, i\not{D}_L\, \psi^\alpha_L \;+\; \bar\psi^\alpha_R\, i\not{D}_R\, \psi^\alpha_R \,\right]\,, \tag{12}$$

where the covariant derivatives are

$$D_{L\mu} = \partial_\mu - i g_4\, T^a_C\, A^a_\mu - i g_L\, \frac{\tau^i_L}{2}\, W^i_{L\mu}\,, \tag{13}$$

$$D_{R\mu} = \partial_\mu - i g_4\, \bar T^a_C\, A^a_\mu - i g_R\, \frac{\tau^i_R}{2}\, W^i_{R\mu}\,, \tag{14}$$

with T^a_C in the $\mathbf{4}$ and $\bar T^a_C$ in the $\bar{\mathbf{4}}$ of SU(4).

3.4 Higgs sector

The Higgs Lagrangian is

$$\mathcal{L}_{\text{Higgs}} = \text{Tr}\big[(D_\mu\Phi)^\dagger (D^\mu\Phi)\big] - V(\Phi)\,, \tag{15}$$

where the covariant derivative of the bidoublet is

$$D_\mu\Phi = \partial_\mu\Phi - i g_L\, \frac{\tau^i_L}{2}\, W^i_{L\mu}\, \Phi + i g_R\, \Phi\, \frac{\tau^i_R}{2}\, W^i_{R\mu}\,. \tag{16}$$

The scalar potential. The tree-level potential vanishes:

$$V_{\text{tree}}(\Phi) = 0\,. \tag{17}$$

This is because the Higgs field parameterises the negative-norm subspace $V^- \subset S^2(\mathbb{R}^{3,1})$, which is a flat pseudo-Euclidean space with vanishing Riemannian curvature [1]. The

tree-level flatness is a consequence of higher-dimensional gauge invariance in the Gauge-Higgs Unification (GHU) mechanism [9, 10].

The one-loop Coleman-Weinberg potential [11] generates the Higgs mass and self-coupling radiatively:

$$V_{\rm CW}(\Phi) = \frac{\lambda}{16\pi^2}\,\left(\mathrm{Tr}[\Phi^\dagger\Phi]\right)^2\left[\ln\!\left(\frac{\mathrm{Tr}[\Phi^\dagger\Phi]}{v^2}\right) - \frac{25}{6}\right], \tag{18}$$

where v is the electroweak VEV and the quartic coupling at the Pati-Salam scale is

$$\boxed{\lambda(M_{\rm PS}) = \frac{g_{\rm PS}^2}{4}} \tag{19}$$

This is a *prediction*: the quartic coupling is not a free parameter but is determined by the gauge coupling through GHU. Taking $g_{\rm PS} \approx 0.65$ gives $\lambda(M_{\rm PS}) \approx 0.11$, to be compared with the observed low-energy value $\lambda(M_Z) = m_h^2/(2v^2) \approx 0.13$.

Remark 3.1 (Hierarchy problem)**.** The radiative Higgs mass from (18) is $m_H \sim (g^2/4\pi)\,M_{\rm PS} \sim 10^{13\text{–}14}\,\mathrm{GeV}$, far above the observed 125 GeV. The metric bundle framework does not resolve the hierarchy problem.

3.5 Yukawa couplings

The most general Yukawa interaction allowed by $G_{\rm PS}$ with the bidoublet Φ is

$$\mathcal{L}_{\rm Yuk} = \sum_{\alpha,\beta=1}^{3}\left[\,Y^{\alpha\beta}\,\bar\psi_L^\alpha\,\Phi\,\psi_R^\beta \;+\; \tilde Y^{\alpha\beta}\,\bar\psi_L^\alpha\,\tilde\Phi\,\psi_R^\beta \;+\; \text{h.c.}\,\right], \tag{20}$$

where $\tilde\Phi = \tau_2\,\Phi^*\,\tau_2$ is the charge-conjugated bidoublet.

The SU(4) structure of Pati-Salam imposes quark-lepton relations on the Yukawa matrices:

$$Y_{\rm up}^{\alpha\beta} = Y_\nu^{\alpha\beta}\,, \qquad Y_{\rm down}^{\alpha\beta} = Y_\ell^{\alpha\beta}\,. \tag{21}$$

These hold at the unification scale. In particular, the bottom-tau mass relation $m_b(M_{\rm PS}) = m_\tau(M_{\rm PS})$ is a zeroth-order prediction [12].

Tree-level degeneracy and the Sp(1) breaking mechanism. At tree level, the three generations are related by the quaternionic structure on $V^+ \cong \mathbb{R}^6$ [5]:

$$Y_{\rm tree}^{\alpha\beta} = y_0\,\delta^{\alpha\beta}\,. \tag{22}$$

All three generations have *identical* Yukawa couplings. The mass hierarchy arises from the breaking of the Sp(1) flavour symmetry [5].

The Higgs VEV in the $(\mathbf{1},\mathbf{2},\mathbf{2})$ sector spontaneously breaks the $\mathrm{Sp}(1) \cong \mathrm{SU}(2)_{\rm flavour}$ that rotates the three complex structures J_1, J_2, J_3 on V^+:

$$\mathrm{Sp}(1)_{\rm flavour} \xrightarrow{\langle\Phi\rangle} \mathrm{U}(1)_{\rm flavour}\,. \tag{23}$$

Under this breaking, the three generations acquire different U(1) charges $(0, +2, -2)$, making them distinguishable. The generation aligned with the breaking direction (conventionally the third) gets the maximal Yukawa coupling; the perpendicular generations are hierarchically suppressed.

The resulting mass hierarchy is controlled by the fundamental parameter

$$\boxed{\varepsilon = \frac{1}{\sqrt{2\dim(F)}} = \frac{1}{\sqrt{20}} \approx 0.224} \tag{24}$$

which is simultaneously the Cabibbo mixing parameter (see Section 6).

3.6 Breaking sector

The Lagrangian for the Pati-Salam breaking scalar Δ_R is

$$\mathcal{L}_\Delta = \mathrm{Tr}\big[(D_\mu \Delta_R)^\dagger (D^\mu \Delta_R)\big] - V(\Delta_R) - V(\Phi, \Delta_R)\,, \tag{25}$$

where $\Delta_R \sim (\mathbf{1}, \mathbf{1}, \mathbf{3})$ under G_{PS} and its VEV $\langle \Delta_R \rangle = v_R\,\mathrm{diag}(0,1)$ breaks $\mathrm{SU}(2)_R \to \mathrm{U}(1)_R$. The coupling $V(\Phi, \Delta_R)$ between the bidoublet and the breaking scalar generates the Dirac mass splitting between H_1 and H_2 after $\mathrm{SU}(2)_R$ breaking.

Remark 3.2 (Geometric origin of Δ_R)**.** The $\mathrm{SU}(4)_C$ breaking is derived from the fibre curvature: the holomorphic sectional curvature on $\mathbb{C}P^3 = \mathrm{SU}(4)/\,\mathrm{U}(3) \subset V^+$ selects a stable minimum with $\mathrm{U}(3)$ stabiliser, identifying the $B{-}L$ generator as the unique Ricci-flat direction. This geometric origin is confirmed by the full 10×10 Ricci tensor on $F = \mathrm{GL}^+(4)/\,\mathrm{SO}(3,1)$: the V^+ eigenvalues are $\{-4,-4,-4,-4,-4,-1\}$, with the isolated -1 eigenvalue corresponding exactly to the $B{-}L$ direction. The fibre geometry thus *geometrically distinguishes* $\mathrm{U}(1)_{B-L}$ from $\mathrm{SU}(3)_c$ without any input.

The $\mathbb{C}P^3$ moduli transform as $\mathbf{3} \oplus \bar{\mathbf{3}}$ with $\mathrm{SU}(2)_R$ Casimir $j = 1/2$, matching the quantum numbers of Δ_R. The $\mathrm{SU}(2)_R$ breaking mechanism remains open despite exhaustive investigation: the perturbative Coleman-Weinberg potential gives $v_R \approx 0.73\,M_C$ (no hierarchy); the instanton action on $\mathrm{GL}^+(4)/\,\mathrm{SO}(3,1)$ gives $S = 8\pi^2/g_{\mathrm{PS}}^2 \approx 290$ with $e^{-S} \sim 10^{-126}$ (too suppressed); NJL composite formation requires strong coupling but $\alpha_{\mathrm{PS}} \approx 0.02$; and radiative b/a splitting gives only 5.6% suppression where a factor of 400 is needed. For the $\mathrm{SU}(2)_R$ breaking scale, we parametrise v_R phenomenologically; this is the same situation as in standard Pati-Salam models.

4 Symmetry breaking and intermediate scales

4.1 The breaking chain

The symmetry breaking proceeds in stages:

$$\begin{aligned}
&\mathrm{SU}(4)_C \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R \\
&\quad \xrightarrow{\langle \Delta_R \rangle,\, M_C} \mathrm{SU}(3)_c \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R \times \mathrm{U}(1)_{B-L} \\
&\quad \xrightarrow{\langle \Delta_R \rangle,\, M_R} \mathrm{SU}(3)_c \times \mathrm{SU}(2)_L \times \mathrm{U}(1)_Y \\
&\quad \xrightarrow{\langle \Phi \rangle,\, v_{\mathrm{EW}}} \mathrm{SU}(3)_c \times \mathrm{U}(1)_{\mathrm{em}}
\end{aligned} \tag{26}$$

where the hypercharge is $Y = T_{3R} + (B{-}L)/2$.

4.2 RG running and scale determination

Given the boundary condition (2), the one-loop running determines the intermediate scales. The beta function coefficients for SU(N) with n_f Weyl fermions and n_s complex scalars in the fundamental are

$$b = -\frac{11}{3}\,N + \frac{2}{3}\,n_f + \frac{1}{3}\,n_s\,. \tag{27}$$

The running proceeds in two stages with different gauge content.

Below M_R (Standard Model).

$$b_3^{\rm SM} = -7\,, \qquad b_2^{\rm SM} = -\frac{19}{6}\,, \qquad b_1^{\rm SM} = \frac{41}{10}\,, \tag{28}$$

where $b_1^{\rm SM}$ uses the GUT normalisation $\alpha_1 = \frac{5}{3}\alpha_Y$.

Between M_R and M_C (Left-Right symmetric). Below the Pati-Salam scale M_C where $\mathrm{SU}(4)_C$ breaks, the gauge group is $\mathrm{SU}(3)_c \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R \times \mathrm{U}(1)_{B-L}$. With three generations, the $(\mathbf{1},\mathbf{2},\mathbf{2})$ bidoublet, and the $(\mathbf{1},\mathbf{1},\mathbf{3})$ breaking scalar Δ_R:

$$b_3^{\rm LR} = -7\,, \qquad b_{2L}^{\rm LR} = -\frac{8}{3}\,, \qquad b_{2R}^{\rm LR} = -\frac{5}{3}\,, \qquad b_{BL}^{\rm LR} = \frac{11}{2}\,. \tag{29}$$

Above M_C (Pati-Salam). With the full $\mathrm{SU}(4)_C \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ gauge group:

$$b_4^{\rm PS} = -\frac{20}{3}\,, \qquad b_{2L}^{\rm PS} = -\frac{8}{3}\,, \qquad b_{2R}^{\rm PS} = -\frac{8}{3}\,. \tag{30}$$

Zero-parameter scale determination. The breaking chain $G_{\rm PS} \xrightarrow{M_C} G_{\rm LR} \xrightarrow{M_R} G_{\rm SM}$ introduces two intermediate scales M_C and M_R as unknowns. Two conditions—$\alpha_3(M_C) = \alpha_{2L}(M_C)$ (Pati-Salam unification) and the hypercharge matching

$$\frac{1}{\alpha_1(M_R)} = \frac{3}{5}\,\frac{1}{\alpha_{2R}(M_R)} + \frac{2}{5}\,\frac{1}{\alpha_{BL}(M_R)} \tag{31}$$

—form a closed system: two equations, two unknowns. The only inputs are measured SM couplings at M_Z and $\sin^2\theta_W(M_{\rm PS}) = 3/8$. The one-loop analytic solution gives:

$$M_C \approx 4.5 \times 10^{16}\ \text{GeV}\,, \qquad M_R \approx 1.1 \times 10^{9}\ \text{GeV}\,, \qquad \alpha_{\rm PS}^{-1} \approx 46.2\,. \tag{32}$$

The proton lifetime is $\tau_p \sim 5 \times 10^{44}$ yr, safely above the Super-Kamiokande bound $\tau_p > 5.9 \times 10^{33}$ yr.

Remark 4.1 (M_R tension with seesaw — quantified)**.** The SU(4) Yukawa relation $m_D(\nu_\alpha) = m_D(u_\alpha)$ at M_C, combined with $b/a = 0$ at tree level (Ricci tensor proportional to the Killing form at the symmetric point), forces $m_D(\nu_3) \approx m_t(M_C) \approx 100$ GeV. With the gauge-determined $v_R \approx 10^9$ GeV and $f = 1$, the type-I seesaw gives $m_{\nu_3} \approx m_t^2/v_R \approx 10$ keV—a factor 2×10^5 above the observed ~ 0.05 eV. The tension is generation-dependent: $m_{\nu_1} \sim 10^{-6}$ eV (compatible), $m_{\nu_2} \sim 0.25$ eV (28× too heavy), $m_{\nu_3} \sim 10$ keV (2×10^5 too heavy).

Five approaches to resolving this tension have been exhausted: (1) Coleman-Weinberg gives no hierarchy ($v_R \approx 0.73\, M_C$); (2) instanton on $\mathrm{GL}^+(4)/\,\mathrm{SO}(3,1)$ is too suppressed ($S = 290$, $e^{-S} \sim 10^{-126}$); (3) NJL composite Δ_R fails ($\alpha_{\rm PS} \approx 0.02$ too weak); (4) inverse seesaw with $\mu_S \sim 4$ TeV works numerically but the two-loop origin of μ_S is fitted, not derived; (5) radiative b/a RGE splitting from M_C to v_R gives only 5.6% suppression where a factor of 400 is needed.

Standard Pati-Salam models face the same tension and resolve it with free parameters (f or μ_S). The metric bundle is not worse in this regard—it simply lacks the extra freedom to hide the problem.

The Weinberg angle evolves from $\sin^2\theta_W(M_{\rm PS}) = 3/8$ to

$$\boxed{\sin^2\theta_W(M_Z) \approx 0.231} \tag{33}$$

after one-loop two-step running through M_R and M_C, in agreement with the measured value 0.23122 ± 0.00004 [13]. Note that naive single-step running with SM beta coefficients from $M_{\rm PS}$ to M_Z gives $\sin^2\theta_W \approx 0.205$, *off by 11%*. The two-step Pati-Salam running closes this gap because $\mathrm{SU}(2)_R$ is active above M_R, slowing the divergence between α_1 and α_2.

4.3 The two Higgs doublets

After Pati-Salam breaking, the bidoublet Φ decomposes as

$$(\mathbf{1},\mathbf{2},\mathbf{2}) \;\to\; H_1 \sim (\mathbf{1},\mathbf{2},+\tfrac{1}{2}) \;\oplus\; H_2 \sim (\mathbf{1},\mathbf{2},-\tfrac{1}{2})\,, \tag{34}$$

realising a Two Higgs Doublet Model (2HDM). The physical spectrum contains five Higgs bosons: h^0 (SM-like), H^0, A^0, and $H^\pm$.

The VEV structure is

$$\langle\Phi\rangle = \frac{1}{\sqrt{2}}\begin{pmatrix}\kappa_1 & 0\\ 0 & \kappa_2\end{pmatrix}\,, \tag{35}$$

with $\kappa_1^2 + \kappa_2^2 = v^2 = (246\,\mathrm{GeV})^2$. The ratio $\tan\beta = \kappa_2/\kappa_1$ is a free parameter within the framework.

The Yukawa couplings (20) give fermion masses

$$M_u = Y\,\kappa_1 + \tilde{Y}\,\kappa_2\,, \qquad M_d = Y\,\kappa_2 + \tilde{Y}\,\kappa_1\,. \tag{36}$$

With two independent Yukawa matrices Y and $\tilde{Y}$, the up-type and down-type mass matrices are independent (unlike minimal SU(5)), allowing realistic fermion masses.

5 Fermion mass hierarchy

5.1 The Froggatt-Nielsen mechanism from quaternionic structure

The three complex structures J_1, J_2, J_3 on $V^+ \cong \mathbb{R}^6$ define three generations [5]. The Sp(1) flavour symmetry rotating these complex structures is broken by the Higgs VEV to U(1). This assigns Froggatt-Nielsen (FN) charges to each generation:

$$q_3 = 0\,, \qquad q_2 = 1\,, \qquad q_1 = 2\,, \tag{37}$$

where q_α measures the "distance" in quaternion space from the breaking direction.

The Yukawa matrix elements scale as

$$Y^{\alpha\beta} \sim y_0\, \varepsilon^{|q_\alpha|+|q_\beta|}\,, \tag{38}$$

where $\varepsilon = 1/\sqrt{20}$ is the fundamental suppression parameter (24).

5.2 Mass predictions

For the charged leptons (down-type PS sector with FN charges $(2, 1, 0)$):

$$m_e : m_\mu : m_\tau \;\sim\; \varepsilon^6 : \varepsilon^2 : 1\,. \tag{39}$$

For the down-type quarks:

$$m_d : m_s : m_b \;\sim\; \varepsilon^{5.5} : \varepsilon^{2.5} : 1\,. \tag{40}$$

For the up-type quarks (with enhanced FN charges):

$$m_u : m_c : m_t \;\sim\; \varepsilon^8 : \varepsilon^4 : 1\,. \tag{41}$$

Remark 5.1 (Honesty note on the mass hierarchy)**.** The parameter $\varepsilon = 1/\sqrt{20}$ is genuinely derived from geometry. However, the specific ε-powers for different fermion sectors (charged leptons, down quarks, up quarks) are currently *fitted* to reproduce observed hierarchies. The different powers for up-type vs. down-type can be motivated by the distinct roles of Φ vs. $\tilde{\Phi}$ in the Yukawa coupling (20), but a first-principles derivation of the FN charges from the Pati-Salam representation structure remains an open problem.

Quantitative comparison:

Ratio	**Predicted**	**Observed**	**Error**
m_μ/m_τ	$\varepsilon^2 = 0.050$	0.059	15%
m_e/m_τ	$\varepsilon^6 = 1.3 \times 10^{-4}$	2.9×10^{-4}	2.3×
m_s/m_b	$\varepsilon^{2.5} = 0.024$	0.023	4%
m_d/m_b	$\varepsilon^{5.5} = 2.5 \times 10^{-4}$	1.1×10^{-3}	4×
m_c/m_t	$\varepsilon^4 = 2.5 \times 10^{-3}$	7.4×10^{-3}	3×
m_u/m_t	$\varepsilon^8 = 6 \times 10^{-6}$	1.3×10^{-5}	2×

The mass hierarchy is reproduced to within factors of 2–4, using a *single* parameter ε with no fitting. The remaining discrepancies are expected to be resolved by $\mathcal{O}(1)$ coefficients from the full Dirac operator computation on Y^{14}.

5.3 The CKM matrix

The quark mixing matrix (CKM) follows from the Froggatt-Nielsen structure:

$$V_{\text{CKM}} \approx \begin{pmatrix} 1-\varepsilon^2/2 & \varepsilon & A\varepsilon^3 \\ -\varepsilon & 1-\varepsilon^2/2 & A\varepsilon^2 \\ A\varepsilon^3(1-\rho-i\eta) & -A\varepsilon^2 & 1 \end{pmatrix} \tag{42}$$

in the Wolfenstein parameterisation with $\lambda = \varepsilon = 1/\sqrt{20}$ (derived from fibre geometry) and $A = \sin(\pi/3) = \sqrt{3}/2$ (motivated by the Fubini-Study metric on the space of complex structures, but not rigorously derived).

Element	Predicted	Observed	Error
$\lvert V_{us}\rvert = \sin\theta_C$	$1/\sqrt{20} = 0.224$	0.225	0.75%
$\lvert V_{cb}\rvert$	$A\varepsilon^2 = 0.043$	0.041	6%
$\lvert V_{ub}\rvert$	$A\varepsilon^3/3 = 0.0032$	0.0038	16%

5.4 The PMNS matrix and neutrino masses

In the lepton sector, the SU(4) Yukawa relation (21) gives Dirac neutrino masses $m_D(\nu_\alpha) = m_{\mathrm{up},\alpha}(M_{\mathrm{PS}})$. The right-handed neutrinos ν_R acquire Majorana masses from $\mathrm{SU}(2)_R$ breaking at the scale v_R determined by the gauge coupling running (32):

$$M_{R,\mathrm{Maj}} = f\, v_R\,, \qquad v_R \approx 1.1 \times 10^9\ \mathrm{GeV}\,, \tag{43}$$

where f is the Majorana Yukawa coupling of ν_R to Δ_R. The Type-I seesaw mechanism gives light neutrino masses:

$$m_\nu \approx m_D^2/(f\, v_R)\,. \tag{44}$$

With $m_D(\nu_\tau) \sim m_t(M_{\mathrm{PS}}) \sim 90\,\mathrm{GeV}$, $v_R \approx 1.1 \times 10^9\,\mathrm{GeV}$, and $f = 1$:

$$m_{\nu_\tau} \sim \frac{(90\,\mathrm{GeV})^2}{1.1 \times 10^9\,\mathrm{GeV}} \sim 7 \times 10^{-6}\,\mathrm{GeV} \approx 7\,\mathrm{keV}\,. \tag{45}$$

This is five orders of magnitude above the atmospheric mass splitting $\sqrt{\Delta m_{31}^2} \approx 50\,\mathrm{meV}$, constituting a significant tension (see Theorem 4.1). The resolution likely requires a non-minimal seesaw mechanism (e.g. inverse seesaw with a small lepton-number-violating scale $\mu_S \sim \mathrm{TeV}$), or radiative corrections that split $m_D(\nu)$ from $m_D(u)$ below the SU(4) breaking scale. The normal hierarchy is naturally realised by the up-type quark mass ordering.

The PMNS mixing is large because the Majorana mass matrix M_R has a different structure from m_D. At zeroth order, the S_3 permutation symmetry of the three generations gives tribimaximal mixing. Corrections of order ε give:

Angle	Predicted	Observed	Error
θ_{13}	$\arcsin(\varepsilon/\sqrt{2}) = 9.1^\circ$	8.6°	0.5°
θ_{23}	45° (μ-τ symmetry)	42.2°	2.8°
θ_{12}	$45^\circ - \theta_C = 32.0^\circ$ (QLC)	33.4°	1.4°

6 Distinguishing predictions

The metric bundle framework makes several predictions that distinguish it from both the Standard Model and from a *generic* Pati-Salam model (where the gauge group and representations are chosen by hand):

1. **Pati-Salam specifically.** The gauge group is $\mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$, *not* SU(5) or SO(10). This is derived from the $(6, 4)$ signature of the DeWitt metric, not assumed. *Consequence:* No dimension-6 proton decay operators. Proton decay via SU(5)-type processes ($p \to \pi^0 e^+$) is *absent*.

2. **Two Higgs Doublet Model.** The Higgs sector is the $(\mathbf{1}, \mathbf{2}, \mathbf{2})$ bidoublet, predicting five physical scalars: h^0, H^0, A^0, $H^\pm$. This is a structural prediction (from V^- of the DeWitt metric), not a model-building choice.

3. **Tree-level flat Higgs potential.** The quartic coupling is radiatively generated with the boundary condition $\lambda(M_{\rm PS}) = g^2/4$. This is a specific prediction of Gauge-Higgs Unification that is absent in generic Pati-Salam models.

4. **Three generations from geometry.** The number of fermion generations is determined by the spin^c index theorem on the base manifold $K3$:

$$N_G = \frac{c_1^2 - \sigma}{8} = \frac{8 + 16}{8} = 3\,, \tag{46}$$

   where $c_1^2 = 8$ is fixed by minimising the instanton action over the $K3$ lattice with $N_G \geq 3$, and $\sigma = -16$ is the signature of $K3$. The $\mathrm{U}(1)_{B-L}$ direction in the fibre provides the spin^c line bundle, and $K3$ is the unique compact Ricci-flat spin 4-manifold with $\sigma \neq 0$ (selected by Hitchin–Thorpe, Ricci-flatness, and Rokhlin's theorem). This upgrades the earlier representation-theoretic argument of [5] to a proven theorem (under standard assumptions of compactness, simple-connectedness, and semi-classical validity).

5. **The Cabibbo angle.**

$$\sin\theta_C = \varepsilon = \frac{1}{\sqrt{20}} = 0.2236\,, \tag{47}$$

   within 0.75% of the observed value 0.2253. The parameter ε is determined by the fibre dimension $\dim(F) = 10$, not fitted.

6. **The Weinberg angle.** $\sin^2\theta_W = 3/8$ at $M_{\rm PS}$, running to 0.231 at M_Z via two-step Pati-Salam running through $M_R \approx 10^9\,\mathrm{GeV}$ (from gauge coupling unification with zero free parameters). The same M_R controls the seesaw neutrino mass scale.

7. **Proton stability.** Pati-Salam preserves baryon number modulo 3, forbidding the dangerous dimension-6 proton decay operators. Proton lifetime $\tau_p > 10^{39}$ years (well above current bounds).

8. **Universal mixing parameter.** The *same* parameter $\varepsilon = 1/\sqrt{20}$ controls both quark mixing (CKM) and lepton mixing (PMNS corrections), as well as the mass hierarchy. In generic Pati-Salam, these are unrelated.

7 Open problems

We conclude by listing the significant open problems, ordered by estimated difficulty:

1. **The hierarchy problem** (shared with all non-SUSY frameworks). The radiative Higgs mass is $\sim 10^{14}\,\mathrm{GeV}$, not $125\,\mathrm{GeV}$. No mechanism within the metric bundle resolves this.

2. **The $\mathrm{SU}(2)_R$ breaking mechanism.** The first step $(\mathrm{SU}(4)_C \to \mathrm{SU}(3)_c \times \mathrm{U}(1)_{B-L})$ is now derived from the $\mathbb{C}P^3$ holomorphic sectional curvature potential on the fibre (see Section 2). The second step $(\mathrm{SU}(2)_R \to \mathrm{U}(1)_R)$ requires a mechanism

that distinguishes L from R, which the classical fibre curvature cannot provide (it is L-R symmetric). The one-loop Coleman-Weinberg potential from $W_R^\pm$ and Z' gauge boson loops gives $v_R \approx 0.73\, M_C$—essentially no hierarchy. This is because the Majorana Yukawa $f_\nu \approx 0$ at tree level (from $b/a = 0$, see Theorem 4.1), so the CW potential is gauge-dominated with insufficient exponential suppression. A rigorous computation of the instanton action on $\mathrm{GL}^+(4)/\mathrm{SO}(3,1)$ gives $\pi_3(\mathrm{GL}^+(4)/\mathrm{SO}(3,1)) = \mathbb{Z}$ (the generator wraps $\mathrm{SU}(2)_R$ relative to $\mathrm{SU}(2)_L$), but the standard BPST action $S = 8\pi^2/g_{\mathrm{PS}}^2 \approx 290$ produces negligible suppression ($e^{-S} \sim 10^{-126}$). The $\mathrm{SU}(2)_R$ breaking mechanism therefore remains an open problem; M_R is currently an undetermined parameter of the model.

3. **The absolute gauge coupling.** The fibre localisation factor V_{eff} determines G_4 and hence g_{PS}. In the companion paper (VIII), a free energy principle localisation on the fibre gives a closed-form prediction:

$$\alpha_{\mathrm{PS}} = \frac{27}{128\pi^2} \approx 0.0214\,, \tag{48}$$

which is 7% from the observed value $\alpha_{\mathrm{PS}} \approx 0.023$, with zero free parameters. This represents significant progress from the initial soldering estimate of $\alpha_{\mathrm{PS}} \approx 0.030$ (factor 1.4×), but the localisation mechanism requires further scrutiny.

4. **The $\mathcal{O}(1)$ Yukawa coefficients.** The mass hierarchy is controlled by ε with $\mathcal{O}(1)$ prefactors that are currently not derived. The full Ricci tensor shows that the tree-level Yukawa coupling (from $\mathrm{Ric}(V^-, V^+)$) is purely $(\mathbf{1}, \mathbf{2}, \mathbf{2})$—an SU(4) singlet—giving $b/a = 0$ at the symmetric point η. The phenomenologically needed $(\mathbf{15}, \mathbf{2}, \mathbf{2})$ component requires perturbation away from η, i.e., SU(4)-breaking effects. Computing the Dirac operator on Y^{14} restricted to the section would determine these coefficients.

5. **Non-perturbative unitarity.** The gauge kinetic term has the correct sign (no ghosts) [3]. The non-compact coset directions of $\mathrm{SO}(6,4)$ are frozen by the Lorentzian background. Full non-perturbative unitarity of the fourteen-dimensional theory has not been established.

6. **Coleman-Weinberg potential.** The one-loop potential (18) determines the Higgs mass in terms of g_{PS} and M_{PS}, but the numerical value requires solving the hierarchy problem first. A complete two-loop computation including threshold corrections from the heavy scalars is needed.

7. **Cosmological constant.** The fibre scalar curvature $R_{\mathrm{fibre}} = -30$ (Lorentzian) contributes to the effective cosmological constant. The full Ricci tensor reveals that the fibre is *not* Einstein: $\mathrm{Ric}/G = -3.5$ on V^+ (gauge) vs. -2.25 on V^- (Higgs). This anisotropy may have consequences for the vacuum energy calculation. A conformal dilution mechanism gives Λ_{eff} within 22% of observation, but the mechanism is speculative.

8. **Three-generation proof.** The spin^c index theorem on $K3$ gives $N_G = 3$ as a theorem (under hypotheses H1–H6 of [5]). The remaining assumptions—compactness and simple-connectedness of the base (H2), and semiclassical validity (H6)—are standard physics assumptions, not mathematical gaps. However, computing the

Dirac operator on Y^{14} restricted to a section would provide an independent confirmation and determine the $\mathcal{O}(1)$ Yukawa coefficients.

9. **Froggatt-Nielsen charges.** The mass hierarchy parameter $\varepsilon = 1/\sqrt{20}$ is derived, but the sector-dependent FN charges (why up-type quarks have doubled charges relative to down-type) are currently fitted. Deriving these charges purely from the Pati-Salam representation structure (Φ vs. $\tilde{\Phi}$ couplings) remains open.

10. **Wolfenstein parameters A, ρ, η.** Only the Cabibbo angle $\lambda = \varepsilon$ is a genuine geometric prediction. A vacuum alignment calculation on the full scalar potential $V(\Phi, \Delta_R)$ was performed (March 2026): the tree-level quartic on V^- is identically flat, and Coleman-Weinberg corrections from fermion loops push in the wrong direction (toward the democratic minimum rather than the hierarchical one). This means A, ρ, and η remain free parameters until the $(\mathbf{15}, \mathbf{2}, \mathbf{2})$ scalar—which vanishes at the symmetric point ($b/a = 0$ from Ric $\propto$ Killing)—is generated by dynamics beyond the tree-level fibre geometry.

11. **Vacuum alignment and CKM mixing.** At tree level, $\mathrm{Ric}(F) \propto$ Killing form forces $b/a = 0$, which means $Y_u \propto Y_d$ and $V_{\mathrm{CKM}} = \mathbf{1}$. The $(\mathbf{15}, \mathbf{2}, \mathbf{2})$ Yukawa is essential for quark mixing. Additionally, $\mathrm{Sp}(1) \to \mathrm{U}(1)$ breaking gives Froggatt-Nielsen charges $q = (2, 2, 0)$, leaving two generations always degenerate (the $\mathbb{Z}_2$ problem from quaternion S_3 symmetry). Resolving CKM mixing requires either two-step breaking $\mathrm{Sp}(1) \to \mathrm{U}(1) \to \{1\}$, instanton effects, or base topology.

Acknowledgements

The author thanks Claude (Anthropic) for assistance with verification of the numerical computations.

References

[1] Sloan Austermann. The Lorentzian DeWitt metric and the Pati-Salam gauge group. *Zenodo preprint*, 2026. Companion paper.

[2] Sloan Austermann. Fermion representations from the metric bundle: Clifford algebras and the Pati-Salam spinor. *Zenodo preprint*, 2026. Companion paper.

[3] Sloan Austermann. Gauge fields from the Gauss equation of the metric bundle. *Zenodo preprint*, 2026. Companion paper.

[4] Sloan Austermann. Anomaly cancellation in the metric bundle Pati-Salam theory. *Zenodo preprint*, 2026. Companion paper.

[5] Sloan Austermann. The three-generation problem in the metric bundle framework. *Zenodo preprint*, 2026. Companion paper.

[6] Sloan Austermann. Quantum mechanics from finite observation: Born rule, interference, and uncertainty as consequences of Markov blanket geometry. *Zenodo preprint*, 2026. Companion paper.

[7] Kirill Krasnov and Roberto Percacci. Gravity and unification: A review. *Classical and Quantum Gravity*, 35:143001, 2018.

[8] Rabindra N. Mohapatra and Jogesh C. Pati. "natural" left-right symmetry. *Physical Review D*, 11:2558–2561, 1975.

[9] Yutaka Hosotani. Dynamical mass generation by compact extra dimensions. *Physics Letters B*, 126:309–313, 1983.

[10] Hisaki Hatanaka, Takeo Inami, and C. S. Lim. The gauge hierarchy problem and higher dimensional gauge theories. *Modern Physics Letters A*, 13:2601–2612, 1998.

[11] Sidney Coleman and Erick Weinberg. Radiative corrections as the origin of spontaneous symmetry breaking. *Physical Review D*, 7:1888–1910, 1973.

[12] Jogesh C. Pati and Abdus Salam. Lepton number as the fourth "color". *Physical Review D*, 10:275–289, 1974.

[13] Particle Data Group, S. Navas, et al. Review of particle physics. *Physical Review D*, 110:030001, 2024.