Fermion Representations from the Metric Bundle: Clifford Algebras and the Pati-Salam Spinor

Sloan Austermann

Independent researcher

sloan@omnisciences.org

March 5, 2026

Abstract

In a companion paper, we showed that the DeWitt supermetric on the space of Lorentzian metrics over four-dimensional spacetime has signature $(6, 4)$, yielding the Pati-Salam gauge group $\mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ as the maximal compact subgroup of $\mathrm{SO}(6, 4)$. In this paper, we derive the fermion content of the theory from the Clifford algebra of the positive-norm sector of the normal bundle. The Clifford algebra $\mathrm{Cl}(\mathbb{R}^6) \otimes \mathbb{C} \cong M_8(\mathbb{C})$ acts on an eight-dimensional spinor $\mathbb{C}^8$, which decomposes under the chirality operator as $\mathbb{C}^8 = S^+ \oplus S^- = \mathbf{4} \oplus \bar{\mathbf{4}}$ of $\mathrm{SU}(4) \cong \mathrm{Spin}(6)$. We prove that the $\mathrm{SU}(3)$ subalgebra defined as the centraliser of a complex structure J in $\mathfrak{so}(6)$ coincides exactly with the $\mathrm{SU}(3)$ from the Pati-Salam breaking $\mathrm{SU}(4) \to \mathrm{SU}(3) \times \mathrm{U}(1)_{B-L}$, with the $\mathrm{U}(1)$ charge reproducing the baryon-minus-lepton number in the ratio $+1/3 : -1$ (triplet : singlet). Combined with the $(\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$ spinor of $\mathrm{Spin}(4) \cong \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$, the full representation is the $\mathbf{16}$ of $\mathrm{Spin}(10) \supset \mathrm{Spin}(6) \times \mathrm{Spin}(4)$: one complete generation of Standard Model fermions including a right-handed neutrino, with all hypercharges correctly reproduced by $Y = (B-L)/2 + T_{3R}$.

Keywords: Clifford algebra, Pati-Salam model, fermion representations, metric bundle, Spin(10) spinor

Contents

1 Introduction **3**

2 The Clifford algebra of the positive-norm sector **3**
2.1 Setup . . . 3
2.2 Gamma matrices and bivectors . . . 4

3 The exceptional isomorphism $\mathrm{Spin}(6) \cong \mathrm{SU}(4)$ **4**
3.1 Chirality and the $4 \oplus \bar{4}$ decomposition . . . 4

4 The complex structure and $\mathrm{SU}(3) \times \mathrm{U}(1)$ **4**
4.1 The Hodge complex structure . . . 4
4.2 The centraliser of J in $\mathfrak{so}(6)$ . . . 5

5 Equivalence of the Clifford and Pati-Salam $\mathrm{SU}(3)$ **5**

6 The 4 decomposition under $\mathrm{SU}(3) \times \mathrm{U}(1)_{B-L}$ **6**

7 One complete generation: the 16 of $\mathrm{Spin}(10)$ **6**
7.1 Combining the two sectors . . . 6
7.2 Standard Model fermions . . . 6
7.3 Hypercharge verification . . . 7

8 The complex structure and chirality **7**

9 The three-generation problem **8**

10 Discussion **8**
10.1 Summary of results . . . 8
10.2 Combined with Paper I . . . 9
10.3 Open problems . . . 9

1 Introduction

The Pati-Salam gauge group $\mathrm{SU}(4)\times\mathrm{SU}(2)_L\times\mathrm{SU}(2)_R$ was derived in [1] from the geometry of the metric bundle $Y^{14} = \mathrm{Met}(X^4)$ over a four-dimensional Lorentzian spacetime. The key result is that the DeWitt supermetric [2] evaluated on a Lorentzian background has signature $(6, 4)$ on the ten-dimensional fibre of symmetric two-tensors. The normal bundle of a metric section $g\colon X \hookrightarrow Y$ therefore has structure group $\mathrm{SO}(6, 4)$, whose maximal compact subgroup is $\mathrm{SO}(6) \times \mathrm{SO}(4) \cong \mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$.

In this companion paper, we address the fermion sector: what representations of the gauge group arise naturally from the metric bundle geometry? We show that the Clifford algebra of the six-dimensional positive-norm sector provides exactly one generation of Standard Model fermions, with all quantum numbers correctly determined.

The key results are:

1. The Clifford algebra $\mathrm{Cl}(\mathbb{R}^6, g_+) \otimes \mathbb{C} \cong M_8(\mathbb{C})$ of the positive-norm sector gives a spinor $\mathbb{C}^8$.
2. Under the exceptional isomorphism $\mathrm{Spin}(6) \cong \mathrm{SU}(4)$, the chirality decomposition $\mathbb{C}^8 = S^+ \oplus S^-$ corresponds to $\mathbf{4} \oplus \bar{\mathbf{4}}$ of $\mathrm{SU}(4)$.
3. A complex structure J on $\mathbb{R}^6$ (from the Hodge star on 2-forms) defines $\mathrm{SU}(3) \subset \mathrm{SU}(4)$ as the centraliser of J in $\mathrm{SO}(6)$, and this $\mathrm{SU}(3)$ is identical to the $\mathrm{SU}(3)$ from Pati-Salam breaking $\mathrm{SU}(4) \to \mathrm{SU}(3) \times \mathrm{U}(1)_{B-L}$.
4. The $\mathrm{U}(1)$ charge on $S^+ = \mathbf{4}$ has the pattern $\{+1/3, +1/3, +1/3, -1\}$, matching baryon-minus-lepton number.
5. Combined with the $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ spinor from $\mathrm{Spin}(4)$, the full representation is the $\mathbf{16}$ of $\mathrm{Spin}(10)$: one complete SM generation including ν_R.

Relation to division algebra approaches. The use of Clifford algebras to derive Standard Model fermion representations has a long history, notably in the work of Furey [3,4], Dixon [5], and Stoica [6], who use $\mathrm{Cl}(6)$ or the division algebras $\mathbb{C}\otimes\mathbb{H}\otimes\mathbb{O}$ to obtain one generation. Our approach differs in that the $\mathrm{Cl}(6)$ arises *geometrically* from the positive-norm sector of the DeWitt metric on the space of Lorentzian metrics, rather than being postulated algebraically. The complex structure J that breaks $\mathrm{SO}(6) \to \mathrm{U}(3)$ has a natural geometric origin: it is induced by the Hodge star on 2-forms, which is parallel with respect to the Levi-Civita connection.

2 The Clifford algebra of the positive-norm sector

2.1 Setup

The Lorentzian DeWitt metric on $S^2(\mathbb{R}^{3,1})$ has signature $(6, 4)$. The six positive-norm directions span a subspace $V^+ \cong \mathbb{R}^6$ with a positive-definite metric. The Clifford algebra of this subspace is

$$\mathrm{Cl}(V^+) = \mathrm{Cl}(6, 0) \cong M_8(\mathbb{R}) \,, \tag{1}$$

the algebra of real 8×8 matrices. Its complexification is

$$\mathrm{Cl}_6(\mathbb{C}) = \mathrm{Cl}(6, 0) \otimes_{\mathbb{R}} \mathbb{C} \cong M_8(\mathbb{C}) \,, \tag{2}$$

which has a unique irreducible module $\mathbb{C}^8$.

2.2 Gamma matrices and bivectors

We construct $\mathrm{Cl}(6,0)$ explicitly using the standard tensor product construction. Let $\sigma_x, \sigma_y, \sigma_z$ be the Pauli matrices and $I_2 = \mathbf{1}_2$. The six gamma matrices are:

$$\begin{aligned} \gamma_1 &= \sigma_x \otimes I_2 \otimes I_2\,, & \gamma_2 &= \sigma_y \otimes I_2 \otimes I_2\,, \\ \gamma_3 &= \sigma_z \otimes \sigma_x \otimes I_2\,, & \gamma_4 &= \sigma_z \otimes \sigma_y \otimes I_2\,, \\ \gamma_5 &= \sigma_z \otimes \sigma_z \otimes \sigma_x\,, & \gamma_6 &= \sigma_z \otimes \sigma_z \otimes \sigma_y\,. \end{aligned} \tag{3}$$

These satisfy $\{\gamma_i, \gamma_j\} = 2\delta_{ij}\,\mathbf{1}_8$ (verified numerically to machine precision).

The *bivectors*

$$\Sigma_{ij} = \frac{i}{2}\,\gamma_i\gamma_j\,, \qquad 1 \le i < j \le 6\,, \tag{4}$$

are 15 Hermitian 8×8 matrices that generate $\mathfrak{so}(6) \cong \mathfrak{su}(4)$ in the spinor representation. The algebra closure $[\Sigma_{ij}, \Sigma_{kl}] \subset \mathrm{span}\{\Sigma_{mn}\}$ is verified with leakage $< 10^{-15}$.

3 The exceptional isomorphism $\mathrm{Spin}(6) \cong \mathrm{SU}(4)$

3.1 Chirality and the $4 \oplus \bar{4}$ decomposition

The chirality (volume) element of $\mathrm{Cl}(6,0)$ is

$$\Gamma_7 = (-i)^3\,\gamma_1\gamma_2\gamma_3\gamma_4\gamma_5\gamma_6\,, \tag{5}$$

normalised so that $\Gamma_7^2 = \mathbf{1}_8$. It commutes with all bivectors Σ_{ij} and therefore defines a decomposition of the spinor space:

$$\mathbb{C}^8 = S^+ \oplus S^-\,, \qquad S^\pm = \ker(\Gamma_7 \mp \mathbf{1})\,. \tag{6}$$

Both S^+ and S^- are four-dimensional (verified numerically).

Proposition 3.1. *Under the exceptional isomorphism* $\mathrm{Spin}(6) \cong \mathrm{SU}(4)$*:*

$$S^+ \cong \mathbf{4} \quad (\textit{fundamental})\,, \qquad S^- \cong \bar{\mathbf{4}} \quad (\textit{anti-fundamental})\,. \tag{7}$$

The 15 bivectors Σ_{ij}*, restricted to* S^+*, generate* $\mathfrak{su}(4)$ *in the fundamental representation.*

Proof. The bivectors Σ_{ij} are traceless Hermitian 8×8 matrices that commute with Γ_7 and hence preserve the $S^\pm$ decomposition. Their restrictions to S^+ are traceless Hermitian 4×4 matrices spanning a 15-dimensional Lie algebra. Since $\dim \mathfrak{su}(4) = 15$ and the matrices satisfy the $\mathfrak{so}(6)$ commutation relations, the restriction to S^+ realises $\mathfrak{su}(4)$ in the fundamental representation. That S^- carries $\bar{\mathbf{4}}$ follows from complex conjugation. □

4 The complex structure and $\mathrm{SU}(3) \times \mathrm{U}(1)$

4.1 The Hodge complex structure

The positive-norm sector $V^+ \cong \mathbb{R}^6$ carries a natural complex structure from the Hodge star operator. In four-dimensional Riemannian geometry, the Hodge star on 2-forms satisfies $*^2 = +1$ and decomposes $\Lambda^2(\mathbb{R}^4) = \Lambda^2_+ \oplus \Lambda^2_-$ into self-dual and anti-self-dual

2-forms, each of dimension 3. Identifying V^+ with the six-dimensional Weyl sector of the normal bundle, the Hodge star induces a complex structure $J\colon V^+ \to V^+$ with $J^2 = -\mathbf{1}$, making $V^+ \cong \mathbb{C}^3$.

In the Clifford algebra, J is represented by

$$J = \Sigma_{12} + \Sigma_{34} + \Sigma_{56}\,, \tag{8}$$

which pairs the gamma matrices into complex pairs: (γ_1, γ_2), (γ_3, γ_4), (γ_5, γ_6).

4.2 The centraliser of J in $\mathfrak{so}(6)$

Definition 4.1. The *centraliser* of J in $\mathfrak{so}(6)$ is

$$\mathfrak{u}(3) = \ker(\mathrm{ad}_J) = \{X \in \mathfrak{so}(6) : [X, J] = 0\}\,. \tag{9}$$

Proposition 4.2. $\dim \mathfrak{u}(3) = 9$. *The centraliser decomposes as* $\mathfrak{u}(3) = \mathfrak{su}(3) \oplus \mathfrak{u}(1)$, *where* $\mathfrak{u}(1) = \mathbb{R} \cdot J$ *and* $\dim \mathfrak{su}(3) = 8$. *The complement* $\mathfrak{m} = \{X \in \mathfrak{so}(6) : JX + XJ = 0\}$ *has dimension 6.*

Proof. We compute ad_J as a linear map on the 15-dimensional space $\mathfrak{so}(6)$ (spanned by the bivectors Σ_{ij}) and find its kernel. The matrix of ad_J in the bivector basis has rank 6, so $\dim \ker(\mathrm{ad}_J) = 15 - 6 = 9$. The generator J itself lies in the kernel and generates a $\mathfrak{u}(1)$ factor. Projecting out J, the remaining 8-dimensional subspace is verified to close under the Lie bracket with leakage $< 10^{-15}$, confirming it is $\mathfrak{su}(3)$. □

This is verified numerically: $\dim \ker(\mathrm{ad}_J) = 9$, and after projecting out J, exactly 8 linearly independent generators remain, closing under commutation to machine precision.

5 Equivalence of the Clifford and Pati-Salam SU(3)

The Clifford route to SU(3) (centraliser of J in SO(6)) and the Pati-Salam route (SU(3) $\subset$ SU(4) from the breaking $\mathrm{SU}(4) \to \mathrm{SU}(3) \times \mathrm{U}(1)_{B-L}$) produce the *same* subgroup.

Theorem 5.1 (SU(3) equivalence)**.** *Let J be the complex structure* (8)*. Under the exceptional isomorphism* $\mathfrak{so}(6) \cong \mathfrak{su}(4)$*:*

1. *J maps to a diagonal generator of* $\mathfrak{su}(4)$*, proportional to* $\mathrm{diag}(1, 1, 1, -3)$*.*
2. $\ker(\mathrm{ad}_J) \cong \mathfrak{u}(3) = \mathfrak{su}(3) \oplus \mathfrak{u}(1)$*, where* SU(3) *is the standard upper-left embedding* $\mathrm{SU}(3) \hookrightarrow \mathrm{SU}(4)$*.*
3. *The* U(1) *factor generated by J is* $\mathrm{U}(1)_{B-L}$ *(baryon minus lepton number).*

Therefore $\mathrm{SU}(3)_{\mathrm{Clifford}} = \mathrm{SU}(3)_{\mathrm{PS}}$*.*

Proof. (1) Under the isomorphism $\mathfrak{so}(6) \cong \mathfrak{su}(4)$, the bivectors Σ_{ij} restricted to $S^+ \cong \mathbf{4}$ give 4×4 Hermitian matrices generating $\mathfrak{su}(4)$. The restriction of $J = \Sigma_{12}+\Sigma_{34}+\Sigma_{56}$ to S^+ is a diagonal matrix with eigenvalues $\{+1/2, +1/2, +1/2, -3/2\}$ (up to normalisation), which is proportional to $\mathrm{diag}(1, 1, 1, -3)$.

(2) The centraliser of $\mathrm{diag}(1, 1, 1, -3)$ in SU(4) consists of block-diagonal matrices $\left(\begin{smallmatrix} A & 0 \\ 0 & e^{i\theta} \end{smallmatrix}\right)$ with $A \in \mathrm{SU}(3)$, $e^{i\theta} = (\det A)^{-1}$. This is $\mathrm{U}(3) = \mathrm{SU}(3) \times \mathrm{U}(1)$, where SU(3) acts on the first three components. This is the standard Pati-Salam breaking $\mathrm{SU}(4) \to \mathrm{SU}(3)_c \times \mathrm{U}(1)_{B-L}$.

(3) We verify the U(1) charge structure computationally. The eigenvalues of J restricted to S^+ are $\{+1/2, +1/2, +1/2, -3/2\}$ (numerically: $\{0.5, 0.5, 0.5, -1.5\}$). The ratio of the singlet charge to the triplet charge is $-3/2 \div 1/2 = -3$. With the standard normalisation $B-L = (2/3) \cdot (\text{eigenvalue of } J|_{S^+})$, the charges are:

$$\text{triplet: } B-L = +\tfrac{1}{3}\,, \qquad \text{singlet: } B-L = -1\,, \tag{10}$$

matching quarks ($B = 1/3$, $L = 0$) and leptons ($B = 0$, $L = 1$).

Computational verification: the 8 $\mathfrak{su}(3)$ generators on $\mathbb{C}^8$ restrict to 8 linearly independent generators on $S^+ \cong \mathbf{4}$, all commuting with $J|_{S^+}$. The subalgebra closure on S^+ has leakage $< 10^{-15}$. □

6 The 4 decomposition under $\mathrm{SU}(3) \times \mathrm{U}(1)_{B-L}$

Proposition 6.1. *Under* $\mathrm{SU}(3) \times \mathrm{U}(1)_{B-L} \subset \mathrm{SU}(4)$*:*

$$\mathbf{4} = \mathbf{3}_{+1/3} \oplus \mathbf{1}_{-1}\,, \qquad \bar{\mathbf{4}} = \bar{\mathbf{3}}_{-1/3} \oplus \mathbf{1}_{+1}\,. \tag{11}$$

Proof. The quadratic Casimir $C_2(\mathrm{SU}(3)) = \sum_{a=1}^{8} T_a^2$ restricted to $S^+ = \mathbf{4}$ has eigenvalues $\{0, c, c, c\}$ where $c = 4/3$ in our normalisation. The three eigenvectors with $C_2 = 4/3$ span the $\mathbf{3}$ of SU(3); the eigenvector with $C_2 = 0$ is the SU(3) singlet. The $\mathrm{U}(1)_{B-L}$ charges, read off from $J|_{S^+}$, are $+1/3$ on the triplet and -1 on the singlet. □

The decomposition of $S^- = \bar{\mathbf{4}}$ follows by charge conjugation: $\bar{\mathbf{3}}_{-1/3} \oplus \mathbf{1}_{+1}$.

Physically:

- The $\mathbf{3}_{+1/3}$ comprises three quark colors with $B - L = +1/3$.
- The $\mathbf{1}_{-1}$ is a lepton with $B - L = -1$.
- This is the defining feature of the Pati-Salam model: leptons as the "fourth color" [7].

7 One complete generation: the 16 of Spin(10)

7.1 Combining the two sectors

The full normal bundle has $V = V^+ \oplus V^- \cong \mathbb{R}^{6,4}$. The positive-norm sector $V^+ \cong \mathbb{R}^6$ gives $\mathrm{Spin}(6) \cong \mathrm{SU}(4)$ with spinors $S^\pm = \mathbf{4} \oplus \bar{\mathbf{4}}$. The negative-norm sector $V^- \cong \mathbb{R}^4$ gives $\mathrm{Spin}(4) \cong \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ with spinors $(\mathbf{2}, \mathbf{1}) \oplus (\mathbf{1}, \mathbf{2})$.

The total spin group is $\mathrm{Spin}(6) \times \mathrm{Spin}(4) \subset \mathrm{Spin}(10)$, and the Spin(10) Weyl spinor decomposes as

$$\mathbf{16} = (\mathbf{4}, \mathbf{2}, \mathbf{1}) \oplus (\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2}) \tag{12}$$

under $\mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$.

7.2 Standard Model fermions

Breaking further via $\mathrm{SU}(4) \to \mathrm{SU}(3)_c \times \mathrm{U}(1)_{B-L}$:

$$(\mathbf{4}, \mathbf{2}, \mathbf{1}) \to (\mathbf{3}, \mathbf{2}, \mathbf{1})_{+1/3} \oplus (\mathbf{1}, \mathbf{2}, \mathbf{1})_{-1}\,, \tag{13}$$

$$(\bar{\mathbf{4}}, \mathbf{1}, \mathbf{2}) \to (\bar{\mathbf{3}}, \mathbf{1}, \mathbf{2})_{-1/3} \oplus (\mathbf{1}, \mathbf{1}, \mathbf{2})_{+1}\,. \tag{14}$$

After $\mathrm{SU}(2)_R \to \mathrm{U}(1)_R$ breaking and forming $Y = (B-L)/2 + T_{3R}$:

SM representation	$\mathrm{SU}(3)_c \times \mathrm{SU}(2)_L \times \mathrm{U}(1)_Y$	Y	Particle
Q_L	$(\mathbf{3}, \mathbf{2}, +1/6)$	$+1/6$	(u_L, d_L)
u_R	$(\mathbf{3}, \mathbf{1}, +2/3)$	$+2/3$	up-type quark
d_R	$(\mathbf{3}, \mathbf{1}, -1/3)$	$-1/3$	down-type quark
L_L	$(\mathbf{1}, \mathbf{2}, -1/2)$	$-1/2$	(ν_L, e_L)
e_R	$(\mathbf{1}, \mathbf{1}, -1)$	-1	charged lepton
ν_R	$(\mathbf{1}, \mathbf{1}, 0)$	0	right-handed neutrino

This is *one complete generation* of Standard Model fermions, with all 16 Weyl spinors and the correct hypercharges.

Remark 7.1 (Right-handed neutrino)**.** The **16** of Spin(10) automatically includes a right-handed neutrino ν_R, which is a singlet under the SM gauge group. This is a generic prediction of Pati-Salam and SO(10) models, and the metric bundle framework inherits it. The existence of ν_R is consistent with neutrino oscillation data, which require non-zero neutrino masses that can be naturally accommodated via a seesaw mechanism with $M_{\nu_R} \sim M_{\mathrm{PS}}$.

7.3 Hypercharge verification

The hypercharge formula $Y = (B-L)/2 + T_{3R}$ is verified for each fermion:

Fermion	$B-L$	T_{3R}	Y	Check
u_L	$+1/3$	0	$+1/6$	✓
d_L	$+1/3$	0	$+1/6$	✓
u_R	$+1/3$	$+1/2$	$+2/3$	✓
d_R	$+1/3$	$-1/2$	$-1/3$	✓
ν_L	-1	0	$-1/2$	✓
e_L	-1	0	$-1/2$	✓
ν_R	-1	$+1/2$	0	✓
e_R	-1	$-1/2$	-1	✓

All eight Standard Model hypercharges are correctly reproduced. The formula $Y = (B-L)/2+T_{3R}$ is not imposed by hand—it follows from the embedding $\mathrm{U}(1)_Y \subset \mathrm{SU}(4) \times \mathrm{SU}(2)_R$ via the Pati-Salam structure.

8 The complex structure and chirality

The complex structure J on the positive-norm sector V^+ has a natural geometric origin: the Hodge star operator $*$ on 2-forms in four-dimensional Riemannian geometry.

Proposition 8.1. *The Hodge star $*$ on $\Lambda^2(T^*X)$ satisfies $\nabla * = 0$, where ∇ is the Levi-Civita connection. Therefore, the decomposition $\Lambda^2 = \Lambda^2_+ \oplus \Lambda^2_-$ (self-dual/anti-self-dual) is* parallel*: the Levi-Civita connection preserves the splitting.*

Proof. The Hodge star is defined from the metric g and the volume form ϵ. Both are parallel with respect to ∇: $\nabla g = 0$ (metric compatibility) and $\nabla \epsilon = 0$ (consequence of metric compatibility and torsion-freeness). Therefore $\nabla * = 0$. □

The complex structure J induced by $*$ on the normal bundle is therefore parallel: $\nabla^{\perp} J = 0$. This ensures that $\mathrm{SU}(3) \subset \mathrm{SO}(6)$ is a *true gauge symmetry*, not merely a pointwise artifact—the holonomy of the normal connection preserves the U(3) structure.

Remark 8.2 (Chirality from orientation)**.** The choice of J versus $-J$ (equivalently, the choice of which sector is self-dual versus anti-self-dual) is determined by the orientation of X. The orientation fixes which SU(2) factor is "left" and which is "right." This provides a geometric origin for parity violation: the chirality of the weak interaction is tied to the orientation of spacetime.

9 The three-generation problem

The Clifford algebra construction gives exactly *one* generation. This is because $\mathrm{Cl}_6(\mathbb{C}) \cong M_8(\mathbb{C})$ has a unique irreducible module $\mathbb{C}^8$ (by the Artin-Wedderburn theorem), and there is no algebraic mechanism for replication.

The observation of three generations of fermions in nature remains unexplained within the metric bundle framework. We briefly discuss possible approaches.

Index theorem. If fermions are zero modes of a Dirac operator $\not{D}_Y$ on Y restricted to the section, the number of generations is

$$n_{\mathrm{gen}} = \mathrm{index}(\not{D}_Y\big|_{\sigma_g(X)}) = \int_X \hat{A}(TX) \wedge \mathrm{ch}(N)\,, \tag{15}$$

where N is the normal bundle. If this index equals 3 for topological reasons (depending on the topology of X and the Chern classes of N), three generations would be topologically protected. Computing this index requires specifying the topology of spacetime.

Complex structures. The space of compatible complex structures on $\mathbb{R}^6$ is $\mathrm{SO}(6)/\,\mathrm{U}(3) \cong \mathbb{C}P^3$, with $\pi_2(\mathbb{C}P^3) = \mathbb{Z}$. Topologically distinct complex structures could give rise to distinct generations.

Division algebras. In the Dixon-Furey approach [3, 5], three generations arise from the three imaginary quaternionic units in $\mathbb{C} \otimes \mathbb{H} \otimes \mathbb{O}$. Whether this mechanism can be embedded in the metric bundle framework remains an open question.

Status. The three-generation problem is the most significant open problem in the metric bundle programme. Its resolution (or the demonstration of its irresolvability) would substantially affect the viability assessment of the framework.

10 Discussion

10.1 Summary of results

We have shown that the Clifford algebra of the positive-norm sector of the DeWitt metric naturally produces one complete generation of Standard Model fermions:

1. $\mathrm{Cl}_6(\mathbb{C})$ gives a spinor $\mathbb{C}^8 = \mathbf{4} \oplus \bar{\mathbf{4}}$ of $\mathrm{SU}(4) \cong \mathrm{Spin}(6)$.
2. A parallel complex structure J breaks $\mathrm{SU}(4) \to \mathrm{SU}(3) \times \mathrm{U}(1)_{B-L}$.

3. The SU(3) from J is *identical* to the Pati-Salam SU(3) (proven).
4. $B - L$ charges are $+1/3$ (quarks) and -1 (leptons)—correct.
5. Combined with $\mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ from Spin(4): the **16** of Spin(10).
6. All SM hypercharges reproduced via $Y = (B-L)/2 + T_{3R}$.
7. Right-handed neutrino ν_R is automatically included.

10.2 Combined with Paper I

Together with the results of Paper I [1], the metric bundle framework now accounts for:

- **Gauge group:** $\mathrm{SU}(4) \times \mathrm{SU}(2)_L \times \mathrm{SU}(2)_R$ (Pati-Salam) from the DeWitt signature $(6, 4)$.
- **Coupling unification:** $g_4 = g_L = g_R$ from equal Dynkin indices, giving $\sin^2 \theta_W = 3/8 \to 0.231$.
- **Higgs sector:** $(\mathbf{1}, \mathbf{2}, \mathbf{2})$ bidoublet from the negative-norm modes (Two Higgs Doublet Model).
- **Fermion content:** One generation of 16 Weyl fermions with correct quantum numbers (this paper).
- **Gravity:** Correct Einstein-Hilbert, Yang-Mills, and torsion terms from the Gauss equation.

10.3 Open problems

The principal open problems, in order of severity, are:

1. **Three generations.** Paper V [8] proposes that three inequivalent complex structures on $\mathbb{R}^6$ give $N_G = 3$, consistent with the anomaly constraint $N_G \equiv 0 \pmod 3$ from Paper IV [9]; see Section 9 for discussion.

2. **Hierarchy problem** ($m_H \ll M_{\mathrm{PS}}$; no resolution).

3. **Fermion masses** (tree-level Yukawa couplings are degenerate due to (1) quaternionic symmetry; mass hierarchy requires explicit breaking).

4. **Anomaly cancellation.** Paper IV [9] verifies that all gauge and gravitational anomalies cancel for Pati-Salam with $N_G = 3$, and that the Witten SU(2) anomaly is absent.

5. **CKM/PMNS matrices** (require inter-generation mixing, which depends on the symmetry-breaking mechanism for the (1) flavour symmetry).

6. **Quantum mechanics.** Paper VI [10] derives the Born rule, interference, and uncertainty from the $(6, 4)$ DeWitt signature via finite-agent observation geometry; implications for the fermion sector remain to be explored.

Acknowledgements

The author thanks Claude (Anthropic) for assistance with the numerical computations and cross-checks.

References

[1] Sloan Austermann. The Lorentzian DeWitt metric and the Pati-Salam gauge group. *Zenodo preprint*, 2026. Companion paper.

[2] Bryce S. DeWitt. Quantum theory of gravity. I. The canonical theory. *Physical Review*, 160:1113–1148, 1967.

[3] Cohl Furey. Standard model physics from an algebra? 2016.

[4] Cohl Furey. Three generations, two unbroken gauge symmetries, and one eight-dimensional algebra. *Physics Letters B*, 785:84–89, 2018.

[5] Geoffrey M. Dixon. *Division Algebras: Octonions, Quaternions, Complex Numbers and the Algebraic Design of Physics*. Kluwer Academic, 1994.

[6] Ovidiu Cristinel Stoica. The standard model algebra – leptons, quarks, and gauge from the complex clifford algebra Cl_6. *Advances in Applied Clifford Algebras*, 28:52, 2018.

[7] Jogesh C. Pati and Abdus Salam. Lepton number as the fourth "color". *Physical Review D*, 10:275–289, 1974.

[8] Sloan Austermann. The three-generation problem in the metric bundle framework. *Zenodo preprint*, 2026. Companion paper.

[9] Sloan Austermann. Anomaly cancellation in the metric bundle Pati-Salam theory. *Zenodo preprint*, 2026. Companion paper.

[10] Sloan Austermann. Quantum mechanics from finite observation: Born rule, interference, and uncertainty as consequences of Markov blanket geometry. *Zenodo preprint*, 2026. Companion paper.