

# Chapter 1

## Functions

In calculus, one of the first thing a student has to understand is the concept of functions. Yet, it is one of the most overlooked part of high-school mathematics.

### 1.1 What is a function?

#### **Definition 1 - Function.**

A function is a relation between a set of inputs to a set of outputs. For a given input, it has one and only one particular output

It is often written in the following form –  $y = f(x)$ , where  $x$  is the input and  $y$  is the output.

The letters  $x$ ,  $y$ , and  $f$  are arbitrary placeholders. You can substitute these with anything. For many real-world applications, they are meant to be substituted in order to obtain tangible values. In fact, they can be substituted with a different set of placeholders without losing the intended meaning.

Let us look at a function that doubles the input.

$$y = f(x) = 2x \tag{1.1}$$

Here,  $x$  can take any number. It could be an integer, say,  $x = 2$ . Then,

$$y = f(2) = 2 \times 2 = 4$$

What if  $x$  is a complex number, say,  $x = 5 + 3i$ ? In that case,

$$y = f(5 + 3i) = 2 \times (5 + 3i) = 10 + 6i$$

It must be noted that, no matter what we plug-in as  $x$ , the output (or  $y$ ) is the same for a given value of  $x$ .

Let us consider a very different kind of function. What if we want to compute the number of unique letters a word has? Let us call this function  $c$ .

$$y = c(x) = \text{number of unique letters in } x \quad (1.2)$$

Here are some computed cases for  $x = \text{"HELLO"}$ ,  $x = \text{"OVER"}$ , and  $x = \text{"WATERMELON"}$  respectively.

$$y = c(\text{"HELLO"}) = 4$$

$$y = c(\text{"OVER"}) = 4$$

$$y = c(\text{"WATERMELON"}) = 9$$

Just like the earlier example, we observe that the output for a given input is fixed.  $c(\text{"Hello"})$  will never be 2, 3, 5, 6, or anything other than 4.

There is another aspect that must be highlighted. The input to this function cannot be anything under the sun. For example,  $c(x)$  makes no sense when  $x$  is a real number, a complex number, or even a word from the Indian languages. This restriction to the type of input that a function can accept, gives rise to the domain of the function.

### Definition 2 - Domain of a function.

The domain of a function is the set of input values for which the function is defined.

The output of  $c(x)$  will invariably be a positive integer. You can't have it return a floating point number. I can't think of any word whose unique letter count is 3.14156. Can you?

This brings us to the definition of codomain.

### Definition 3 - Codomain.

Codomain is the set of into which all of the output of the function is constrained to fall.

Clearly, the codomain of  $c(x)$  in equation (1.2) is the set of natural numbers. Or, as they are often denoted in mathematics,  $y \in \mathbb{N}$ .

The description of a function may not be so simple. The choice of the function's form may depend on the input itself. Let us look at an example.

$$y = f(x) = \begin{cases} x, & \text{if } x \geq 0 \\ 0, & \text{otherwise} \end{cases} \quad (1.3)$$

For this function, if you insert a number that is negative, the output is 0. The graph of the function looks like this –

:insert graph here:

These kind of functions have a name. They are called conditionals.

## 1.2 Understanding discontinuities

Conditionals do not always behave well. You can easily construct a conditional that is broken. Let us look at one such case.

$$y = f(x) = \begin{cases} 5, & \text{if } x > 4 \\ 0, & \text{if } x < 4 \end{cases} \quad (1.4)$$

Clearly, the function tells us nothing about a situation when we feed it  $x = 4$  as input. We can argue that 4 is not the domain of the function as it is not described by the function itself.

Fair enough. let us rectify the function in equation (1.4) and re-define the first part of the conditional with a  $\geq$  sign instead of a  $>$  sign.

$$y = f(x) = \begin{cases} 5, & \text{if } x \geq 4 \\ 0, & \text{if } x < 4 \end{cases} \quad (1.5)$$

The curve looks like this –

:insert graph here:

We now have a complete definition of the function. Still, there is a nagging problem. What would happen if we try to approach the value from left. Let us assume a situation where you are asked to run towards  $x = 4$  from  $x = 0$  and asked to halve your speed as you get closer and closer to  $x = 4$ . (Let's say, there is a force field at  $x=4$ .) You would never reach  $x = 4$  in any finite time, but you would get close, very, very close. You would be at what mathematically is referred to as  $x = 4-$ . The value of  $y$  you will observe from here would be 0. Similarly, if you did the same exercise while running from  $x = 8$  to  $x = 4$ , you would reach a point that can be written as  $x = 4+$ . Here you would observe that the value of the function is 5.

This is a type of discontinuity. There is also a name for such kind of discontinuity. It's called a jump discontinuity.

What about the following function? What happens when we plug-in  $x = 1$ ?

$$y = g(x) = \frac{1}{x-1} \quad (1.6)$$

:insert graph here:

Studying its graph, we notice that the function is not defined at  $x = 1$ . This is not because someone has goofed around with the function at  $x = 1$  but by the very nature of  $g(x)$ . The output vanishes at  $-\infty$  at  $x = 1-$  and reappears at  $+\infty$  at  $x = 1+$  leaving no way for us to figure out what exactly happens at  $x = 1$ . There is a name for this kind of discontinuity – asymptotic discontinuity. There is another type of discontinuity that must be mentioned. It is known as a point discontinuity. Let us take the following function –

$$y = f(x) = \begin{cases} 5, & \text{if } x > 4 \\ 0, & \text{if } x = 4 \\ 5, & \text{if } x < 4 \end{cases} \quad (1.7)$$

:insert graph here:

This is a diabolical function. If you approach it from the right ( $x = 4+$ ), you land up with a value of 5. If you approach it from left ( $x = 4-$ ), you also land up with a value of 5. However, if you parachute and land exactly at  $x = 4$ , you obtain a value of 0!

We will encounter the concept of discontinuity as we dwelve deeper into calculus. In fact, study of discontinuities is an important aspect of this subject.