EASY-TO-USE AI CALCULATOR PREDICTING 5Y WEIGHT TRAJECTORIES AFTER BARIATRIC SURGERY: A SOPHIA STUDY



Patrick Saux¹, Pierre Bauvin², Violeta Raverdy², Philippe Preux¹, François Pattou².








¹Inria, Univ. Lille, CNRS, Centrale Lille, UMR 9198-CRIStAL, F-59000, Lille, France. ²Univ. Lille, Inserm, CHU Lille, Institut Pasteur de Lille, U1190 - EGID, F-59000, Lille, France.

Goal: predict the outcome of surgery




Results


- 7 (simple) attributes are predictive of post-surgery weight loss:
 - Weight (preop),
 - •Height,
 - •Age,
 - Type of intervention,
 - Type II diabetes (T2D),
 - Duration of T2D,
 - •Smoking.

- **A** Lower weight loss with
 - •Age,
 - •SG (after 1y),
 - •T2D,
 - Longer (more severe) T2D.





A companion tool for patients and care providers




https://bwtp.univ-lille.fr


- Individualized trajectory (dotted line),
 - "Green zone": where the majority of patients are (IQR),
 - Preop: visualize expected weight loss,
 - **Postop**: flag complications (patient out of green zone).

An international study


Validation: 8 cohorts + 2 RCT studies 10,000 patients, 3 continents




Impact of RYGB vs SG



Impact of T2D



Validation



MAD: accuracy in BMI points for a standard patient.