

All 16 models, modes R0-R2 — left: do we find the same root VALUES? right: do we NUMBER them the same way?
Are the roots right?
root-finder tolerance
3layer_normal R0
3layer_normal R1
3layer_normal R2
grad_smooth R0
grad_smooth R1
grad_smooth R2
grad_steep R0
grad_steep R1
grad_steep R2
grad_sqrt_fine R0
grad_sqrt_fine R1
grad_sqrt_fine R2
lvz_weak R0
lvz_weak R1
lvz_weak R2
lvz_strong R0
lvz_strong R1
lvz_strong R2
lvz_deep R0
lvz_deep R1
lvz_deep R2
lid_over_channel R0
lid_over_channel R1
lid_over_channel R2
grad_with_lvz R0
grad_with_lvz R1
grad_with_lvz R2
tailings_dry R0
tailings_dry R1
tailings_dry R2
tailings_saturated R0
tailings_saturated R1
tailings_saturated R2
lvz_saturated R0
lvz_saturated R1
lvz_saturated R2
2layer_normal_split R0
2layer_normal_split R1
2layer_normal_split R2
near_homogeneous_split R0
near_homogeneous_split R1
near_homogeneous_split R2
37 of 42 curves within the root-finder tolerance; only 2 exceed 0.1 m/s
10^{-5}
10^{-4}
10^{-3}
10^{-2}
10^{-1}
10^{0}
worst |Δc| to the nearest disba root (m/s)
Is the mode numbering the same?
disba duplicates roots here
disba duplicates roots here
disba duplicates roots here
disba duplicates roots here
0
20
40
60
80
100
periods where the mode index matches disba (%)