Editorial 🖰



PRACTICE

Submissions

COMPETE IOBS LEADERBOARD

Leaderboard

Practice > Interview Preparation Kit > Greedy Algorithms > Minimum Absolute Difference in an Array

Minimum Absolute Difference in an Array ☆

Consi	der an array of integers, $m{arr} = [m{arr}[0], m{arr}[1], \dots, m{arr}[n]$	– $1]].$ We define the absolute difference between two elements, $oldsymbol{a[i]}$
and a	$m{\imath}[j]$ (where $m{i} eq m{j}$), to be the absolute value of $m{a}[m{i}] - m{a}[m{j}]$.	

Discussions

Given an array of integers, find and print the minimum absolute difference between any two elements in the array. For example, given the array arr = [-2, 2, 4] we can create 3 pairs of numbers: [-2, 2], [-2, 4] and [2, 4]. The absolute differences for these pairs are |(-2)-2|=4, |(-2)-4|=6 and |2-4|=2. The minimum absolute difference is 2.

Function Description

Problem

Complete the minimumAbsoluteDifference function in the editor below. It should return an integer that represents the minimum absolute difference between any pair of elements.

minimumAbsoluteDifference has the following parameter(s):

- n: an integer that represents the length of arr
- · arr: an array of integers

Input Format

The first line contains a single integer **n**, the size of **arr**.

The second line contains n space-separated integers arr[i].

Constraints

- $2 \le n \le 10^5$
- $-10^9 \le arr[i] \le 10^9$

Print the minimum absolute difference between any two elements in the array.

Sample Input 0

3 -7 0

Sample Output 0

3

With n=3 integers in our array, we have three possible pairs: (3,-7), (3,0), and (-7,0). The absolute values of the differences between these pairs are as follows:

- |3 -7| ⇒ 10
- |3 − 0| ⇒ 3
- |-7 0| ⇒ 7

Notice that if we were to switch the order of the numbers in these pairs, the resulting absolute values would still be the same. The smallest of these possible absolute differences is 3.

Sample Input 1

Sample Output 1

Explanation 1

The smallest absolute difference is |-54--53|=1.

Sample Input 2

1 -3 71 68 17

Sample Output 2

Explanation 2

The minimum absolute difference is |71 - 68| = 3

Author shashank21i Difficulty Easy Max Score 15 Submitted By 49592 NEED HELP? View discussions ♥ View top submissions RATE THIS CHALLENGE **公公公公公** MORE DETAILS ■ Download problem statement ■ Download sample test cases Suggest Edits Ff 🟏 in

```
▼ 50 € 7 €
                                                        JavaScript (Node.js)
  1
      'use strict':
 2
     const fs = require('fs');
 3
     process.stdin.resume();
     process.stdin.setEncoding('utf-8');
     let inputString = '';
     let currentLine = 0;
 10
     process.stdin.on('data', inputStdin => {
 11
 12
        inputString += inputStdin;
 13
 14
 15
     process.stdin.on('end', function() {
      inputString = inputString.replace(/\s*$/, '')
 16
            .split('\n')
 17
            .map(str => str.replace(/\s*$/, ''));
 18
 19
 20
 21
     });
 23
     function readLine() {
 24
       return inputString[currentLine++];
 25
 26
     // Complete the minimumAbsoluteDifference function below.
 27
 28
     function minimumAbsoluteDifference(arr) {
 29
 30
 31
     }
 32
Run Code
                                                                             Submit Code
```

Contest Calendar | Blog | Scoring | Environment | FAQ | About Us | Support | Careers | Terms Of Service | Privacy Policy | Request a Feature