

Sum of Squares: Part 1

Apoorv Vikram Singh

February 2, 2021

Introduction

Reason two basic problems about polynomial inequalities

1. Feasibility of polynomial system.

$$\begin{aligned} p_1(\mathbf{x}) &\geq 0 \\ &\vdots \\ p_m(\mathbf{x}) &\geq 0 \end{aligned} \tag{1}$$

Is there an \mathbf{x} such that (1) is satisfied?

2. Checking non-negativity: Is $q(\mathbf{x}) \geq 0$, $\forall \mathbf{x}$ satisfying (1) ?

Justification: Feasibility

Feasibility checking is highly expressive.

- Example: MaxCut.
- . Input: $G = (V, E)$, $|V| = n$.
- . Goal: Find $S \subseteq V$ such that $|E(S, \bar{S})|$ is maximized.
- . Polynomial Feasibility: For some $\beta \in \mathbb{Z}^+$,

$$\frac{1}{4} \sum_{(i,j) \in E} (\mathbf{x}_i - \mathbf{x}_j)^2 = \beta |E|$$

$$\mathbf{x}_i^2 = 1, \quad \forall 1 \leq i \leq n.$$

- . $n + 1$ degree-2 polynomials. Enumerate over polynomial number of values of β and solve MaxCut.

Other examples: MaxClique, Max 3-SAT, Knapsack. Therefore, polynomial feasibility checking problem is *NP*-Hard.

Goal

- ▶ Analyze a relaxation for the feasibility problem, and try to find interesting situations where one can get a poly-time algorithms.

Justification: Non-Negativity

- ▶ Checking positivity: Given $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ with rational coefficients, decide if:
 - $f \geq 0$, $\forall \mathbf{x} \in \{-1, 1\}^n$, or,
 - find an $\mathbf{x} \in \{-1, 1\}^n$ such that $f(\mathbf{x}) \leq 0$.
- ▶ Example MaxCut: Decide if $\text{MaxCut} \leq c$.
 - . Let $f_G(\mathbf{x}) \stackrel{\text{def}}{=} \frac{1}{4} \sum_{(i,j) \in E} (\mathbf{x}_i - \mathbf{x}_j)^2$.
 - . Decide if $c - f_G(\mathbf{x}) \geq 0$, $\forall \mathbf{x} \in \{-1, 1\}^n$.

Certifying Non-Negativity

Given $f : \{-1, 1\}^n \rightarrow \mathbb{R}$, find an “efficiently verifiable” certificate of non-negativity.

SoS Certificates

Definition (SoS cert of non-neg (or) SoS proof of non-neg)

A degree- d SoS certificate of non-negativity of $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ is a list of polynomials $g_1, \dots, g_r : \{-1, 1\}^n \rightarrow \mathbb{R}$, such such that

- $\deg(g_i) \leq d/2$, and
- $f(\mathbf{x}) = \sum_{i \leq r} g_i^2(\mathbf{x}), \forall \mathbf{x} \in \{-1, 1\}^n$.

Efficiently Verifiable (?)

Polynomials f, g_1, \dots, g_r are represented as a vector of coefficients.

1. How large if r ? ($\leq n^d$, see later)
2. How large are coefficients of g_i ?

Efficiently Verifiable

Proposition (Efficiently Verifiable)

Suppose $r \leq n^d$, all coefficients of g_i are bounded in magnitude by $2^{\text{poly}(n^d)}$. Then the identity $f = \sum_{i \leq r} g_i^2$ over all $\mathbf{x} \in \{-1, 1\}^n$ can be checked in $\text{poly}(n^d)$ time.

Proof.

- Given g_i , can compute g_i^2 , and $\sum_{i \leq r} g_i^2$ in polynomial time.
- Check if $(f - \sum_{i \leq r} g_i^2)(\mathbf{x}) = 0, \forall \mathbf{x} \in \{-1, 1\}^n$.
- Using the fact that coefficient vector representation is unique, just check if $f - \sum_{i \leq r} g_i^2 = \mathbf{0}$



Fact (Unique Representation)

$\forall f : \{-1, 1\}^n \rightarrow \mathbb{R}$, there exists a unique representation of f : The multi-linear representation of f

$$f = \sum_{S \subseteq [n]} \hat{f}(S) \prod_{i \in S} x_i.$$

(this representation is its Fourier transform)

\implies coefficient vector representation is unique.

(multilinear representation exists because $x_i^2 = 1$)

Are non-negative functions always certifiable?

Proposition (Certifiability of non-negative functions)

Let $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ be non-negative over $\{-1, 1\}^n$. Then, there exists a $\deg(2n)$ -SoS certificate of non-negativity.

Proof.

- Consider $g : \{-1, 1\}^n \rightarrow \mathbb{R}$, and $g(\mathbf{x}) = \sqrt{f(\mathbf{x})}$.
- Every function on $\{-1, 1\}^n$ is a polynomial of $\deg \leq n$.
- $f = g^2 \implies \deg(2n)$ -SoS Certificate.



Tensor Notation

- Suppose vector $\mathbf{v} \in \mathbb{R}^n$.
- $\mathbf{v}^{\otimes 2} \in \mathbb{R}^{n^2}$, where $\mathbf{v}(i, j) = \mathbf{v}_i \mathbf{v}_j$.
- $\mathbf{v}^{\otimes k} \in \mathbb{R}^{n^k}$.

Proving Efficient Verifiability

Theorem (PSD Matrices and SoS Certificates)

$f : \{-1, 1\}^n \rightarrow \mathbb{R}$ has a $\deg(d)$ -SoS certificate of non-negativity iff there exists a matrix A such that $A \succcurlyeq 0$, and

$$f(\mathbf{x}) = \left\langle (1, \mathbf{x})^{\otimes \frac{d}{2}}, A \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle .$$

► Parsing Notation:

- $(1, \mathbf{x}) \in \mathbb{R}^{(n+1)}$.
- $(1, \mathbf{x})^{\otimes \frac{d}{2}}$: populate in a vector all possible monomials in the variable \mathbf{x} of degree at most $d/2$.
- $A \in \mathbb{R}^{(n+1)^{d/2} \times (n+1)^{d/2}}$.

Proof

- If Part:

$$\begin{aligned} f(\mathbf{x}) &= \left\langle (1, \mathbf{x})^{\otimes \frac{d}{2}}, A \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle \\ &= \left\langle (1, \mathbf{x})^{\otimes \frac{d}{2}}, (B^\top B) \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle \\ &= \left\langle B \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}}, B \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle \\ &= \left\| B \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\|^2. \end{aligned} \tag{2}$$

- Let $g_i(\mathbf{x}) = \left\langle e_i, B \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle$, i.e., i -th entry of the vector.
- B is a matrix of constants, applied to monomials of degree at most $d/2$, therefore, $\deg(g_i) \leq d/2$.
- Therefore,

$$f(\mathbf{x}) = \sum_{i=1}^{(n+1)^{\otimes d/2}} g_i^2(\mathbf{x}).$$

Proof Cont...

- Only if Part: Suppose f has a degree- d SoS certificate.

$$f = \sum_{i \leq r} g_i^2, \text{ and } \underbrace{g_i(\mathbf{x})}_{\deg \leq d/2} = \left\langle \mathbf{v}_i, (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle.$$

$$\begin{aligned} f(\mathbf{x}) &= \sum_{i \leq r} \left\langle \mathbf{v}_i, (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle^2 \\ &= \left\langle (1, \mathbf{x})^{\otimes \frac{d}{2}}, \underbrace{\left(\sum_{i \leq r} \mathbf{v}_i \mathbf{v}_i^\top \right)}_{\succeq 0} \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle. \end{aligned}$$



Efficient Verifiability

Corollary (Bound on r)

If $f : \{-1, 1\}^n \rightarrow \mathbb{R}$ has a degree- d SoS certificate, then it has a certificate with $r \leq (n+1)^{d/2}$.

Proof.

Follows from (2):

$$f(\mathbf{x}) = \left\| B \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\|^2.$$



Efficient Verifiability

Lemma (Bit-complexity of SoS proofs)

Suppose f has a degree- d certificate over $\{-1, 1\}^n$. Then, f has a degree- d SoS certificate with bit complexity $\text{poly}(n^d)$.

- ▶ $\{-1, 1\}^n$ is important, and doesn't necessarily hold for other domains.

Proof.

Since we are given f , we know that it can be efficiently represented. Therefore, we try to bound the entries of A in terms of f .

- $f(\mathbf{x}) = \left\langle (1, \mathbf{x})^{\otimes \frac{d}{2}}, A \cdot (1, \mathbf{x})^{\otimes \frac{d}{2}} \right\rangle$, for some A .
- $f = \sum_u \hat{f}_u \mathbf{x}_u$, where $\mathbf{x}_u = \prod_{i \in u} \mathbf{x}_i$.
- Expanding the inner product, we see $\hat{f}_u = \sum_{S, T} A_{S, T}$, such that $S \Delta T = u$, and $|S|, |T| \leq d/2$.
- $\hat{f}_\emptyset = \sum_S A_{S, S} = \text{tr}(A) = \sum_i \underbrace{\lambda_i(A)}_{\geq 0}$.
- $\|A\|_F^2 = \sum_{S, T} A_{S, T}^2 = \sum_i \lambda_i^2(A) \leq \hat{f}_\emptyset^2$.



References I



Mitra, A. (2015).

Super-resolution, extremal functions and the condition number of vandermonde matrices.

In *Proceedings of the Forty-Seventh Annual ACM Symposium on Theory of Computing*, STOC '15, page 821–830, New York, NY, USA. Association for Computing Machinery.