

# TNF $\alpha$ and the TNF Receptor Superfamily: Structure-Function Relationship(s)

HAITHAM T. IDRIS AND JAMES H. NAISMITH\*

*Centre for Biomolecular Sciences, The University, St. Andrews, Fife KY16 9ST, Scotland, United Kingdom*

**KEY WORDS** TNF $\alpha$ ; receptor; cytokine; apoptosis; crystal; structure

**ABSTRACT** Tumour Necrosis Factor  $\alpha$  (TNF $\alpha$ ), is an inflammatory cytokine produced by macrophages/monocytes during acute inflammation and is responsible for a diverse range of signalling events within cells, leading to necrosis or apoptosis. The protein is also important for resistance to infection and cancers. TNF $\alpha$  exerts many of its effects by binding, as a trimer, to either a 55 kDa cell membrane receptor termed TNFR-1 or a 75 kDa cell membrane receptor termed TNFR-2. Both these receptors belong to the so-called TNF receptor superfamily. The superfamily includes FAS, CD40, CD27, and RANK. The defining trait of these receptors is an extra cellular domain comprised of two to six repeats of cysteine rich motifs. Additionally, a number of structurally related “decoy receptors” exist that act to sequester TNF molecules, thereby rescuing cells from apoptosis. The crystal structures of TNF $\alpha$ , TNF $\beta$ , the extracellular domain of TNFR-1 (denoted sTNFR-1), and the TNF $\beta$  sTNFR-1 complex have been defined by crystallography. This article will review the structure/function relationships of the TNF $\alpha$  and the TNF receptor superfamily. It will also discuss insights as to how structural features play a role in the pleiotropic effects of TNF $\alpha$ . *Microsc. Res. Tech.* 50:184–195, 2000. © 2000 Wiley-Liss, Inc.

## BACKGROUND

Hints of the existence of a biological factor mediating tumour necrotic activity date back to the 18th century following observations that cancer patients, exposed to severe bacterial infection, underwent occasional shrinkage of their tumours (Coley, 1893). Lymphotoxin was discovered in 1968 as a soluble protein produced by T-lymphocytes (Granger and Williams, 1968). In 1975, experiments demonstrated that a protein, termed “Tumour Necrosis Factor” (TNF $\alpha$ ), was released into the circulation of animals subsequent to stimulation of their reticuloendothelial system and lipopolysaccharide challenge (Carswell et al., 1975). This protein has been demonstrated to cause rapid necrotic regression of certain forms of tumours (hence its name). The necrotic ability of TNF $\alpha$  did not prove true for all tumour types and the therapeutic potential of TNF $\alpha$  was hampered by its systemic toxicity at effective concentrations. This discriminatory TNF $\alpha$  mediated cytotoxicity may be related to the tumour cell’s ability to undergo nitrotyrosinylated-tubulin mediated microtubule dysfunction (Idriss, 2000). However, the specific necrotic effects of TNF $\alpha$  on tumour cells were increased by the presence of Interferon (Fiers, 1991). Both TNF $\alpha$  and lymphotoxin (termed TNF $\beta$  and also as LT $\alpha$ ) define two structurally and functionally related proteins (Aggarwal et al., 1991; Armitage, 1994; Bazzoni and Beutler, 1996; Beyaert and Fiers, 1994; Fiers, 1991; Vilcek and Lee, 1991). We use the abbreviation TNF to refer to both proteins. Both proteins were isolated in 1984 from activated macrophages and T cells, respectively (Gray et al., 1984; Pennica et al., 1984). Their cDNA sequence revealed 30% homology in the encoded amino acid residues. The structure of TNF $\alpha$  and TNF $\beta$  have been determined and both proteins were found as a “pear”/

“cone” shaped trimers (Fig. 1; Eck and Sprang, 1989; Eck et al., 1992; Jones et al., 1989). TNF receptors, TNFR-1 and TNFR-2, were cloned and expressed as 55 and 75 kDa proteins, respectively (Dembic et al., 1990; Loetscher et al., 1990; Schall et al., 1990; Smith et al., 1990). The X-ray structure of TNF $\beta$  in complex with TNFR-1 provided a break through in the molecular understanding of how TNF functions (Fig. 2; Banner et al., 1993). The TNF trimer binds three receptor molecules, one at each of three TNF monomer-monomer interfaces (Fig. 3). The extracellular domain of the receptor is an elongated molecule composed of 3 disulphide-containing 40 residue motifs (Banner et al., 1993). Only three of four extracellular modules of TNFR-1 were visualised. The first structure of the receptor on its own also contained disorder in the C-terminus but did hint at a possible mechanism for signal transduction (Naismith et al., 1995). This was followed up by a high-resolution structure of the entire extracellular domain of receptor (Naismith et al., 1996). This work determined the structure of the fourth motif, which is different in structure to the other three. This led to the novel method of aligning and predicting the structure of related receptors (Naismith and Sprang, 1998). Such an approach has been recently used in modelling the CD40/CD40L interaction (Singh et al., 1998).

Contract grant sponsor: Hoescht Marion Roussel (France).

\*Correspondence to: James H. Naismith, Centre for Biomolecular Sciences, The University, St. Andrews, Fife KY16 9ST, Scotland, UK. E-mail: naismith@st-and.ac.uk (JHN), hi@st-and.ac.uk (HTI).

Received 30 September 1999; accepted in revised form 29 December 1999




Fig. 1. Van der Waals surface of the lymphotoxin-TNF $\beta$  (*left*) and the TNF $\alpha$  (*right*) trimers. Subunits are coloured white, grey, and charcoal. Mutagenic "hot spots" associated with changes in cytotoxicity or receptor binding, are shown at the front in grey. On TNF $\beta$ , these are Tyr-108 (on white subunit) and Asp-50 (on black subunit). In TNF $\alpha$ , mutagenesis of residues 84–91 (on white subunit), 30–40 (lower group on charcoal subunit), and 143–149 (upper group on charcoal subunit) can alter cytotoxicity. Reproduced from Eck et al. (1992) with permission of the publisher. (Note: color version of artwork is available online.)

### BIOLOGICAL ROLE(S) OF TNF $\alpha$

The biological functions of TNF $\alpha$  are varied and the mechanism of action is somewhat complex. This protein, conferring resistance to certain type of infections on the one hand and causing pathological complications on the other, carries out contradictory roles (Fiers, 1991). This may be related to the varied signaling pathways that are activated. Interestingly, mutant phenotype of knock-out mice lacking the gene for TNF $\alpha$  showed no resistance to leishmania and failed to form germinal centres in their spleens (Goldfeld and Tsai, 1996). Moreover, TNFR knockout mice displayed an increased susceptibility to microbial infection and a suppressed inflammatory response upon challenge with bacterial endotoxins (Acton et al., 1996; Steinshamn et al., 1996). This clearly points to an important role for TNF $\alpha$  in conferring immunity and a proper inflammatory response.

TNF $\alpha$  plays several therapeutic roles within the body, which include immunostimulation, resistance to infection agents, resistance to tumours (Aggarwal et al., 1991; Vilcek and Lee, 1991), sleep regulation (Krueger et al., 1998), and embryonic development (Wride and Sanders, 1995). On the other hand, parasitic, bacterial and viral infections become more pathogenic or fatal due to TNF circulation (Fiers, 1991). However, the major role of TNF seems to be as an important mediator in resistance against such infections. Indeed, in experimental leishmaniasis, insufficient TNF $\alpha$  is associated with progressive disease and death (Goldfeld and Tsai, 1996). It seems that only abhorrent situations, such as overreaction of the host or dysfunction of natural autoregulatory networks, lead to the aforementioned deleterious effects (Fiers, 1991). Many cell types infected with viruses become sensitised to TNF effects implicating this cytokine in viral disease (Czarniecki, 1993). TNF may contribute towards resistance of infection through activation of neutrophils and platelets, enhancement of macro-





Fig. 2. **A:** Complete structure of the extracellular domain of TNFR-1. Reproduced from Naismith et al. (1996) with permission of the publisher. **B:** Van der Waals surface of the TNF $\beta$  trimer (magenta) complexed with its receptor (purple); water molecules are shown in blue. From Banner et al. (1993), prepared using Rasmol. (Note: color version of artwork is available online.)

phage/NK cell killing abilities, and stimulation of the immune system (Fiers, 1991). TNF also plays a possible pathological role in a number of autoimmune diseases such as graft vs. host rejection, and rheumatoid arthritis (Beutler and Bazzoni, 1998; Beutler, 1999). Additionally, TNF shows anti-malignant cell cytotoxicity especially in combination with Interferon (Fiers, 1991; Gruss and Dower, 1995). Such a combination can cure aggressive non-immunogenic tumours in animal models. However, as high concentrations of TNF $\alpha$  are toxic to the host (maximum tolerated dose in humans 200  $\mu\text{g}/\text{m}^2$ ), it is necessary to increase the therapeutic index either by decreasing toxicity (e.g., pre-treatment with IL-1) or by increasing effectiveness. This may be possible through mutations that reduce systematic cytotoxicity and increase TNF's effectiveness in selectively eliminating tumour cells (Cha et al., 1998). Efficacy may be heightened with some chemotherapeutic drugs (Balkwill, 1992). So far clinical trials using human TNF $\alpha$  or TNF $\alpha$ /Interferon  $\gamma$ , reported no success (Hieber and Heim, 1994).

TNF $\alpha$  is also involved in physiologic sleep regulation. Administration of exogenous TNF $\alpha$  was shown to induce increased sleep, whilst inhibition of TNF $\alpha$  reduced spontaneous sleep. TNF $\alpha$  mRNA is produced in diurnal rhythm in brain with the highest levels occurring during peak sleep periods. Absence of the TNFR-1 receptors resulted in reduced sleep in animal models. Similar effects were observed with the cytokine IL-1 (Shoham et al., 1987). TNF mediated apoptosis also seems a part of normal embryonic development (Wride and Sanders, 1995).

TNF can induce necrotic or apoptotic cell death (Beyaert and Fiers, 1994). Necrosis is characteristic with cell swelling, organelle destruction, and cell lysis.



Apoptosis is perceived as a conserved and biochemically driven process of cell death with distinct morphological features (Steller, 1995). Apoptosis is characterised by cell shrinkage, formation of apoptotic bodies, and, typically, formation of inter nucleosomal DNA fragmentation. Necrosis is observed with the necrotic haemorrhage of transplantable methylcholanthrene-induced sarcomas in animal models (Beyaert and Fiers, 1994). Apoptosis is seen as a general TNF mediated cytotoxic phenomenon occurring through the TNFR-1 signalling pathway. However, it has been proposed that TNFR-2 indirectly mediates cytotoxicity through endogenous production of TNF and autotropic or paratropic activation of TNFR-1 (Grell et al., 1999). Many adaptor proteins are involved in the complex signalling pathways, which initiate apoptosis (Baker and Reddy, 1996, 1998; Yuan, 1997). However, the full signalling network leading to, and controlling, apoptosis remains to be elucidated. Activation of proteases is postulated as a general mechanism of apoptosis (Thornberry and Lazebnik, 1998). TNF $\alpha$  can also lead to cell survival through activation of NF- $\kappa$ B. Many TNF related ligands have been identified thus far (Armitage, 1994; Bazzoni and Beutler, 1996; Orlinich and Chao, 1998). TNF related proteins such as RANKL are required for (osteoclast) cell differentiation, which is necessary for bone resorption (Takahashi et al., 1999). Certain TNF-like receptors (e.g., HVEM) serve as cell entry points into cells during infection (Marsters et al., 1997; Montgomery et al., 1996).

#### MECHANISM(S) OF ACTION

TNF $\alpha$  exerts its effect(s) by binding to, as a trimer, and clustering high-affinity receptors present in great numbers on most cell membranes (Loetscher et al., 1991a; Schoenfeld et al., 1991). The ligand/receptor complex is rapidly internalised via clathrin-coated pits and ends up in secondary lysosomes where it is degraded. Binding of TNF $\alpha$  to the 75 kDa TNFR-2 is not sufficient to attain cytotoxicity (Thoma et al., 1990), but rather binding to the 55 kDa TNFR-1 is sufficient to attain TNF $\alpha$  mediated cell killing (Van Ostade et al., 1993). TNF $\alpha$  exerts its effects by activating a number of secondary proteins that provoke a variety of re-

Fig. 3. Three models describing TNF-induced receptor signalling. **A:** Trimerisation hypothesis: the juxtaposition of the three receptors results from binding of a single TNF trimer, leading to cell signalling. **B:** Expanding network hypothesis: binding of the TNF trimer to the receptor dimer generates an expanding hexagonal array of ligand/receptor complex. Each TNF trimer engages three receptor dimers. "Capping" of the receptors triggers the biological response. **C:** Molecular-switch hypothesis: each receptor dimer is an activatable unit. Receptor activation occurs in response to: (1) TNF ligand binding to subunit B of the receptor, (2) Subunit A disengaging from subunit B, resulting in the receptor binding to another part of the TNF trimer. These events lead to conformational changes (e.g., homodimerisation of the receptor's cytoplasmic "death domains"), resulting in signal transduction. The graph shows a hypothetical profile depicting the free energy associated with conformational changes between "On" and "Off" states of the receptor. A large energy barrier preventing transition from the "Off" to the "On" states, is lowered upon ligand binding. The free energy of the "On" state (defining the stable ligand/receptor complex), may be lower than the energy of the "Off" state and this may be irreversible. Reproduced from Bazzoni and Beutler (1996) with permission of the publisher. (Note: color version of artwork is available online.)






Fig. 4. Structure of the murine TNF locus. Coding portion of the exons are shaded and inset shows location of TNF $\alpha$  5' flanking region elements potentially involved in transcriptional regulation of TNF $\alpha$  gene. Reproduced from Shakhov et al. (1990), with permission of the publisher.

sponses within the cell such as activation of gene transcription and/or production of reactive oxygen or nitrogen radicals (e.g., NO). Activated proteins include G-Protein, transcription factors (e.g., NF- $\kappa$ B, AP-1), protein kinases (e.g., CK II, erk-1, erk-2, and MAP2), phospholipases (e.g. PLA<sub>2</sub>, PLC, PLD and sphingomyelinase) mitochondrial proteins (e.g., manganese superoxide dismutase), and serine and cysteine proteases, known as caspases (Beyaert and Fiers, 1994; Darnay and Aggarwal, 2000; Thornberry and Lazebnik, 1998; Vilcek and Lee, 1991). Several members of the TNF receptor superfamily possess intracellular "death domains," which are protein interaction domains each consisting of 65–80 amino acids that participate in TNF $\alpha$  mediated apoptosis. These "death domains" function to activate caspases (cysteine proteases that cleave certain proteins after specific aspartic acid residues). Caspases are recruited to the activated receptor (together with other signalling molecules) by adaptor proteins such as TRADD and FADD (for review see Ashkenazi and Dixit, 1998, 1999; Yuan, 1997). Interestingly, the TNF $\alpha$  trimer undergoes, at a relatively low pH and in the absence of receptor binding, conformational change that allows it to form a functional ion channel (Baldwin et al., 1996; Kagan et al., 1992). This shows that the molecule performs many subtle and extraordinarily diverse functions. Additionally, TNF binding mediates alterations in cell volume associated with necrosis/apoptosis by inducing regulation of the activities of a number of anion/cation channels (Lang et al., 1998).

#### CHARACTERISATION OF THE TNF $\alpha$ GENE AND MOLECULE:

The TNF $\alpha$  gene is a single copy gene, preceded by the TNF $\beta$  on the short arm of human chromosome 6 and murine chromosome 17 (Spriggs et al., 1992). It is closely linked within the major histocompatibility complex genes. TNF $\alpha$  is always 3' to the TNF $\beta$  gene and is around 3 Kb long. It consists of four exons, arranged over approximately 3 kb of DNA, interrupted by three introns. More than 80% of the mature TNF sequence is encoded in the fourth exon, while exons I and II contain almost entirely leader peptide sequence. The TNF $\alpha$  mRNA seems to be transcribed in a wide variety of cells. The 5' flanking region of the TNF $\alpha$  gene contains several  $\kappa$ B homologous sequence enhancer elements and a MHC class II like "Y box" (Fig. 4; Shakhov et al., 1990). These sequences seem to be involved in transcriptional activation. The gene also seems to be regulated at the translational level through the UA-rich sequence in the 3' untranslated region of the human TNF $\alpha$  mRNA. The mRNA translates into a 157 amino acid protein in human and 156 in murine cells (Spriggs et al., 1992; Vilcek and Lee, 1991).

TNF $\alpha$  exists in two forms, a membrane bound and a soluble form; each form possibly carries out a distinct physiological role (Beyaert and Fiers, 1994; Watts et al., 1997). Human TNF $\alpha$  is expressed as a precursor molecule of a 157 amino acid, preceded by a 76 amino acid presequence (79 in murine). This presequence is highly conserved and seems to serve to anchor the

TABLE 1. Consensus phosphorylation sites<sup>†</sup> for TNF $\alpha$  and TNFR-1hTNF $\alpha$  233 A.A (precursor molecule):

1 MSTESMIRDV ELAEEALPKK**T**\*GGPQGS\*RRCLFLSLFSLFI VAGATTLFCL LHFVGIVGPQR  
 61 EEFP**RDLS**\*LI SPLAQAVRSS\* **S**\*RT\*PS\*DKPVA HVVANPQAEGLQQLNRRAN ALLANGVELR  
 121 DNQLVVPSEG LYLIYSQVLF KGQGCPSLHV LLTHTISRIA VSYQTKVNL SAIKSPCQRE  
 181 TPEGAEAKPW YEPIYLGGVF QLEKGDRLSA EINR**PDY**\*LDF AESGQVYFGI IAL

hTNFR-1 (55):476 A.A (precursor molecule):

1 MGL**ST**\*VPDLL LPLVLELLV GIYPGSGVIGL VPHLGDKREK**DS**\*VCPQGYI HPQNNISICT  
 61 KCHKGTLYN DCPGPGQDTC CRECESGSFT ASENHLRHCL SCSKCRKEMG QVEISSCTVD  
 121 **RD****T**\*VCGCRKN QYRHYWSENLFQCFNCSLCL NGTVHL**S**\***Q****E** KQNTVCTCHA GFFLRENECV  
 181 SCSNCKKSLE CTKLCLPQIE NVKGTEDSGT TVLLPLVIFV GLCLLSLLFI GLMYRYQRW  
 241 **S**\*KLYSIVCGK **ST**\*PEKEGELE GT**T**\*TKPLAPN PSFSPTPGFT PTLGFSPVPS STFTSSSTYT  
 301 PGDCPNFAAP RREVAPPYQG ADPILATALA SDPIPPLQK WEDSAHK**PQS**\* LDTDDPATLY\*  
 361 AVVENVPPLR WKEFVRRLLGL SDHEIDRLEL QNGRCLREAQ YSMLATWRRR TPRREATLEL  
 421 LGRVLRDMDL LGCLEDIEEA LCGPAALPPA PSLR

<sup>†</sup>Phosphorylation consensus sequences are based on those described by Pearson and Kemp (1991):

CK II sites	( <b>S/T</b> XX E/DX)
PKA sites	( <b>K/R</b> XX S/T)
PKC sites	( <b>S/T</b> X R/K, R/K X <b>S/T</b> , R/K XX <b>S/T</b> )
Tyr kinase sites	(X E/D <b>Y</b> X).

\*Phosphorylatable residue.

precursor polypeptide in the membrane (Vilcek and Lee, 1991). Hence unlike TNF $\beta$ , which is secreted, TNF $\alpha$  is first produced as a type II membrane protein. Amino acids -44 to -26 of the TNF $\alpha$  presequence comprise the hydrophobic transmembrane region and residues -76 to -50 comprise the intracytoplasmic region. The unprocessed protein has a molecular mass of 26 kDa and is proteolytically cleaved (possibly with a serine protease between amino acids -14 and -1) to give a 17 kDa active form. The mature form of TNF $\alpha$  lacks any methionine residues. Processed (cleaved) TNF $\alpha$  exists in solution as a homotrimer of total molecular mass of 52 kDa (other estimates between 45–65 kDa) and this is the form that binds and cross-links receptors (Smith and Baglioni, 1987). Each of the three subunits contains a disulphide bridge, which is not essential for biological activity. Human TNF $\alpha$  binds strongly to TNF receptors with a disassociation constant  $K_d$  of around 0.5 and 0.1 nM for TNFR-1 and TNFR-2, respectively. Binding to the TNFR-1 receptor selectively mediates the toxic effects of TNF $\alpha$  (Beyaert and Fiers, 1994; Fiers, 1991; Vilcek and Lee, 1991). A TNF $\beta$  related ligand termed LT $\beta$  has been identified, which is capable of forming heterotrimer with TNF $\beta$  (LT $\alpha_2\beta$  or LT $\beta_2\alpha$ ) and binding to TNF receptors (Browning et al., 1993).

Not many post-translational modifications have been reported to occur on TNF $\alpha$ . Human TNF $\alpha$  lacks N-glycosylation, which is observed in TNF $\beta$  (Smith and Baglioni, 1992; Vilcek and Lee, 1991). The primary sequence of TNF $\alpha$  shows the presence of a number of phosphorylation consensus sites (Table 1), offering a potential mechanism for regulating trimer formation and/or mode of receptor binding. Indeed, phosphorylation of the 26-kDa membrane bound human TNF $\alpha$  has been previously reported in monocytes (Pocsik et al.,

1995). Further, a recent study demonstrated that membrane bound TNF $\alpha$  is phosphorylated by CK I and dephosphorylated in the presence of soluble dimeric TNFR-1, thereby offering a mechanism for reverse signalling. Dephosphorylation occurs through phosphatase activation as this was sensitive to phosphatase inhibitors (Watts et al., 1999).

TNF $\alpha$  exists in a more diverse range of cells than TNF $\beta$ . TNF $\alpha$  is produced in a wide variety of hematopoietic and non-hematopoietic cells, both normal and malignant. These include macrophages, CD4<sup>+</sup> and CD8<sup>+</sup> T-lymphocytes, B-lymphocytes, LAK cells, NK cells, neutrophils, astrocytes, endothelial cells, smooth muscle cells, and a number of non-hematopoietic tumour cell lines (Vilcek and Lee, 1991). Indeed, the classical cell line for testing TNF cytotoxicity is the murine fibrosarcoma cell line L929, although others (such as the WEHI164 clone) show heightened sensitivity. Some cells can be induced to produce the two forms of TNF. Many of TNF $\alpha$ 's activities are species-independent, but some are species specific (Fiers, 1991).

## TNF RECEPTORS AND THE TNF RECEPTOR SUPERFAMILY

TNF $\alpha$  binds to two cognate receptors of molecular weight 55 and 75 kDa (Loetscher et al., 1991b). Human TNFR-1 consists of 434 amino acids whilst TNFR-2 comprises 439 residues. These receptors share very limited homology in the extracellular region. The TNF receptor superfamily is defined by the presence of repeating units of cysteine clusters. No homology is seen in the intracellular regions of the proteins (reviewed in Beutler and Bazzoni, 1998). Lack of homology in the intracellular regions suggests the two receptors activate distinct signalling pathways. TNF receptors are

TABLE 2. Summary of TNF Receptor Family Members

Receptor	Other names	A.A.	M.wt. kDa	Ligand	Death domain	Reference
CAR 1	N/A	368	39	Unknown	Yes	Brojatsch et al. (1996)
CD 27	N/A	240	45–55	CD27L	No	Camerini et al. (1991); Armitage (1994)
CD 30	N/A	577	120	CD30L	No	Durkop et al. (1992); Armitage (1994)
CD 40	N/A	256	40	CD40L	No	Armitage (1994)
CD 95	FAS, APO-1	319	45	CD95L	Yes	Itoh et al. (1991); Armitage (1994)
CD134	OX 40	272 <sup>1</sup>	50	OX 40L	No	Mallett et al. (1990)
CD137	4-1BB	256	35	4-1BBL	No	Vinay and Kwon (1998)
DR 3	WSL1, TRAMP, Apo3, LARD	417	54	Apo3L	Yes	Screaton et al. (1997)
DR 4	TRAILR-1	468 <sup>2</sup>	32	Apo2L/TRAIL	Yes	Pan et al. (1997b)
DR 5	TRICK2, TRAILR-2, Apo2, KILLER	411	45	Apo2L/TRAIL	Yes	MacFarlane et al. (1997)
DR 6	N/A	655	72	Unknown	Yes	Pan et al. (1998a)
GITR	AITR	228	25	GITRL	No	Nocentini et al. (1997)
HveA	HVE1, HveM	283	30.5	TRAF	No	Marsters et al. (1997)
LT $\beta$ R	N/A	435	60	LT $\beta$	No	Crowe et al. (1994)
NGFR	N/A	425	75	NGF	Yes	Radeke et al. (1990)
RANK	TRANCE-R	616	80 <sup>3</sup>	RANKL	No	Anderson et al. (1997)
TNFR-1	CD120a, R55, R60	460	55	TNF $\alpha$ , TNF $\beta$	Yes	Loetscher et al. (1990); Armitage (1994)
TNFR-2	CD120b, R75, R80	476	75	TNF $\alpha$ , TNF $\beta$	No	Smith et al. (1990); Armitage (1994)
DcR1	TRID, TRAIL-R3, LIT	299	50	TRAIL	No	Pan et al. (1997a); MacFarlane et al. (1997)
DcR2	TRUNDD, TRAIL-R4	386	42	Apo2L	No	Pan et al. (1998b)
DcR3	N/A	300	33	FasL	No	Pitti et al. (1998)
OCIF	OPG, TR1	401	60	OPGL/RANKL	No	Tsuda et al. (1997); Simonet et al. (1997)
CrmB	N/A	355	48	TNF $\alpha$ , TNF $\beta$	No	Hu et al. (1994)
CrmC	N/A	186	25	TNF $\alpha$	No	Smith et al. (1996)

<sup>1</sup>Mouse.<sup>2</sup>Including 24 A.A. signal peptide.<sup>3</sup>60 kDa predicted (Murine).

present on almost all known cell types with few exceptions, such as erythrocytes and unstimulated T lymphocytes. Receptor density ranges from 200–10,000 per cell. There seems to be no correlation between the number of receptors present on a cell and the magnitude or direction of the TNF-induced response (Beyaert and Fiers, 1994). Soluble forms of TNFR-1 and TNFR-2 have been identified in human urine and in the serum of cancer patients. These probably regulate TNF's activity by decoying the ligand, thereby down-regulating the generated signals (Engelmann et al., 1990).

Many membrane bound receptors (including TNFR-1 and 2) have been identified thus far, forming what is known as the TNF receptor superfamily (Bazan, 1993). Their molecular weight is in the range of 50–120 kDa (Table 2). Each molecule contains several disulphide bonds, essential for activity. Members of the TNF receptor family belong to the type I transmembrane glycoproteins. They share a common cysteine-rich motif (about 40 residues) repeated two (e.g., CAR 1) to six (e.g., CD 30) times in the extracellular N-terminal domain (Fig. 2; Beutler and Bazzoni, 1998; Naismith and Sprang, 1998). The average homology in these repeats is around 25% between the various members. The receptors differ in the possession of a cytoplasmic "death domain," which is absent in a number of the many members of the TNFR family (Table 2).

All of these receptors have been cloned from various species. The mode of action of these receptors relies on being cross-linked upon binding of ligand oligomers (e.g., TNF trimers, NGF dimers). They can be activated

through cross-linking with specific monoclonal antibodies (Shalaby et al., 1990). Viruses can utilise many of these receptors as entry points to the cell. A number of related receptors termed "decoy receptors" have also been identified, which function to sequester secreted ligands (Ashkenazi and Dixit, 1999; Beutler, 1999). Decoy receptors (DcR1, DcR2, and DcR3) sequester Trail (DcR1 and DCR2) and Fas (DcR3) ligands, thereby preventing apoptosis (Ashkenazi and Dixit, 1999). In addition, a soluble non-membrane bound decoy receptor termed osteoprotegerin (OPG) has also been identified (Simonet et al., 1997).

Not many posttranslational modifications have been reported for the TNFR-1 family. TNFR-1 and TNFR-2 are both N-glycosylated, but only TNFR-2 is O-glycosylated (Fiers, 1991). All other membrane receptors are also glycosylated. Neither has intrinsic kinase activity or is known to be phosphorylated by a membrane bound or cytoplasmic kinases, but recently murine TNFR-1 has been shown to be preferentially phosphorylated by mitogen activated protein kinase (MAPK) on Thr-236 and Ser-270 (Van Linden et al., 2000) and human TNFR-1 to be phosphorylated on Tyr-331 (Darnay and Aggarwal, 1997). The primary sequence of TNFR-1 suggests the presence of a number of potential phosphorylation consensus sites for PKC or CK II (Table 1).

TNFR-1 has ubiquitous distribution present on cells such as fibroblasts and epithelial cells, whilst TNFR-2 presence is more restricted, being confined mainly to cell of hematopoietic origins (Armitage, 1994; Byffell

and Mihatsch, 1993). Certain cell types (e.g., Hep2) express only TNFR-1 (Hohmann et al., 1989). CD40 is expressed as a 50-kDa glycoprotein on a variety of cells including B cells, thymic epithelium, dendritic cells, monocytes, and at a low level on T cells (Armitage, 1994). Fas receptor is expressed on activated T cells, B cells, monocytes, and neutrophils. Fas mRNA is expressed in various organs in mice (Armitage, 1994). CD30 receptor was originally identified in Hodgkin's cells. It is also expressed on a number of non-Hodgkin's lymphoma cells as well as a several virally transformed B- and T-cell lines (Armitage, 1994). CD27 is expressed on the majority of medullary thymocytes and peripheral blood T cells. It is also present on some B and NK cells (Armitage, 1994). The receptor 4-1BB is expressed as a 35 kDa protein on activated CD4<sup>+</sup> and CD8<sup>+</sup> T cells and intestinal intraepithelial T lymphocytes and human lung epithelial cells (Armitage, 1994; Vinay and Kwon, 1999). Ox40 on the other hand was originally described as a 50-kDa protein on activated rodent CD4<sup>+</sup> T cells and thymocytes and hence shows a relatively restricted profile of expression (Armitage, 1994). CAR1 was identified as a chicken cell-surface receptor for cytopathic avian leukosis sarcoma viruses (Brojatsch et al., 1996). LT $\beta$ R is expressed in a wide range of cell types, except lymphocytes (VanArsdale et al., 1997). However, expression of its ligand is restricted to activated lymphocytes. Human GITRL mRNA is found in several peripheral tissues, and hGITRL protein is detected on cultured vascular endothelial cells. The levels of hGITRL mRNA in tissues are generally low; in peripheral blood T cells, however, antigen-receptor stimulation leads to substantial induction of hGITRL transcripts (Nocentini et al., 1997). HVEM mRNA is expressed mainly in lymphocyte-rich tissues, e.g., spleen and also in several non-lymphoid tissues (Marsters et al., 1997). DR3 is abundantly expressed on thymocytes and lymphocytes, expression is restricted predominantly to lymphoid tissues, spleen, thymus, and PBL (Screaton et al., 1997). DR4 and DR5 are present in many tissues, including spleen, thymus, peripheral blood lymphocytes (PBLs), prostate, testis, ovary, uterus, and multiple tissues along the gastrointestinal tract (Pan et al., 1997a,b; MacFarlane et al., 1997; Walczak et al., 1997). TRAIL-R3 (Dc1) is a decoy receptor (MacFarlane et al., 1997); its mRNA has been found in heart, placenta, lung, liver, kidney, spleen, peripheral blood leukocytes, and bone marrow, but at substantially lower amounts in most transformed cell lines (Frank et al., 1999; Golstein, 1997). TRAIL-R4 (DcR2), shows widespread tissue distribution, being expressed both in normal human tissue and tumour cell lines (Degli-Esposti et al., 1997). RANK is expressed on the cell surface of precursor bone cells (Anderson et al., 1997; Nakagawa et al., 1998).

### STRUCTURAL ASPECTS OF THE TNF $\alpha$ PROTEIN

The crystal structure for TNF $\alpha$  was solved almost simultaneously by two research groups (Eck and Sprang, 1989; Jones et al., 1989). Recently, the structure has been described to a very high resolution (Baeyens et al., 1999). TNF $\alpha$  exists as a trimer arranged like a triangular cone, such that each molecule contacts the other two. The interfaces between the monomers in




Fig. 5. Ribbon diagram showing the "cone-shaped" TNF $\alpha$  trimer (A, Eck et al., 1989) and two TNF $\alpha$  mutants (B, Reed et al., 1997; C, Cha et al., 1998). Mutated residues are marked with arrows. Panel (C) was reproduced from Cha et al. (1998) with permission of the publisher. (Note: color version of artwork is available online.)

this analogy would be the vertical edge of the cone. Each monomer consists of two packed  $\beta$ -pleated sheets each of eight, anti-parallel  $\beta$ -strands arranged in a  $\beta$ -jellyroll topology with an N-terminal insertion that contains three additional  $\beta$ -strands (Fig. 5A). The monomer is approximately 60Å long and 30Å wide. The outer sheet is rich in hydrophilic residues, whilst the inner sheet is hydrophobic and contains the C-terminal segment, which is located close to the trimer's central



axis. The jellyroll has a wedged shape with a broad base and a tapering peak and is reminiscent of viral capsid proteins. The "inner" sheet, hidden in the trimer complex, is formed by strands B'-B-I-D-G (based on the historical nomenclature of strands) in correct spatial order. The exposed "outer" sheet is formed by strands C'-C-H-E-F. Murine TNF has now been determined to 1.4Å (Baeyens et al., 1999); although it is materially the same as the human TNF structure, it does provide a more accurate and detailed description of side chain positions. Site-directed mutagenesis identified many residues within the TNF sequence that are necessary for the TNF's cytotoxicity and receptor binding (reviewed by Jones et al., 1991; Sprang and Eck, 1991).

The related cytokines TNF $\beta$  (Banner et al., 1993; Eck et al., 1992) and CD40L (Karpusas et al., 1995) have both been structurally characterised. As was expected, both share the same fold as TNF $\alpha$ . However, there are considerable differences in the surface properties of the three molecules. The differences were particularly marked for CD40L, which binds to CD40 and does not bind to TNFR-1 or TNFR-2.

The crystal structures of TNF mutants that show greater affinity for TNFR-1 binding compared to TNFR-2 or that have diminished binding to both receptors, have been elucidated (e.g., Cha et al., 1998; Reed et al., 1997). One TNF $\alpha$  mutant, where Arg31 was substituted with Asp, shows preferentially decreased binding to TNFR-2 (Fig. 5B). Model complexes of TNF $\alpha$  with receptors TNFR-1 and TNFR-2 suggest that Arg31 of TNF $\alpha$  forms an ionic interaction with a Glu residue in both receptors. Asp31 of the TNF $\alpha$  mutant was postulated to form hydrogen bond interactions with Ser59 or Cys70 of TNFR-1, but not in TNFR-2. The loss of the strong ionic interactions of Arg31 and the electrostatic repulsion of Asp31 explains the reduced binding of the Arg31Asp mutant to TNFR-2. The replacement of the ionic interactions by a weaker hydrogen bond interactions between Asp31 of the TNF $\alpha$  mutant and TNFR-1, compared with no interactions with TNFR-2, explains the observed preferential binding of the Arg31Asp mutant to TNFR-1 over TNFR-2 (Reed et al., 1997).

Another TNF $\alpha$  mutant, termed M3S, contained four changes: a hydrophilic substitution of Leu29Ser, two hydrophobic substitutions of Ser52Ile and Tyr56Phe, and a deletion of the N-terminal seven amino acids that is disordered in wild-type TNF $\alpha$  (Fig. 5C). The mutant exhibited 11- and 71-fold lower binding affinities for the human TNFR-1 and TNFR-2, respectively. It also showed 20 and 10 times lower in vitro cytotoxic effect and in vivo systemic toxicity, respectively, resistance to trypsin proteolysis and increased thermal stability (Cha et al., 1998). The decreased affinity and altered thermal stability were due to the Leu29Ser mutation causing substantial restructuring of the loop containing residues 29–36 into a rigid segment, resulting from the formation of intra- and intersubunit interactions. This rigidity also explained resistance to proteolysis. The Ser52Ile and Tyr56Phe mutations did not induce a noticeable conformational change. This and similarly mutated TNF molecules may be beneficial for cancer therapy, due to their reduced systematic cytotoxicity.

Recently, the structure of the TNF related ligand TRAIL (Apo2L) was independently solved by several

groups to a resolution of 2.8–1.3 Å, both in a complex with DR5 receptor (Hymowitz et al., 1999; Mongkolsapaya et al., 1999) and as a free uncomplexed trimer (Cha et al., 1999; Hymowitz et al., 2000). The structure shows the expected B-sandwich jellyroll topology and has high degree of structural homology to the other TNF family members (TNF $\alpha$ , TNF $\beta$ , and CD40L). Conflicting reports emerged concerning the presence of a unique zinc-binding site within the trimer. Hymowitz et al. (2000) reported the presence of such site, which they suggested was important for both the structure and function of the protein. However, the zinc binding site was not observed by the other investigators (Cha et al., 1999; Mongkolsapaya et al., 1999). Mutagenesis studies identified at least five different residues (Gln 205, Val 207, Tyr 216, Glu 236, and Tyr 237) within TRAIL that are essential for high-affinity receptor binding and/or apoptotic function (Hymowitz et al., 2000).

### STRUCTURE ASPECTS OF THE RECEPTOR-LIGAND COMPLEX AND THE RECEPTOR

The complex between TNF $\beta$  and the extracellular domain of TNFR-1 was the first view of the receptor (Fig. 2B). The complex confirmed cross-linking studies that TNF functioned by aggregating receptors upon binding them. The receptors are highly elongated, almost rod shaped (Fig. 2A). The receptors align their long axis parallel to each other and parallel to and in contact with the three TNF $\beta$  subunit interfaces (Banner et al., 1993). To continue with our earlier cone analogy, three rods lie along the vertical edges of the cone. The extracellular domain of the receptor can be split by sequence analysis into four similar sub-domains. Each sub-domain consists of between 30 to 40 amino acids (Banner et al., 1993; D'Arcy et al., 1993). The structure, which only visualised the N-terminal 3 and 1/2 domains, confirmed this analysis. Each subdomain contained 3 disulphide bonds, which held together three strands connected by loops. The interface between the receptor and TNF $\beta$  involved residues from two TNF monomers and from the second and third subdomain of the receptor. The interface splits into two almost separate contact surfaces. The lower region (Fig. 2B) consists of receptor residues 56 to 73, which form an extended loop like structure, which spans the interface between two TNF $\beta$  molecules. Residues 35–53 from one TNF $\beta$  subunit (denoted A) and residues 106–112 from the other (denoted C subunit) are involved in the contacts (Banner et al., 1993). Two regions of the receptor (residues 77–81 and 107–114) form the upper region and these sit in a cleft formed at the TNF $\beta$  monomer interface. Residues 82–84, 126–130, and 155–161 from A subunit and residues 97, 118–120 and 150–152 of the C subunit of TNF $\beta$  form this interface. The interface is a mixture of both directional hydrophilic contacts and hydrophobic contacts. This mix probably confers both the exquisite specificity and high affinity that characterises the receptor complex. The extracellular domains of the receptors made no contact with each other in the crystal structure. However, it seemed likely that by bringing the receptors close together this would promote aggregation below the membrane.



Shortly after the determination of the complex, the structure of the free receptor was reported at neutral pH (Naismith et al., 1995). The receptor showed extensive hinging movements between the complex and the free form (Naismith et al., 1995) but was unchanged in the overall description of the receptor structure. In this structure again only 3 and 1/2 domains were visualised. The receptor was found in the crystals as two different dimers, both with extensive contacts. This study suggested for the first time that receptors may dimerise in the absence of TNF. It was postulated that this may regulate the signalling by enforcing an "off" state and/or pre-assembling the receptors for signalling. Some biological evidence for this came from separate studies (Boldin et al., 1995). This suggested that in the absence of the extracellular domain the receptor constitutively signals. Further indirect evidence came from Remy et al. (1999), who showed that the erythropoietin receptor (unrelated in structure to the TNF receptor superfamily) dimerises in the absence of ligand and plays a role in signal regulation. Another dimer seen in the crystal structure would be capable of binding TNF $\alpha$  and may allow a 2-D array of receptor complexes to form on the cell surface (Fig. 3B). This is the so-called expanding network hypothesis (Naismith et al., 1995).

The high-resolution structure of the complete extracellular domain of the receptor was determined by Naismith et al. (1996). This structure was determined at low pH but again showed the protein to be a dimer with extensive contacts, but different to the one observed at neutral pH. The fourth module of the extracellular domain was revealed as having a very different structure than the other three modules. The structure emphasised that the elongated receptor deforms, in response to binding another molecule (either TNF $\beta$  or itself), by a series of hinging movements. These movements are very pronounced and can lead to a difference of nearly 20Å in atomic positions (Naismith et al., 1996). This led to a new method being advanced for modelling and aligning the extracellular domains of the TNF receptor superfamily. This new approach radically alters the location of certain residues and allows modelling of CD40 binding of its ligand (Singh et al., 1998). This study is particularly relevant to TNF as both the receptor and ligand share similar structures. Singh et al. (1998) highlighted the importance of the polar residues in controlling the interaction.

A consistent picture of the receptor superfamily emerges. The disulphide bonds form a structural scaffold that can be decorated with short regions of variable length and amino acid sequence. These variable regions confer specificity for the appropriate ligand. The combination of structural conservation and sequence variability allows each member of this large receptor superfamily to recognise its cognate ligand with exquisite specificity, whilst utilising a very small array of topologically distinct units. This recognition is enhanced by the inherent flexibility of the receptor molecule, which can deform to optimise its interaction with ligand through a series of hinging movements.

Most recently the structure of extracellular fragment of DR5 (residues 58–184 or 1–130) complexed with its TRAIL/Ap02L ligand (residues 91–281 or 114–281) emerged from two different groups (Hymowitz et al.,




Fig. 6. Predicted tertiary and quaternary structures of a chimeric TNF inhibitor molecule generated by linking the extracellular domain of TNFR-1 to domains CH<sub>2</sub>, CH<sub>3</sub>, and CH<sub>4</sub> of an IgG Heavy Chain. Reproduced from Bazzoni and Beutler (1996), with permission of the publisher.

1999; Mongkolsapaya et al., 1999). The complex crystallised as a hexameric unit containing three receptors and a ligand trimer. Like TNFR1, DR5 forms an extended rod-like structure consisting of three cysteine-rich domains (CRDs), rather than the four observed for TNFR1. DR5 contains seven disulfide bridges; six were present in structural regions corresponding to CRD2 and CRD3 (three each) of TNFR1 and only one in the region corresponding to the C-terminal half (last disulfide bridge) of CRD1. Binding of Trail to DR5 buries 2,750 Å<sup>2</sup> of solvent-accessible surface area (1,400–1,540 Å<sup>2</sup> from the receptor and 1,350–1,420 Å<sup>2</sup> from the ligand). The residues of contact between the receptor and ligand (i.e., interface), were located in two main patches, covering a total area of 1,790 Å<sup>2</sup> and 890 Å<sup>2</sup>, respectively. The structure suggests common structural features amongst all of the TNF receptors.

The structure of certain accessory proteins that associate with TNF receptors, such as TRAFs (Pullen et al., 1999), have also been determined. TRAF2 (TNFR associated factor 2) binds to the cytoplasmic domain of

TNF receptors such as TNFR-1 and CD40 and mediates signal transduction. The structure of TRAF2 receptor binding domain, complexed with CD40 peptid, showed that TRAF2 forms a mushroom-shaped trimer consisting of a coiled coil and a unique beta-sandwich domain (McWhirter et al., 1999). The CD40 peptide binds in an extended conformation with every side chain in contact with a complementary groove on the rim of each TRAF monomer. The trimer may pre-organise the receptors to ensure recognition of three distinct sites on the TRAF trimer (McWhirter et al., 1999).

### FUTURE PROSPECTS

The structure of many other TNF receptors should soon emerge. Elucidation of the crystal structure of the TNF related receptor, receptor activated NF $\kappa$ B (RANK), is currently underway. RANK is the sole signalling receptor for the ligand RANKL (also known as TRANCE/ODF) and its activation is important for differentiation and activation of osteoclasts, which are necessary for bone resorption and remodelling (Nakagawa et al., 1998; Takahashi et al., 1999). Signalling through RANK leads to activation of the transcription factor NF $\kappa$ B and the protein kinase JNK. Activation of these factors seems to be important for osteoclastogenesis (formation of mature osteoclasts from their precursor stem cells). Understanding the structure/function relationship for the RANK molecule and residues involved in binding RANKL or adaptor protein during signalling will be valuable for designing future drugs that may relieve diseases associated with perturbations of bone density, such as osteopetrosis or osteoporosis.

Generating TNF mutants with decreased systematic cytotoxicity and increased tumour-specific cytotoxicity is also a useful approach in terms of increasing the overall TNF potency against aggressive tumours. Similarly, TNF inhibitor molecules aimed at sequestering "surplus" TNF molecules in the circulation is a useful approach to reduce any harmful side effects of this cytokine. Of particular interest is a chimeric inhibitor molecule engineered such that the ligand binding domain of the TNFR is linked to parts of an antibody molecule, serving as a form of "artificial" decoy receptor (Peppel et al., 1991). Such chimeric inhibitors have now been produced for clinical use in diseases such as rheumatoid arthritis (Bazzoni and Beutler, 1996; Beutler, 1999) and can be delivered using adenovirus vector (Kolls et al., 1994). Chimeric TNFR-IgG molecules (Fig. 6) may prove useful in a range of other diseases. Continued emergence of the crystal structure of TNF ligands and relevant receptors will be invaluable for efficient design of such chimeras.

### ACKNOWLEDGMENTS

We thank Drs. Bruce Beutler, Steve Sprang, and Byung-Ha Oh for permission to use the TNF figures and for reprints. We also thank Drs. Emad Alnemri, Dirk Anderson, Avi Ashkenazi, Paul Crowe, Abraham de Vos, Yvonne Jones, James G. Pan, and Gavin Screaton for communicating information about their work and Drs. Bryant Darnay and Bharat Aggarwal for useful discussion. We apologise for the authors of many relevant original papers we were unable to cite, due to restrictions on allocated space.

### REFERENCES

- Acton RD, Dahlberg PS, Uknis ME, Klaerner HG, Fink GS, Norman JG, Dunn DL. 1996. Differential sensitivity to *Escherichia coli* infection in mice lacking tumor necrosis factor p55 or interleukin-1 p80 receptors. *Arch Surg* 131:1216–1221.
- Aggarwal B, Vilcek J, Eds. 1991. Tumor necrosis factors: structure, function and mechanism. Marcel Dekker Publishers, New York.
- Anderson DM, Maraskovsky E, Billingsley WL, Dougall WC, Tometsko ME, Roux ER, Teepe MC, DuBose RF, Cosman D, Galibert L. 1997. A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature* 390:175–179.
- Armitage RJ. 1994. Tumor necrosis factor receptor superfamily members and their ligands. *Curr Opin Immunol* 6:407–413.
- Ashkenazi A, Dixit VM. 1998. Death receptors: signaling and modulation. *Science* 281:1305–1308.
- Ashkenazi A, Dixit VM. 1999. Apoptosis control by death and decoy receptors. *Curr Opin Cell Biol* 11:255–260.
- Baeyens KJ, De Bondt HL, Raeymaekers A, Fiers W, De Ranter CJ. 1999. The structure of mouse tumour-necrosis factor at 1.4 Å resolution: towards modulation of its selectivity and trimerization. *Acta Crystallogr D Biol Crystallogr* 55:772–778.
- Baker SJ, Reddy EP. 1996. Transducers of life and death: TNF receptor superfamily and associated proteins. *Oncogene* 12:1–9.
- Baker SJ, Reddy EP. 1998. Modulation of life and death by the TNF receptor superfamily. *Oncogene* 17:3261–3270.
- Baldwin RL, Stolowitz ML, Hood L, Wisniewski BJ. 1996. Structural changes of tumor necrosis factor alpha associated with membrane insertion and channel formation. *Proc Natl Acad Sci U S A* 93:1021–1026.
- Balkwill FR. 1992. Tumour necrosis factor and cancer. *Prog Growth Factor Res* 4:121–137.
- Banner DW, D'Arcy A, Janes W, Gentz R, Schoenfeld HJ, Broger C, Loetscher H, Lesslauer W. 1993. Crystal structure of the soluble human 55 kd TNF receptor-human TNF beta complex: implications for TNF receptor activation. *Cell* 73:431–445.
- Bazan JF. 1993. Emerging families of cytokines and receptors. *Curr Biol* 3:603–606.
- Bazzoni F, Beutler B. 1996. The tumor necrosis factor ligand and receptor families. *N Engl J Med* 334:1717–1725.
- Beutler B, Bazzoni F. 1998. TNF, apoptosis and autoimmunity: a common thread? *Blood Cells Mol Dis* 24:216–230.
- Beutler BA. 1999. The role of tumor necrosis factor in health and disease. *J Rheumatol* 26(Suppl) 57:16–21.
- Beyaert R, Fiers W. 1994. Molecular mechanisms of tumor necrosis factor-induced cytotoxicity. What we do understand and what we do not. *FEBS Lett* 340:9–16.
- Boldin MP, Mett IL, Varfolomeev EE, Chumakov I, Shemer-Avni Y, Camonis JH, Wallach D. 1995. Self association of the "death domains" of the p55 tumor necrosis factor (TNF) receptor and FAS/APO1 prompts signalling for TNF and FAS/APO1 effects. *J Biol Chem* 270:387–391.
- Brojatsch J, Naughton J, Rolls MM, Ziegler K, Young JA. 1996. CAR1, a TNFR-related protein, is a cellular receptor for cytopathic avian leukosis-sarcoma viruses and mediates apoptosis. *Cell* 87:845–855.
- Browning JL, Ngam-ek A, Lawton P, DeMarinis J, Tizard R, Chow EP, Hession C, O'Brine-Greco B, Foley SF, Ware CF. 1993. Lymphotoxin beta, a novel member of the TNF family that forms a heteromeric complex with lymphotoxin on the cell surface. *Cell* 72:847–856.
- Camerini D, Walz G, Loenen WA, Borst J, Seed B. 1991. The T cell activation antigen CD27 is a member of the nerve growth factor/tumor necrosis factor receptor gene family. *J Immunol* 147:3165–3169.
- Carswell EA, Old LJ, Kassel RL, Green S, Fiore N, Williamson B. 1975. An endotoxin-induced serum factor that causes necrosis of tumors. *Proc Natl Acad Sci U S A* 72:3666–3670.
- Cha SS, Kim JS, Cho HS, Shin NK, Jeong W, Shin HC, Kim YJ, Hahn JH, Oh BH. 1998. High resolution crystal structure of a human tumor necrosis factor-alpha mutant with low systemic toxicity. *J Biol Chem* 273:2153–2160.
- Cha SS, Kim MS, Choi YH, Sung BJ, Shin NK, Shin HC, Sung YC, Oh BH. 1999. 2.8 Å resolution crystal structure of human TRAIL, a cytokine with selective antitumor activity. *Immunity* 11:253–261.
- Coley WB. 1893. The treatment of malignant tumors by repeated inoculations of erysipelas: with a report of ten original cases. *Am J Med Sci* 105:487–511.
- Crowe PD, VanArsdale TL, Walter BN, Ware CF, Hession C, Ehrenfels B, Browning JL, Din WS, Goodwin RG, Smith CA. 1994. A



- lymphotoxin-beta-specific receptor [see comments]. *Science* 264: 707-710.
- Czarniecki CW. 1993. The role of tumor necrosis factor in viral disease. *Antiviral Res* 22:223-258.
- D'Arcy A, Banner DW, Janes W, Winkler FK, Loetscher H, Schonfeld HJ, Zulauf M, Gentz R, Lesslauer W. 1993. Crystallization and preliminary crystallographic analysis of a TNF-beta-55 kDa TNF receptor complex. *J Mol Biol* 229:555-557.
- Darnay BG, Aggarwal BB. 1999. Signal transduction by tumor necrosis factor and tumour necrosis factor related ligands and their receptors. *Ann Rheum Dis* 58:S0-S11.
- Degli-Esposti MA, Dougall WC, Smolak PJ, Waugh JY, Smith CA, Goodwin RG. 1997. The novel receptor TRAIL-R4 induces NF-kappaB and protects against TRAIL-mediated apoptosis, yet retains an incomplete death domain. *Immunity* 7:813-820.
- Dembic Z, Loetscher H, Gubler U, Pan YC, Lahm HW, Gentz R, Brockhaus M, Lesslauer W. 1990. Two human TNF receptors have similar extracellular, but distinct intracellular, domain sequences. *Cytokine* 2:231-237.
- Durkop H, Latza U, Hummel M, Eitelbach F, Seed B, Stein H. 1992. Molecular cloning and expression of a new member of the nerve growth factor receptor family that is characteristic for Hodgkin's disease. *Cell* 68:421-427.
- Eck MJ, Sprang SR. 1989. The structure of tumor necrosis factor-alpha at 2.6 Å resolution. Implications for receptor binding. *J Biol Chem* 264:17595-17605.
- Eck MJ, Ultsch M, Rinderknecht E, de Vos AM, Sprang SR. 1992. The structure of human lymphotoxin (tumor necrosis factor-beta) at 1.9-Å resolution. *J Biol Chem* 267:2119-2122.
- Engelmann H, Novick D, Wallach D. 1990. Two tumor necrosis factor-binding proteins purified from human urine. Evidence for immunological cross-reactivity with cell surface tumor necrosis factor receptors. *J Biol Chem* 265:1531-1536.
- Fiers W. 1991. Tumor necrosis factor. Characterization at the molecular, cellular and in vivo level. *FEBS Lett* 285:199-212.
- Frank S, Kohler U, Schackert G, Schackert HK. 1999. Expression of TRAIL and its receptors in human brain tumors [published erratum appears in *Biochem Biophys Res Commun* 1999 May 27;259(1):230]. *Biochem Biophys Res Commun* 257:454-459.
- Goldfeld AE, Tsai EY. 1996. TNF-alpha and genetic susceptibility to parasitic disease. *Exp Parasitol* 84:300-303.
- Goldstein P. 1997. Cell death: TRAIL and its receptors. *Curr Biol* 7:R750-3.
- Granger GA, Williams TW. 1968. Lymphocyte cytotoxicity in vitro: activation and release of a cytotoxic factor. *Nature* 218:1253-1254.
- Gray PW, Aggarwal BB, Benton CV, Bringman TS, Henzel WJ, Jarrett JA, Leung DW, Moffat B, Ng P, Svedersky LP, et al. 1984. Cloning and expression of cDNA for human lymphotoxin, a lymphokine with tumour necrosis activity. *Nature* 312:721-724.
- Grell M, Zimmermann G, Gottfried E, Chen CM, Grunwald U, Huang DC, Wu Lee YH, Durkop H, Engelmann H, Scheurich P, Wajant H, Strasser A. 1999. Induction of cell death by tumour necrosis factor (TNF) receptor 2, CD40 and CD30: a role for TNF-R1 activation by endogenous membrane-anchored TNF. *Embo J* 18:3034-3043.
- Gruss HJ, Dower SK. 1995. Tumor necrosis factor ligand superfamily: involvement in the pathology of malignant lymphomas. *Blood* 85: 3378-3404.
- Hieber U, Heim ME. 1994. Tumor necrosis factor for the treatment of malignancies. *Oncology* 51:142-153.
- Hohmann HP, Remy R, Brockhaus M, van Loon AP. 1989. Two different cell types have different major receptors for human tumor necrosis factor (TNF alpha). *J Biol Chem* 264:14927-14934.
- Hu FQ, Smith CA, Pickup DJ. 1994. Cowpox virus contains two copies of an early gene encoding a soluble secreted form of the type II TNF receptor. *Virology* 204:343-356.
- Hymowitz SG, Christinger HW, Fuh G, Ultsch M, O'Connell M, Kelley RF, Ashkenazi A, de Vos AM. 1999. Triggering cell death: the crystal structure of Apo2L/TRAIL in a complex with death receptor 5. *Mol Cell* 4:563-571.
- Hymowitz SG, Christinger HW, Fuh G, Ultsch M, O'Connell M, Kelley RF, Ashkenazi A, de Vos AM. 2000. A unique zinc-binding site revealed by a high resolution X-ray structure of homotrimeric Apo2L/TRAIL. *Biochemistry* 39:633-640.
- Idriss H. 2000. Do TNF  $\alpha$  insensitive cancer cells escape  $\alpha$ -tubulin nitro-tyrosinylation? *Nitric Oxide* 4:1-3.
- Itoh N, Yonehara S, Ishii A, Yonehara M, Mizushima S, Sameshima M, Hase A, Seto Y, Nagata S. 1991. The polypeptide encoded by the cDNA for human cell surface antigen Fas can mediate apoptosis. *Cell* 66:233-243.
- Jones EY, Stuart DI, Walker NP. 1989. Structure of tumour necrosis factor. *Nature* 338:225-228.
- Jones EY, Stuart DI, Walker NPC. 1991. Crystal Structure of TNF. In: Aggarwal BB, Vilcek J, editors. *Tumor necrosis factors: structure, function and mechanism*, Marcel Dekker, Publishers, New York.
- Kagan BL, Baldwin RL, Munoz D, Wisniewski BJ. 1992. Formation of ion-permeable channels by tumor necrosis factor-alpha. *Science* 255:1427-1430.
- Karpusas M, Hsu YM, Wang JH, Thompson J, Lederman S, Chess L, Thomas D. 1995. 2 Å crystal structure of an extracellular fragment of human CD40 ligand [erratum appears in *Structure* 1995 Dec 15;3(12):1426]. *Structure* 3:1031-1039.
- Kolls J, Peppel K, Silva M, Beutler B. 1994. Prolonged and effective blockade of tumor necrosis factor activity through adenovirus-mediated gene transfer [published erratum appears in *Proc Natl Acad Sci USA* 1994 Mar 1; 91(5):1979. *Proc Natl Acad Sci USA* 91:215-219.
- Krueger JM, Fang J, Taishi P, Chen Z, Kushikata T, Gardi J. 1998. Sleep. A physiologic role for IL-1 beta and TNF-alpha. *Ann N Y Acad Sci* 856:148-159.
- Lang F, Lepple-Wienhues A, Paulmichl M, Szabo I, Siemen D, Gulbins E. 1998. Ion channels, cell volume, and apoptotic cell death. *Cell Physiol Biochem* 8:285-292.
- Loetscher H, Pan YC, Lahm HW, Gentz R, Brockhaus M, Tabuchi H, Lesslauer W. 1990. Molecular cloning and expression of the human 55 kd tumor necrosis factor receptor. *Cell* 61:351-359.
- Loetscher H, Gentz R, Zulauf M, Lustig A, Tabuchi H, Schlaeger EJ, Brockhaus M, Gallati H, Manneberg M, Lesslauer W. 1991a. Recombinant 55-kDa tumor necrosis factor (TNF) receptor. Stoichiometry of binding to TNF alpha and TNF beta and inhibition of TNF activity. *J Biol Chem* 266:18324-18329.
- Loetscher H, Steinmetz M, Lesslauer W. 1991b. Tumor necrosis factor: receptors and inhibitors. *Cancer Cells* 3:221-226.
- MacFarlane M, Ahmad M, Srinivasula SM, Fernandes-Alnemri T, Cohen GM, Alnemri ES. 1997. Identification and molecular cloning of two novel receptors for the cytotoxic ligand TRAIL. *J Biol Chem* 272:25417-25420.
- Mallett S, Fossum S, Barclay AN. 1990. Characterization of the MRC OX40 antigen of activated CD4 positive T lymphocytes: a molecule related to nerve growth factor receptor. *Embo J* 9:1063-1068.
- Marsters SA, Ayres TM, Skubatch M, Gray CL, Rothe M, Ashkenazi A. 1997. Herpesvirus entry mediator, a member of the tumor necrosis factor receptor (TNFR) family, interacts with members of the TNFR-associated factor family and activates the transcription factors NF-kappaB and AP-1. *J Biol Chem* 272:14029-14032.
- McWhirter SM, Pullen SS, Holton JM, Crute JJ, Kehry MR, Alber T. 1999. Crystallographic analysis of CD40 recognition and signaling by human TRAF2. *Proc Natl Acad Sci U S A* 96:8408-8413.
- Mongkolsapaya J, Grimes JM, Chen N, Xu XN, Stuart DI, Jones EY, Screaton GR. 1999. Structure of the TRAIL-DR5 complex reveals mechanisms conferring specificity in apoptotic initiation. *Nat Struct Biol* 6:1048-1053.
- Montgomery RI, Warner MS, Lum BJ, Spear PG. 1996. Herpes simplex virus-1 entry into cells mediated by a novel member of the TNF/NGF receptor family. *Cell* 87:427-436.
- Naismith JH, Sprang SR. 1998. Modularity in the TNF-receptor family. *Trends Biochem Sci* 23:74-79.
- Naismith JH, Devine TQ, Brandhuber BJ, Sprang SR. 1995. Crystallographic evidence for dimerization of unliganded tumor necrosis factor receptor. *J Biol Chem* 270:13303-13307.
- Naismith JH, Devine TQ, Kohno T, Sprang SR. 1996. Structures of the extracellular domain of the type I tumor necrosis factor receptor. *Structure* 4:1251-1262.
- Nakagawa N, Kinoshita M, Yamaguchi K, Shima N, Yasuda H, Yano K, Morinaga T, Higashio K. 1998. RANK is the essential signaling receptor for osteoclast differentiation factor in osteoclastogenesis. *Biochem Biophys Res Commun* 253:395-400.
- Nocentini G, Giunchi L, Ronchetti S, Krausz LT, Bartoli A, Moraca R, Migliorati G, Riccardi C. 1997. A new member of the tumor necrosis factor/nerve growth factor receptor family inhibits T cell receptor-induced apoptosis. *Proc Natl Acad Sci U S A* 94:6216-6221.
- Orlinick JR, Chao MV. 1998. TNF-related ligands and their receptors. *Cell Signal* 10:543-551.
- Pan G, Ni J, Wei YF, Yu G, Gentz R, Dixit VM. 1997a. An antagonist decoy receptor and a death domain-containing receptor for TRAIL [see comments]. *Science* 277:815-818.
- Pan G, O'Rourke K, Chinnaiyan AM, Gentz R, Ebner R, Ni J, Dixit VM. 1997b. The receptor for the cytotoxic ligand TRAIL. *Science* 276:111-113.
- Pan G, Bauer JH, Haridas V, Wang S, Liu D, Yu G, Vincenz C, Aggarwal BB, Ni J, Dixit VM. 1998a. Identification and functional



- characterization of DR6, a novel death domain-containing TNF receptor. *FEBS Lett* 431:351–356.
- Pan G, Ni J, Yu G, Wei YF, Dixit VM. 1998b. TRUNDD, a new member of the TRAIL receptor family that antagonizes TRAIL signalling. *FEBS Lett* 424:41–45.
- Pearson RB, Kemp BE. 1991. Protein kinase phosphorylation site sequences and consensus specificity motifs: tabulations. *Methods Enzymol* 200:62–81.
- Pennica D, Nedwin GE, Hayflick JS, Seeburg PH, Derynck R, Palladino MA, Kohr WJ, Aggarwal BB, Goeddel DV. 1984. Human tumour necrosis factor: precursor structure, expression and homology to lymphotoxin. *Nature* 312:724–729.
- Peppl K, Crawford D, Beutler B. 1991. A tumor necrosis factor (TNF) receptor-IgG heavy chain chimeric protein as a bivalent antagonist of TNF activity. *J Exp Med* 174:1483–1489.
- Pitti RM, Marsters SA, Lawrence DA, Roy M, Kischkel FC, Dowd P, Huang A, Donahue CJ, Sherwood SW, Baldwin DT, Godowski PJ, Wood WI, Gurney AL, Hillan KJ, Cohen RL, Goddard AD, Botstein D, Ashkenazi A. 1998. Genomic amplification of a decoy receptor for Fas ligand in lung and colon cancer. *Nature* 396:699–703.
- Pocsik E, Duda E, Wallach D. 1995. Phosphorylation of the 26 kDa TNF precursor in monocytic cells and in transfected HeLa cells. *J Inflamm* 45:152–160.
- Pullen SS, Labadia ME, Ingraham RH, McWhirter SM, Everdeen DS, Alber T, Crute JJ, Kehry MR. 1999. High-affinity interactions of tumor necrosis factor receptor-associated factors (TRAFs) and CD40 require TRAF trimerization and CD40 multimerization [In Process Citation]. *Biochemistry* 38:10168–10177.
- Radeke HH, Meier B, Topley N, Flöge J, Habermehl GG, Resch K. 1990. Interleukin 1- $\alpha$  and tumor necrosis factor- $\alpha$  induce oxygen radical production in mesangial cells. *Kidney Int* 37:767–775.
- Reed C, Fu ZQ, Wu J, Xue YN, Harrison RW, Chen MJ, Weber IT. 1997. Crystal structure of TNF- $\alpha$  mutant R31D with greater affinity for receptor R1 compared with R2. *Protein Eng* 10:1101–1107.
- Remy I, Wilson IA, Michnick SW. 1999. Erythropoietin receptor activation by a ligand-induced conformation change. *Science* 283:990–993.
- Ryffel B, Mihatsch MJ. 1993. TNF receptor distribution in human tissues. *Int Rev Exp Pathol* 34 Pt B:149–156.
- Schall TJ, Lewis M, Koller KJ, Lee A, Rice GC, Wong GH, Gatanaga T, Granger GA, Lentz R, Raab H, et al. 1990. Molecular cloning and expression of a receptor for human tumor necrosis factor. *Cell* 61:361–370.
- Schoenfeld HJ, Poeschl B, Frey JR, Loetscher H, Hunziker W, Lustig A, Zulauf M. 1991. Efficient purification of recombinant human tumor necrosis factor beta from *Escherichia coli* yields biologically active protein with a trimeric structure that binds to both tumor necrosis factor receptors. *J Biol Chem* 266:3863–3869.
- Screaton GR, Xu XN, Olsen AL, Cowper AE, Tan R, McMichael AJ, Bell JI. 1997. LARD: a new lymphoid-specific death domain containing receptor regulated by alternative pre-mRNA splicing. *Proc Natl Acad Sci U S A* 94:4615–4619.
- Shakhov AN, Collart MA, Vassalli P, Nedospasov SA, Jongeneel CV. 1990. Kappa B-type enhancers are involved in lipopolysaccharide-mediated transcriptional activation of the tumor necrosis factor alpha gene in primary macrophages. *J Exp Med* 171:35–47.
- Shalaby MR, Sundan A, Loetscher H, Brockhaus M, Lesslauer W, Espevik T. 1990. Binding and regulation of cellular functions by monoclonal antibodies against human tumor necrosis factor receptors. *J Exp Med* 172:1517–1520.
- Shoham S, Davenne D, Cady AB, Dinarello CA, Krueger JM. 1987. Recombinant tumor necrosis factor and interleukin 1 enhance slow-wave sleep. *Am J Physiol* 253:R142–149.
- Simonet WS, Lacey DL, Dunstan CR, Kelley M, Chang MS, Luthy R, Nguyen HQ, Wooden S, Bennett L, Boone T, Shimamoto G, DeRose M, Elliott R, Colombero A, Tan HL, Trail G, Sullivan J, Davy E, Bucay N, Renshaw-Gegg L, Hughes TM, Hill D, Pattison W, Campbell P, Boyle WJ, et al. 1997. Osteoprotegerin: a novel secreted protein involved in the regulation of bone density [see comments]. *Cell* 89:309–319.
- Singh J, Garber E, Van Vlijmen H, Karpusas M, Hsu YM, Zheng Z, Naismith JH, Thomas D. 1998. The role of polar interactions in the molecular recognition of CD40L with its receptor CD40. *Protein Sci* 7:1124–1135.
- Smith CA, Davis T, Anderson D, Solam L, Beckmann MP, Jerzy R, Dower SK, Cosman D, Goodwin RG. 1990. A receptor for tumor necrosis factor defines an unusual family of cellular and viral proteins. *Science* 248:1019–1023.
- Smith CA, Hu FQ, Smith TD, Richards CL, Smolak P, Goodwin RG, Pickup DJ. 1996. Cowpox virus genome encodes a second soluble homologue of cellular TNF receptors, distinct from CrmB, that binds TNF but not LT  $\alpha$ . *Virology* 223:132–147.
- Smith RA, Baglioni C. 1987. The active form of tumor necrosis factor is a trimer. *J Biol Chem* 262:6951–6954.
- Smith RA, Baglioni C. 1992. Characterization of TNF receptors. *Immunol Ser* 56:131–147.
- Sprang SR, Eck MJ. 1991. The structure of tumor necrosis factor: perspectives on biological activity. In: Beutler B, editor. *The tumor necrosis factors. The molecules in their emerging role in medicine*. New York: Raven Press.
- Spriggs DR, Deutsch S, Kufe DW. 1992. Genomic structure, induction, and production of TNF- $\alpha$ . *Immunol Ser* 56:3–34.
- Steinshamn S, Bemelmans MH, van Tits LJ, Bergh K, Buurman WA, Waage A. 1996. TNF receptors in murine *Candida albicans* infection: evidence for an important role of TNF receptor p55 in antifungal defense. *J Immunol* 157:2155–2159.
- Steller H. 1995. Mechanisms and genes of cellular suicide. *Science* 267:1445–1449.
- Takahashi N, Udagawa N, Suda T. 1999. A new member of tumor necrosis factor ligand family, ODF/OPGL/TRANCE/RANKL, regulates osteoclast differentiation and function. *Biochem Biophys Res Commun* 256:449–455.
- Thoma B, Grell M, Pfizenmaier K, Scheurich P. 1990. Identification of a 60-kD tumor necrosis factor (TNF) receptor as the major signal transducing component in TNF responses. *J Exp Med* 172:1019–1023.
- Thornberry NA, Lazebnik Y. 1998. Caspases: enemies within. *Science* 281:1312–1316.
- Tsuda E, Goto M, Mochizuki S, Yano K, Kobayashi F, Morinaga T, Higashio K. 1997. Isolation of a novel cytokine from human fibroblasts that specifically inhibits osteoclastogenesis. *Biochem Biophys Res Commun* 234:137–142.
- Van Arsdale TL, Van Arsdale SL, Force WR, Walter BN, Mosialos G, Kieff E, Reed JC, Ware CF. 1997. Lymphotoxin-beta receptor signaling complex: role of tumor necrosis factor receptor-associated factor 3 recruitment in cell death and activation of nuclear factor kappaB. *Proc Natl Acad Sci U S A* 94:2460–2465.
- Van Linden AA, Cottin V, Leu C, Riches DW. 2000. Phosphorylation of the membrane proximal region of tumor necrosis factor receptor CD120a (p55) at ERK consensus sites. *J Biol Chem* 275:6996–7003.
- Van Ostade X, Vandenabeele P, Everaerd B, Loetscher H, Gentz R, Brockhaus M, Lesslauer W, Tavernier J, Brouckaert P, Fiers W. 1993. Human TNF mutants with selective activity on the p55 receptor [see comments]. *Nature* 361:266–269.
- Vilcek J, Lee TH. 1991. Tumor necrosis factor. New insights into the molecular mechanisms of its multiple actions. *J Biol Chem* 266:7313–7316.
- Vinay DS, Kwon BS. 1999. Differential expression and costimulatory effect of 4-1BB (CD137) and CD28 molecules on cytokine-induced murine CD8(+) Tc1 and Tc2 cells. *Cell Immunol* 192:63–71.
- Walczak H, Degli-Esposti MA, Johnson RS, Smolak PJ, Waugh JY, Boiani N, Timour MS, Gerhart MJ, Schooley KA, Smith CA, Goodwin RG, Rauch CT. 1997. TRAIL-R2: a novel apoptosis-mediating receptor for TRAIL. *Embo J* 16:5386–5397.
- Watts AD, Hunt NH, Hambly BD, Chaudhri G. 1997. Separation of tumor necrosis factor alpha isoforms by two-dimensional polyacrylamide gel electrophoresis. *Electrophoresis* 18:1086–1091.
- Watts AD, Hunt NH, Wanigasekara Y, Bloomfield G, Wallach D, Roufogalis BD, Chaudhri G. 1999. A casein kinase I motif present in the cytoplasmic domain of members of the tumour necrosis factor ligand family is implicated in 'reverse signalling'. *Embo J* 18:2119–2126.
- Wride MA, Sanders EJ. 1995. Potential roles for tumour necrosis factor alpha during embryonic development. *Anat Embryol (Berl)* 191:1–10.
- Yuan J. 1997. Transducing signals of life and death. *Curr Opin Cell Biol* 9:247–251.