

Regression Analysis Of Population Drinking Dataset

Sayan Mukherjee

Contents

1	Introduction	2
2	Exploratory Data Analysis	2
3	Regression Analysis	6
3.1	Fitting a Linear Model	6
3.2	Checking Model Assumptions	7
3.3	Detecting Influential Points	12
3.4	Remedies For Influential Points	16
3.5	Collinearity	18
3.6	Remedies For Collinearity	20
3.7	Model Selection	21
3.8	Shrinkage Methods	31
3.9	Robust Regression Methods	33

1 Introduction

About The Data

Here we have 46 observations from different places of the following quantiles :-

Urban Population
Late Births
Wine Consumption Per Capita
Liquor Consumption Per Capita
Cirrhosis Death Rate

Here our response variable is Cirrhosis Death Rate and others are all covariates. To get an overview of the data, we first perform the exploratory data analysis.

2 Exploratory Data Analysis

Loading the Dataset

To get an overview of the data, we first load it in R and print first few values:-

```
library(MASS)
library(lattice)
library(olsrr)
library(car)
library(Lipack)
X = read.csv(file = "D:\\PG files\\Projects\\Regression-Analysis-Project-main\\population_drinking1.txt",
header=TRUE,sep = "\\t")
names(X) <- c("Ind","Ind_1","Urban population", "Late births", "Wine consumption per capita", "Liquor consumption per cap", "Cirrhosis death rate")
head(X)
```

	Ind	Ind_1	Urban population	Late births	Wine consumption per capita	
1	1	1	44	33.2		5
2	2	1	43	33.8		4
3	3	1	48	40.6		3
4	4	1	52	39.2		7
5	5	1	71	45.5		11
6	6	1	44	37.5		9

	Liquor consumption per capita	Cirrhosis death rate
1	30	41.2
2	41	31.7
3	38	39.4
4	48	57.5
5	53	74.8
6	65	59.8

Here we have “Cirrhosis death rate” as the response and “Urban population (A1)”, “Late births(A2)”, “Wine consumption per capita(A3)”, “Liquor consumption per capita(A4)” as the covariates.

Type of the covariates

To know type of each covariates, we use the str() function :-

```
str(X)
```

```
'data.frame': 46 obs. of 7 variables:
 $ Ind      : int  1 2 3 4 5 6 7 8 9 10 ...
 $ Ind_1    : int  1 1 1 1 1 1 1 1 1 1 ...
 $ Urban population : int  44 43 48 52 71 44 57 34 70 54 ...
 $ Late births : num  33.2 33.8 40.6 39.2 45.5 37.5 44.2 31.9 45.6 45.9 ...
 $ Wine consumption per capita : int  5 4 3 7 11 9 6 3 12 7 ...
 $ Liquor consumption per capita: int  30 41 38 48 53 65 73 32 56 57 ...
 $ Cirrhosis death rate : num  41.2 31.7 39.4 57.5 74.8 59.8 54.3 47.9 77.2 56.6 ...
```

So the dataset contains no factor covariate hence we can perform multiple linear regression here. For ease of indexing, we name the columns as “I”, “1”, “A1”, “A2”, “A3”, “A4”, “Y”.

```
names(X) <- c("I", "1", "A1", "A2", "A3", "A4", "Y")
```

5-number Summary of Covariates

To get an idea of the values of each covariate we calculate the 5-number summary for each of them :-

```
summary(X[, -c(1,2)])
```


A1		A2		A3		A4	
Min.	:27.00	Min.	:31.20	Min.	: 2.00	Min.	: 26.00
1st Qu.	:44.25	1st Qu.	:35.62	1st Qu.	: 6.25	1st Qu.	: 41.50
Median	:55.00	Median	:42.25	Median	:10.00	Median	: 56.00
Mean	:56.26	Mean	:41.48	Mean	:11.59	Mean	: 57.50
3rd Qu.	:65.00	3rd Qu.	:45.83	3rd Qu.	:15.75	3rd Qu.	: 68.75
Max.	:87.00	Max.	:56.10	Max.	:31.00	Max.	:149.00

Y	
Min.	: 28.00
1st Qu.	: 48.90
Median	: 57.65
Mean	: 63.49
3rd Qu.	: 75.70
Max.	:129.90

Pairwise Scatterplots

To get an idea of the relationship between covariates and response, we make pairwise scatterplots using pairs() function :-

```
pairs(X[, -c(1,2)])
```



This plot clearly indicates linear relationship between the covariates and response also. This might lead to the problem of multicollinearity which we will formally diagnose.

Correlation Between Covariates

We calculate the correlation between the covariates and response to get even better idea of linear dependence between them :-


	A1	A2	A3	A4
A1	1.0000000	0.8432812	0.6786230	0.4402957
A2	0.8432812	1.0000000	0.6398407	0.6863643
A3	0.6786230	0.6398407	1.0000000	0.6759206
A4	0.4402957	0.6863643	0.6759206	1.0000000

We can see that the correlations are high between many predictors which can lead to problem of multicollinearity.

Partial Residual Plots

To get an idea of the nature of relationship between the covariates and response, we make the partial residual plot for all the covariates :-

```
crPlots(lm(Y~A1+A2+A3+A4,data = X))
```



Conclusion

The plot indicates the linear relationship between the covariates and response.


Hence, we will fit the usual multiple linear regression model with no higher order polynomial terms.

Later we will see if other models with interaction terms are better or not.

Boxplots of Covariates and Response

We draw the boxplots for different covariates to get idea of presence of outlier / high leverage points :-

```
boxplot(X[, -c(1,2)])
```



Here also we get some indication of possible presence of those points.

3 Regression Analysis

3.1 Fitting a Linear Model

Fitting a Linear Model to the Dataset

We fit a linear model of the form :-

$$\mathbf{Y}^{n \times 1} = \mathbf{X}^{n \times p} \boldsymbol{\beta}^{p \times 1} + \boldsymbol{\epsilon}^{n \times 1}$$

where $n = 46$ (total number of observed responses) and $p = 5$ where columns of $\mathbf{X} = [\mathbf{1}_n \quad \mathbf{x}_1 \quad \mathbf{x}_2 \quad \mathbf{x}_3 \quad \mathbf{x}_4]$ and $\boldsymbol{\beta} = (\beta_0 \quad \cdots \quad \beta_4)$ each corresponding to the 4 different covariates.

We fit a linear model based on the given dataset in R and then verify the different assumptions of it.

Features of the fitted model

We fit the linear model specified before in the dataset using `lm()` function and to get an idea about the estimates we use the `summary()` function :-

```
colnames(X)=c("I","1","A1","A2","A3","A4","Y")
attach(X)
reg <- lm(Y~A1+A2+A3+A4)
summary(reg)
```

Call:
lm(formula = Y ~ A1 + A2 + A3 + A4)

Residuals:

	Min	1Q	Median	3Q	Max
	-18.8723	-6.7803	0.1507	7.3252	16.4419

Coefficients:

	Estimate	Std. Error	t value	Pr(> t)
(Intercept)	-13.96310	11.40035	-1.225	0.2276
A1	0.09829	0.24407	0.403	0.6893
A2	1.14838	0.58300	1.970	0.0556 .
A3	1.85786	0.40096	4.634	3.61e-05 ***
A4	0.04817	0.13336	0.361	0.7198

Signif. codes: 0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 10.61 on 41 degrees of freedom
Multiple R-squared: 0.8136, Adjusted R-squared: 0.7954
F-statistic: 44.75 on 4 and 41 DF, p-value: 1.951e-14

Explanation of the fitted model

As we can see only the coefficients β_2, β_3 for covariates “A2”, “A3” are statistically significant.


This does not imply that other covariates are insignificant since there maybe many problems that are hidden in the model.

So, before concluding anything we verify all the assumptions of a linear model.

Obs vs Fitted Values

We plot the observed vs fitted values to get some idea about prediction :-

```
plot(1:nrow(X),X$Y,type = "o",pch = 20,ylab = "observed & predicted values",xlab = "Index")
lines(1:nrow(X),reg$fitted.values,type = "o",pch = 20,col = "red",lty = 2)
abline(v = 1:nrow(X),lty = 2,col = rgb(0,1,0,alpha = 0.3))
legend("topleft",legend = c("Obs","Pred"),fill = c("black","red"))
```



The fit is good except a few observations.


There may be many reasons for this which we will eventually look into.

3.2 Checking Model Assumptions

QQ-plot of residuals

We now plot the sorted residuals (quantiles) against the population quantiles of a normal distribution :-

```
resi<-residuals(reg)
qqnorm(resi,ylab="Residuals",main="")
qqline(resi)
```



We can see the qq-plot indicates light tailed residuals with possible deviation from normality.

There may be some outlier points present which we will verify later.

Shapiro-Wilk Test

We test the following hypothesis H_0 : residuals are normally distributed against H_1 : H_0 is false using Shapiro-Wilk test in R as :-

```
shapiro.test(resi)

Shapiro-Wilk normality test


data:  resi
W = 0.95987, p-value = 0.1133
```

Though the p-value is more than 0.1 but this doesn't give strong evidence in favour of H_0 so we will further check for presence of correlation between the errors and other issues also.

Checking Homoskedasticity Assumptions

First to check homoskedasticity assumption, we make the residuals ($\hat{\epsilon}$) vs fitted (\hat{y}) plots :-

```
plot(fitted(reg), residuals(reg), xlab="Fitted", ylab="Residuals")
abline(h=0)
```

We can see the plot doesn't give indication of presence of heteroskedasticity, hence we will perform confirmatory tests.


[https://online.stat.psu.edu/stat462/node/117/#:~:text=When%20conducting%20a%20residual%](https://online.stat.psu.edu/stat462/node/117/#:~:text=When%20conducting%20a%20residual%20analysis%20of%20variance%20ANOVA,when%20the%20variance%20of%20the%20residuals%20is%20not%20constant,when%20the%20variance%20of%20the%20residuals%20is%20not%20constant)

b_i vs \hat{y}_i plot

A standard technique to detect presence of heteroskedasticity is to plot the quantities $b_i = \frac{e_i^2}{1-h_i}$ against the fitted values \hat{y}_i .

We make the plot using R :-

```
A = as.matrix(X[,-1])
H = A%*%solve(t(A)%*%A)%*%t(A)
H_i = diag(H)
e_i = residuals(reg)
b_i = e_i^2/(1-H_i)
plot(fitted(reg),b_i,pch = 20,main = bquote("Plot of" ~ b[i]^2 ~ "vs fitted values"))
```



This plot gives no indication of any heteroskedasticity present in the residuals.

Breusch-Pagan Test

We perform the Breusch-Pagan test for testing the homoskedasticity assumptions using R :-

```
library(lmtest)
bptest(reg)

studentized Breusch-Pagan test


data:  reg
BP = 2.6929, df = 4, p-value = 0.6105
```

We can see that the p-value of the outcome is satisfactorily high so we can safely assume the error variances to be equal.

ACF plot

If the errors in the model are truly independent, then we will expect the sample autocorrelation coefficients for different lags k to be insignificant.

```
acf(resi,ylab = "",xlab = "",main = "")
title(xlab="Lag", ylab="ACF", line=2)
```




This plot clearly gives indication of no presence of any type of correlation between the residuals.

PACF plot

Similarly, we plot the sample partial autocorrelation coefficients for different lags and got the same kind of observations indicating no presence of correlations.

```
pacf(resi,ylab = "",xlab = "",main = "")
title(xlab="Lag", ylab="PACF", line=2)
```



Durbin-Watson Test

To test the null hypothesis H_0 : errors are uncorrelated against H_1 : errors are correlated, we perform Durbin-Watson test which gives the following results :-

```
require(lmtest)
dwtest(Y~A1+A2+A3+A4,data=X)
```

Durbin-Watson test

data: Y ~ A1 + A2 + A3 + A4

DW = 2.5494, p-value = 0.9734

alternative hypothesis: true autocorrelation is greater than 0

Since the test gives high p-value we can accept H_0 hence the assumption of uncorrelated residuals can be assumed to be satisfied.

Breusch–Godfrey test

To check whether residuals are uncorrelated for higher orders, we perform the Breusch–Godfrey test upto order 20.

```
require(lmtest)
bptest(reg,order = 20)
```

Breusch-Godfrey test for serial correlation of order up to 20

data: reg

LM test = 24.951, df = 20, p-value = 0.2033

Here also the p-value is fairly high favouring the null assumption.

3.3 Detecting Influential Points

Hat Matrix Diagonals

To detect high leverage points, we compute the hat matrix diagonals h_i of the matrix $\mathbf{H} = \mathbf{X}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T$:-

```
hat_d <- hatvalues(reg)
head(sort(hat_d,dec=TRUE))
```

	36	20	12	38	30	39
	0.4994692	0.3042241	0.3015417	0.2998329	0.2078749	0.1721364

We find out if there is any diagonal element with value $> \frac{2p}{n}$ as they should be looked at more closely.

```
n = nrow(X); p = 5
hat_d[hat_d > 2*(p/n)]
```


	12	20	36	38
	0.3015417	0.3042241	0.4994692	0.2998329

Hence we will apply other procedures also to confirm whether these points are influential or not.

Externally Studentized Residuals

We plot the externally studentized residuals using the formula $t_i^2 = r_i^2 \left(\frac{n-p-1}{n-p-r_i^2} \right)$:-


```
stud <- rstudent(reg)
plot(stud,ylim = c(-3,3),pch=20,col = "blue")
abline(h=c(0))
```



If the assumptions are correct i.e. $\epsilon \sim N_n(\mathbf{0}, \sigma^2 \mathbf{I})$ then we should get that $t_i \sim t_{n-p-1}$.

Hence the significant externally studentized residuals will have values $|t_i| > t_{n-p-1; \frac{\alpha}{2}} \iff t_i^2 > F_{n-p-1; \alpha} :-$

```
ols_plot_resid_stud_fit(reg)
```



We can see from the plot that one residual is significant hence we treat that as an outlier.


DFBETAS

After outliers, we check for presence of high leverage points, which can be detected using DFBETAS measure for different parameters $DFBETAS_{ij} = \frac{\hat{\beta}_j - \hat{\beta}^{(i)}_j}{s(i) \sqrt{\sum_i c_{j+1,i}^2}}$

where $\mathbf{C} = ((c_{ij})) = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}$.

We will consider the points for which $|DFBETAS_{ij}| > \frac{2}{\sqrt{n}}$. In the next slide we plot the values for all the 5 coefficients $\beta_i, i = 0, \dots, 4$.


```
par(mfrow = c(2,3))
DFBETAS = dfbetas(reg)
for(i in 1:5)
{
  plot(DFBETAS[,i], main=bquote("DFBETAS for" ~ beta[.(i-1)]), ylab="", ylim=c(-1.5, 1.5), xlab="", pch=
  abline(h=c(-2/sqrt(n), 2/sqrt(n)))
  ind = which(abs(DFBETAS[,i]) > 2/sqrt(n)) # beta_0
  text = text(ind, DFBETAS[ind,i], pos = 3, labels = ind)
}
```



DFFITS

To notice the change in fitted values, we plot the DFFITS values for all the points where $DFFITS_i = t_i \left(\frac{h_i}{1-h_i} \right)^{1/2}$ and we will check for the points for which $|DFFITS_i| > 2\sqrt{\frac{p}{n}}$.

```
ols_plot_dffits(reg)
```




COVRATIO

We also plot the COVRATIO values which are defined as

$$COVRATIO_i = \left(\frac{n-p-1}{n-p} + \frac{t_i^2}{n-p} \right)^{-p} (1-h_i)^{-1}$$

and we consider the points to have high fluence for which $|COVRATIO - 1| > \frac{3p}{n}$.

```
COVRATIO = covratio(reg)
plot(abs(COVRATIO-1),ylab=expression(abs(COVRATIO-1)),pch = 20)
abline(h = 3*p/n)
ind = which(abs(COVRATIO-1) >= 3*p/n)
text(ind,abs(COVRATIO[ind]-1),pos = 1,labels = ind)
```




Cook's D

Lastly, we calculate the Cook's Distance $D_i = \frac{(\hat{\beta}(i) - \hat{\beta})^T \mathbf{X}^T \mathbf{X} (\hat{\beta}(i) - \hat{\beta})}{pS^2} = r_i^2 \frac{h_i}{p(1-h_i)}$ for all the n points.

We flag the points as suspicious for which $D_i > \frac{4}{n}$ here $n = 46, p = 5$. Whose value equals to 0.087.

We plot the values and see if such suspicious points exists or not.

```
COOKSD = cooks.distance(reg)
plot(COOKSD,pch=20)
abline(h = 3*mean(COOKSD))
ind = which(COOKSD > 3*mean(COOKSD)) # beta_0
text = text(ind,COOKSD[ind],pos = 1,labels = ind)
```



We can clearly notice that the points 12,20,30,36 have significant values of D_i . So we will investigate them further.

Conclusion

From all the diagnostics performed for finding influential observations, we can make the following table of our findings :-

Diagnostic Measures	Points Detected
h_i	12, 20, 36, 38
t_i	12
$DFBETAS$	12, 15, 17, 18, 20, 28, 30, 36
$DFFITs$	12, 20, 30, 36
$COVRATIO$	24, 36, 38
$Cook's D$	12, 20, 30, 36

Hence, from the table, we conclude the points 12, 20, 30, 36 to be influential points and we will later remove them from the model and see the changes occurring in all aspects of the fitted linear models.

3.4 Remedies For Influential Points

Removing Influential Points

We remove the influential points and then again fit a linear model with all the covariates and write the summary output of the fitted model here :-

```
summary(lm(Y~A1+A2+A3+A4,data=X[-c(12,20,30,36),]))
```

Call:

```
lm(formula = Y ~ A1 + A2 + A3 + A4, data = X[-c(12, 20, 30, 36),
])
```



```

Residuals:
    Min       1Q   Median       3Q      Max
-16.6468  -5.0683  -0.3998   6.1885  16.7016

Coefficients:
              Estimate Std. Error t value Pr(>|t|)
(Intercept)  20.2021    11.7358   1.721  0.09353 .
A1           0.8783     0.2582   3.401  0.00162 **
A2          -0.7104     0.6205  -1.145  0.25960
A3           1.2890     0.3960   3.255  0.00243 **
A4           0.1489     0.1312   1.134  0.26392
---
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1

Residual standard error: 8.672 on 37 degrees of freedom
Multiple R-squared:  0.8236, Adjusted R-squared:  0.8045
F-statistic: 43.18 on 4 and 37 DF, p-value: 1.867e-13

```

This model has increased value of $R_{adj}^2 = 0.8045$.

Improvements in the fitted model

From the output, we can see that the R_{adj}^2 value has increased from that of the full model which was $= 0.7954$.

Also we can see that the model indicates the estimates of the intercept term $(\hat{\beta}_0)$, variables A1 & A3 $(\hat{\beta}_1, \hat{\beta}_3)$ to be significant.

Whether covariate A2 is significant or not, will be verified later.


Histogram of Residuals

We also plot the histogram of the residual values and can notice it's symmetric about 0 and seems to be normally distributed :-

```

reg_out = lm(Y~A1+A2+A3+A4,data = X[-c(12,20,30,36),])
hist(rstandard(reg_out),breaks = 5,main = "Histogram Of Residual Values")


```



QQ-plot

For checking the assumptions for the residuals we again make the quantile-quantile plot of the standardized residuals :-

```
reg_out = lm(Y~A1+A3+A4,data = X[-c(12,20,30,36),])
qqnorm(rstandard(reg_out))
qqline(rstandard(reg_out))
```



Checking Other Assumptions

We again perform both the Shapiro-Wilk test and Durbin-Watson test for checking normality and presence of correlation between the residuals, respectively. We write down the observations in the following table :-

Tests	Model with influential points	Model without influential points
Shapiro-Wilk	0.1133	0.8429
Durbin-Watson	0.9734	0.6943
Breusch-Pagan	0.6105	0.5742
Breusch-Godfrey	0.2033	0.7973

Hence we can see considerable improvement in the normality assumptions of the residuals whereas the uncorrelated & homoskedasticity assumptions are more or less remains equally acceptable.

Hence, the model can be considered to be better than the previous model as a result of removing the influential points.

3.5 Collinearity

Multicollinearity

Next we consider the problem of multicollinearity that may be present in our dataset as suspected from the pairwise scatterplots.

We calculate the condition number for the scaled and centred model matrix \mathbf{X}^* which is $\kappa(\mathbf{X}^*) = \sqrt{\frac{\lambda_{max}}{\lambda_{min}}}$ where λ_i 's are the eigenvalues of $\mathbf{X}^{*T}\mathbf{X}^* = \mathbf{R}_{xx}$. We calculate $\kappa(\mathbf{X}^*)$ using R :-

```
X_md1 = model.matrix(reg)[-1]
kappa(scale(X_md1))

[1] 8.624355
```

Hence, the square of condition number $\kappa^2(\mathbf{X}^*) \approx 74.379$ is an upper bound for the VIFs which is quite large !

VIF

Now, to determine whether some covariate with corresponding column \mathbf{x}_j , can be predicted accurately using other covariates or not, we compute the variance inflation factors $VIF_j = \frac{1}{1-R_j^2}$ where R_j^2 is the coefficient of determination of the regression of $\mathbf{x}^{(j)}$ on the columns of $\mathbf{X}^{*(j)}$.

We calculate the VIF_j values for $j = 1, 2, 3, 4$ in R :-

```
require("faraway")
vif(lm(Y~A1+A2+A3+A4,data = X[-c(12,20,30,36),]))#high implies collinearity

      A1      A2      A3      A4
9.261059 9.360894 2.857795 2.751179
```

For the variables A1 and A2, we can see that the VIF values are greater than 5 and even close to 10! So we can interpret this as “the standard error of $\widehat{\beta}_1$ and $\widehat{\beta}_2$ would be $\sqrt{9.26} \approx 3.043$ and $\sqrt{9.361} \approx 3.059$ times more (respectively) than it would have been without the presence of collinearity”.

Effect of Influential Points on Collinearity

This is a very interesting observation that we have made in the dataset.

If we remove the influential points and then calculate the VIF values, we get :-

```
require("faraway")
vif(lm(Y~A1+A2+A3+A4,data = X[-c(12,20,30,36),]))#high implies collinearity

      A1      A2      A3      A4
9.261059 9.360894 2.857795 2.751179
```

But if we do the same without removing those points, we get :-

```
require("faraway")
vif(lm(Y~A1+A2+A3+A4,data = X[]))#high implies collinearity

      A1      A2      A3      A4
5.910333 6.748416 3.080737 3.488172
```

So, as we can see the VIF values increased after removal of the influential points.

This is intuitive from the fact that actual linear dependence between the covariates was being slightly nullified by the presence of such influential points.

Demonstrating Effect of Collinearity

To demonstrate how collinearity can affect the estimates badly, we deliberately introduce some random noise in the response observations ($\delta \sim N(0, 1)$) and then fit a linear model and see the changes in the estimate.

```
set.seed(2124)
lmod_per = lm(Y+10*rnorm(nrow(X)-4,s = 5) ~ A1+A2+A3+A4,data = X[-c(12,20,30,36),])
reg$coefficients
```

```
(Intercept)          A1          A2          A3          A4
-13.96310010   0.09828590   1.14837707   1.85786103   0.04817018
```

```
lmod_per$coefficients
```

```
(Intercept)          A1          A2          A3          A4
  1.304038    2.353402   -2.501280   -4.056390    1.437409
```

Hence, we can clearly see the how the estimates change a lot for introducing random noise in the response.

3.6 Remedies For Collinearity

Dealing with Collinearity

To deal with the collinearity present in the dataset, we first try to remove one of the correlated covariates “A1” or “A2” and see what improvements are observed in the variation inflation factors :-

Model	Condition Number (κ)	R_{adj}^2
$Y = \beta_0 + \beta_1 A1 + \beta_2 A2 + \beta_3 A3 + \beta_4 A4$	8.512	0.795
$Y = \beta_0 + \beta_2 A2 + \beta_3 A3 + \beta_4 A4$	5.625	0.755
$Y = \beta_0 + \beta_1 A1 + \beta_3 A3 + \beta_4 A4$	3.951	0.803

We also check for other assumptions between the models :-

Tests	Full Model	Without “A1”	Without “A2”
Shapiro-Wilk	0.1133	0.563	0.2461
Durbin-Watson	0.9734	0.7376	0.7638
Breusch-Pagan	0.6105	0.3742	0.9423

Dealing with Collinearity

Hence, the model with covariates “A1”, “A3”, “A4” seems to be a much better model in terms of both prediction and accuracy of the estimates of β . Also if we calculate the VIF values for the last model, we get them to be considerably small :-

```
vif(reg_out)
```


```
          A1          A3          A4
2.360578  2.491333  1.677148
```

This is also relatable from the fact that initially covariate “A2” had the maximum VIF value and the condition number also decreased significantly due to its removal.

Added Variable Plot

For better understanding the contribution of a covariate in the regression model, we make a scatter plot of $\mathbf{e}^{(i)} = (\mathbf{I} - \mathbf{P}_i) \mathbf{Y}$ against $(\mathbf{I} - \mathbf{P}_i) \mathbf{x}^{(i)}$ where $\mathbf{e}^{(i)}$ are the residuals of the model with variable A_i excluded and $\mathbf{x}^{(i)}$ is the column of observations of A_i . This is also called the added variable plot.

```
avPlots(lm(Y~A1+A2+A3+A4,data = X[-c(12,20,30,36),]))
```



Conclusion

From the 4 plots, we can see that slopes of the fitted lines for the added variable plots of A_1 & A_3 are much more significant than other two plots for A_2 & A_4 .

We can make some important conclusions from here.

This indicates that once predictor A_1 , A_3 , A_4 is included, A_2 can be excluded from the model for the high collinearity present between them.

Similar can be said for A_4 .

Now, for much better conclusions, we perform further model selection procedures based on several criterias.

3.7 Model Selection

Stepwise Selection

We perform the stepwise selection algorithm which performs a forward selection (FS) followed by a backward elimination (BE) using the AIC criterion and get to an optimum model.

We get the following sequence of models in the selection procedure :-

Model	AIC Value
$Y = \beta_0$	372.1803
$Y = \beta_0 + \beta_3 A3$	319.7608
$Y = \beta_0 + \beta_1 A1 + \beta_3 A3$	305.0896

We give the final model as an output we get in R :-

Stepwise Selection Method

Candidate Terms:

1 . A1
2 . A2
3 . A3
4 . A4

Step 0: AIC = 372.1803
Y ~ 1

Variables Entered/Removed:

Enter New Variables

Variable	DF	AIC	Sum Sq	RSS	R-Sq	Adj. R-Sq
A1	1	319.761	11455.113	4316.877	0.726	0.719
A3	1	324.405	10950.375	4821.615	0.694	0.687
A2	1	340.276	8736.385	7035.605	0.554	0.543
A4	1	355.432	5678.829	10093.161	0.360	0.344

- A1 added

Step 1 : AIC = 319.7608
Y ~ A1

Enter New Variables

Variable	DF	AIC	Sum Sq	RSS	R-Sq	Adj. R-Sq
A3	1	305.090	12869.415	2902.575	0.816	0.807
A4	1	318.972	11732.403	4039.587	0.744	0.731
A2	1	321.033	11529.316	4242.673	0.731	0.717

- A3 added

Step 2 : AIC = 305.0896
Y ~ A1 + A3

Remove Existing Variables

Variable	DF	AIC	Sum Sq	RSS	R-Sq	Adj. R-Sq
A3	1	319.761	11455.113	4316.877	0.726	0.719
A1	1	324.405	10950.375	4821.615	0.694	0.687

Enter New Variables

Variable	DF	AIC	Sum Sq	RSS	R-Sq	Adj. R-Sq
A2	1	306.749	12892.877	2879.113	0.817	0.803
A4	1	306.775	12891.077	2880.913	0.817	0.803

No more variables to be added or removed.

Final Model Output

Model Summary

R	0.903	RMSE	8.627
R-Squared	0.816	Coef. Var	14.066
Adj. R-Squared	0.807	MSE	74.425
Pred R-Squared	0.788	MAE	6.755

RMSE: Root Mean Square Error
MSE: Mean Square Error
MAE: Mean Absolute Error

ANOVA

	Sum of Squares	DF	Mean Square	F	Sig.
Regression	12869.415	2	6434.707	86.459	0.0000
Residual	2902.575	39	74.425		
Total	15771.990	41			

Parameter Estimates

model	Beta	Std. Error	Std. Beta	t	Sig.	lower	upper
(Intercept)	9.924	5.106		1.944	0.059	-0.403	20.251
A1	0.640	0.126	0.521	5.078	0.000	0.385	0.895
A3	1.516	0.348	0.447	4.359	0.000	0.813	2.219

Stepwise Summary

Variable	Method	AIC	RSS	Sum Sq	R-Sq	Adj. R-Sq
A1	addition	319.761	4316.877	11455.113	0.72629	0.71945
A3	addition	305.090	2902.575	12869.415	0.81597	0.80653

Hence, this method gives the model containing covariates “A1”, “A3” as the optimum one.

Best Subset Selection

We use different criterions for choosing optimal model among all the 15 possible linear models and plot the diagrams for all of them one by one.


```
X_m = X[,-c(1,2)]
X1 <- X_m[-c(12,20,30,36),]
names(X1) <- c("V1", "V2", "V3", "V4", "V5")
models<-list()
models[["V1"]] <- lm(V5~V1,X1)
models[["V2"]] <- lm(V5~V2,X1)
models[["V3"]] <- lm(V5~V3,X1)
models[["V4"]] <- lm(V5~V4,X1)
models[["V12"]] <- lm(V5~V1+V2,X1)
models[["V13"]] <- lm(V5~V1+V3,X1)
models[["V14"]] <- lm(V5~V1+V4,X1)
models[["V23"]] <- lm(V5~V2+V3,X1)
models[["V24"]] <- lm(V5~V2+V4,X1)
models[["V34"]] <- lm(V5~V3+V4,X1)
models[["V123"]] <- lm(V5~V1+V2+V3,X1)
```

```
models[["V124"]] <- lm(V5~V1+V2+V4,X1)
models[["V134"]] <- lm(V5~V1+V3+V4,X1)
models[["V234"]] <- lm(V5~V2+V3+V4,X1)
models[["V1234"]] <- lm(V5~V1+V2+V3+V4,X1)

mnames<-factor(names(models),levels = names(models))
```


Coefficient of determination (R^2)

```
R2 <- sapply(models, function(fit) summary(fit)$r.squared)
dotplot(R2 ~ mnames, type = "o", pch = 16, auto.key=list(space="right"), xlab="Models", ylab="expressi
```




R^2 & R^2_{adj}

```
adj.R2 <- sapply(models, function(fit) summary(fit)$adj.r.squared)
dotplot(R2 + adj.R2 ~ mnames, type = "o", pch = 16, auto.key=list(space="right"), xlab="Models")
```





Mallow's C_p




Akaike information criterion (AIC)

```
AIC <- sapply(models, function(fit) AIC(fit))
dotplot(AIC ~ mnames, type = "o", pch = 16, xlab="Models")
```




Bayesian information criterion (BIC)

```
BIC <- sapply(models, function(fit) BIC(fit))
dotplot(BIC ~ mnames, type = "o", pch = 16, xlab="Models")
```



Leave-One-Out CV

```
CV = NULL
for(i in 1:15)
{
  X_mdl_mat = model.matrix(models[[i]])
  head(X_mdl_mat)
  Y_vec = X1$V5
  H = X_mdl_mat%*%solve(t(X_mdl_mat)%*%X_mdl_mat)%*%t(X_mdl_mat)
  h = diag(H)
  n_h = nrow(X1)
  CV[i] = (1/n_h)*sum((Y_vec-H%*%Y_vec)^2/(1-h)^2)
}
dotplot(CV ~ mnames, type = "o", pch = 16,xlab="Models")
```



Values of different measures for all the models

We list down the values of R^2 , R_{adj}^2 , Mallows's C_p , AIC , BIC , $CV(1)$ values in one table for all the 15 models for better comparison :-

	R2	adj.R2	Cp	AIC	BIC	CV
V1	0.7262947	0.7194521	19.406493	319.7608	324.9738	112.46225
V2	0.5539177	0.5427657	55.560548	340.2757	345.4888	185.08170
V3	0.6942925	0.6866498	26.118573	324.4050	329.6180	127.88888
V4	0.3600579	0.3440593	96.220397	355.4325	360.6455	261.15078
V12	0.7309995	0.7172046	20.419718	321.0326	327.9832	119.82317
V13	0.8159665	0.8065289	2.598883	305.0896	312.0402	79.76883
V14	0.7438759	0.7307413	17.719047	318.9724	325.9231	110.28385
V23	0.7675782	0.7556591	12.747757	314.8939	321.8445	100.10949
V24	0.5635089	0.5411247	55.548917	341.3629	348.3135	188.38242
V34	0.7084529	0.6935017	25.148610	324.4131	331.3638	128.75523
V123	0.8174541	0.8030425	4.286879	306.7487	315.4370	84.42043
V124	0.7730732	0.7551579	13.595257	315.8890	324.5773	105.20886
V134	0.8173399	0.8029194	4.310820	306.7749	315.4633	83.10493

```
V234 0.7684411 0.7501601 14.566777 316.7376 325.4260 105.04886
V1234 0.8235897 0.8045183 5.000000 307.3127 317.7387 85.14023
```

Conclusion

Now, we write down the optimal models we get from different model selection criteria with corresponding values :-

Criteria	Optimum Model	Value
R_{adj}^2	$Y = \beta_0 + \beta_1 A1 + \beta_3 A3$	0.8065
Mallow's C_p	$Y = \beta_0 + \beta_1 A1 + \beta_3 A3$	2.598
AIC	$Y = \beta_0 + \beta_1 A1 + \beta_3 A3$	305.089
BIC	$Y = \beta_0 + \beta_1 A1 + \beta_3 A3$	312.0402
$CV(1)$	$Y = \beta_0 + \beta_1 A1 + \beta_3 A3$	79.76883

Hence, clearly this indicates among all the linear models, $Y = \beta_0 + \beta_1 A1 + \beta_3 A3$ is optimum based on several criteria.

This is also intuitive from the fact that here we are removing the covariates which had linear dependence.

In terms of terminology of the given dataset, the optimum predictors of $Y =$ Cirrhosis death rate are $A1 =$ Urban population & $A3 =$ Wine consumption per capita.

So the optimum fitted model can be written as :-


$$Y = 9.9241 + 0.6397A1 + 1.5159A3$$

Also we observed that all the covariates in the model are significant.

Obs vs Fitted Values

We plot the observed vs fitted values obtained using this model :-

```
X1 = X[-c(12,20,30,36),]
mod_opt = lm(Y ~ A1+A3, data = X[-c(12,20,30,36),])
plot(1:nrow(X1), X1$Y, type = "o", pch = 20, ylab = "observed & predicted values", xlab = "Index")
lines(1:nrow(X1), mod_opt$fitted.values, type = "o", pch = 20, col = "red", lty = 2)
abline(v = 1:nrow(X1), lty = 2, col = rgb(0,1,0,alpha = 0.3))
legend("topleft", legend = c("Obs", "Pred"), fill = c("black", "red"))
```



Here we can see the prediction is accurate compared to the full model.

Conclusion

Again we perform all the diagnostic tests for this final model and find the following :-

Tests	p-values
Shapiro-Wilk	0.6282
Durbin-Watson	0.645
Breusch-Pagan	0.2723
Breusch-Godfrey	0.9345

All the assumptions seem to be satisfied here. Hence we can really consider this to be a good model.

Models with Interaction Terms

One class of models that we have not considered yet are those with interaction terms (upto second order). Since there can be too many of them, we will not perform best subset selection here. Rather we again perform stepwise regression for choosing an optimal one among them.

In this class, we get the following model :-

```
mod_wt_out = lm(Y ~ A1+A2+A3+A4, data = X[-c(12,20,30,36),])
STEP_REG = stepAIC(mod_wt_out, scope = list(upper = ~(A1+A2+A3+A4)^2, lower = ~1), trace = TRUE)
```

```
Start: AIC=186.12
Y ~ A1 + A2 + A3 + A4

      Df Sum of Sq  RSS   AIC
+ A1:A4  1    186.85 2595.5 185.20
+ A2:A4  1    173.31 2609.0 185.42
```

```

- A4      1      96.77 2879.1 185.56
- A2      1      98.57 2880.9 185.58
+ A3:A4   1      153.19 2629.1 185.74
+ A1:A2   1      134.69 2647.7 186.04
<none>                2782.3 186.12
+ A2:A3   1       77.78 2704.6 186.93
+ A1:A3   1       22.44 2759.9 187.78
- A3      1      796.75 3579.1 194.70
- A1      1      869.80 3652.1 195.55

```

Step: AIC=185.2

Y ~ A1 + A2 + A3 + A4 + A1:A4

	Df	Sum of Sq	RSS	AIC
- A2	1	104.35	2699.8	184.86
<none>			2595.5	185.20
- A1:A4	1	186.85	2782.3	186.12
+ A1:A3	1	56.52	2539.0	186.28
+ A1:A2	1	7.45	2588.0	187.08
+ A3:A4	1	4.07	2591.4	187.14
+ A2:A3	1	3.27	2592.2	187.15
+ A2:A4	1	0.45	2595.0	187.19
- A3	1	674.06	3269.6	192.90

Step: AIC=184.86

Y ~ A1 + A3 + A4 + A1:A4

	Df	Sum of Sq	RSS	AIC
<none>			2699.8	184.86
+ A2	1	104.35	2595.5	185.20
- A1:A4	1	181.07	2880.9	185.58
+ A1:A3	1	78.30	2621.5	185.62
+ A3:A4	1	0.02	2699.8	186.86
- A3	1	1005.28	3705.1	196.15

STEP_REG

Call:

```
lm(formula = Y ~ A1 + A3 + A4 + A1:A4, data = X[-c(12, 20, 30,
36), ])
```

Coefficients:

(Intercept)	A1	A3	A4	A1:A4
36.182148	0.121091	1.366505	-0.448029	0.008963

This model has an adjusted R^2 value equal to 0.8299. But the main problem is this model has very high vif values and many of the predictors are not significant. So we don't consider these type of models.

Conclusion

Hence we conclude our final multiple linear regression model is :-

$$Y = 9.9241 + 0.6397A1 + 1.5159A3$$

Obviously this model also has some drawback and there is no such "best" model that we can have but this performs more or less better than most of the models hence, it is a good one.

Next we use other types of regression models with different interpretations.

3.8 Shrinkage Methods


Ridge Regression

An alternate approach to deal with collinearity is fitting a ridge regression model as it can improve the accuracy of the predictions. The ridge estimate of the model parameters is $\hat{\beta}(k) = (\mathbf{X}^T \mathbf{X} + k\mathbf{I})^{-1} \mathbf{X}^T \mathbf{Y}$ where k is the ridge parameter. For an optimal choice of k , we calculate estimates of prediction errors of the ridge predictors for different choices of k over a set of trial values. This can be expressed as $CV_k(1) = \frac{1}{n} \sum_{i=1}^n \frac{[Y_i - x_i^T \hat{\beta}(k)]^2}{[1 - a_{ii}(k)]^2}$ and choose the k_{opt} for which this quantity is minimum.

We plot the $CV_k(1)$ values for different choices of k :-

```
k_grid = 10^seq(-2,1/2,length.out = 100)
PE = NULL
X_R = as.matrix(X[-c(12,20,30,36),c(2,3,4,5,6)])
Y_R = as.matrix(X[-c(12,20,30,36),c(7)])
n = nrow(X_R)
for(i in 1:length(k_grid))
{
  k = k_grid[i]
  beta_k = solve(t(X_R)%*%X_R + k*diag(rep(1,5)))*%*%t(X_R)%*%Y_R
  A_k = X_R%*%solve(t(X_R)%*%X_R + k*diag(rep(1,5)))*%*%t(X_R)
  Y_ft_R = X_R%*%beta_k
  A_K_diag = diag(A_k)
  PE[i] = (1/n)*sum((Y_R-Y_ft_R)^2/(1-A_K_diag)^2)
}

plot(k_grid,PE,type = "l",ylab = "CV(1)",xlab = "k")
k_opt = k_grid[which(PE == min(PE))]
abline(v = k_opt)
```



We find that the $CV_k(1)$ is minimum for $k \approx 0.3898$ hence, we calculate the corresponding ridge estimates.


The fitted model then becomes :-

$$Y = 11.777 + 0.767A1 + -0.322A2 + 1.324A3 + 0.114A4$$

This model has estimated prediction error ≈ 83.612 .

This is close to the optimum OLS model that we have fitted.

We plot the observed & fitted values with the same index in the x-axis and get the following output :-



Lasso Regression

Another efficient way of model selection is using the Lasso Regression method. Here we minimize the sum of squares $\|Y - X\beta\|^2$ subject to the constraint $\sum_j |\beta_j| \leq \lambda$ for some $\lambda > 0$.

Using R, we find the Lasso Estimates of β where the value of λ is chosen using k -fold cross-validation criteria.

The optimum value of λ chosen by the criteria approximately equals ≈ 0.501 and the model is :-

```
library(glmnet)
x <- as.matrix(X[-c(12,20,30,36),c(3,4,5,6)])
y <- X[-c(12,20,30,36),7]
lambdas <- 10^seq(-1, 5, by = 0.1)

lasso_reg <- cv.glmnet(x,y, alpha = 1, lambda = lambdas, standardize = TRUE, nfolds = 10)

lambda_best <- lasso_reg$lambda.min

lasso_model <- glmnet(x,y, alpha = 1, lambda = 5, standardize = TRUE)
c(lasso_model$a0,t(lasso_model$beta))

$s0
[1] 25.2888

[[2]]
```




```
1 x 4 sparse Matrix of class "dgCMatrix"
      A1 A2      A3 A4
s0 0.4579335 . 1.013119 .
```

Hence the Lasso Estimates of the parameters $\beta = \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \\ \beta_3 \\ \beta_4 \end{pmatrix} = \begin{pmatrix} 25.28 \\ 0.457 \\ 0 \\ 1.013 \\ 0 \end{pmatrix}$.

As we can see from the output, here also, the variables “A2”, “A4” has been dropped and this also gives strong evidence in favour of the optimum linear model.

Here also make the observed and fitted plot for different index values :-



3.9 Robust Regression Methods

We have detected influential points in our dataset, and also removed them to get better models.

Now, we demonstrate using different robust regression methods how they can be used even if we have outliers in our dataset.

So we perform the rest of the methods using the full dataset, without removing any observation.

Least Absolute Deviation

Here we minimize the quantity $\sum_i |e_i(\mathbf{b})|$ i.e. $\hat{\beta}_{LAD} = \underset{\mathbf{b}}{\operatorname{argmin}} \sum_i |e_i(\mathbf{b})|$ where $e_i(\mathbf{b}) = Y_i - \mathbf{x}_i^T \mathbf{b}$.


- The estimated values of $\hat{\beta}_{LAD}$ equals :-

```
library("Lipack")
rmodel_l <- lad(formula = Y~A1+A2+A3+A4,data = X)
rmodel_l$coefficients
```

(Intercept)	A1	A2	A3	A4
-7.19977780	0.42727435	0.61749765	1.96154792	-0.02363183

We plot the observed vs fitted values obtained using this model :-

```
plot(1:nrow(X),X$Y,type = "o",pch = 20,ylab = "observed & predicted values",xlab = "Index")
lines(1:nrow(X),rmodel_l$fitted.values,type = "o",pch = 20,col = "red",lty = 2)
abline(v = 1:nrow(X),lty = 2,col = rgb(0,1,0,alpha = 0.3))
legend("topleft",legend = c("Obs","Pred"),fill = c("black","red"))
```



The plot shows here the predicted values are more or less accurate for all the observations.

- The estimated values of $\hat{\beta}_{LAD}$ equals for A1 & A3:-

```
library("Lipack")
rmodel_l <- lad(formula = Y~A1+A3,data = X)
rmodel_l$coefficients
```

(Intercept)	A1	A3
4.3545455	0.6272727	2.0909091

```
summary(rmodel_l)
```

Call:
lad(formula = Y ~ A1 + A3, data = X)

Residuals:

Min	1Q	Median	3Q	Max
-35.918	-5.514	0.000	5.720	23.845

Coefficients:


	Estimate	Std.Error	Z value	p-value
(Intercept)	4.3545	9.4835	0.4592	0.6461
A1	0.6273	0.2132	2.9417	0.0033

```
A3          2.0909   0.4852   4.3093  0.0000

Degrees of freedom: 46 total; 43 residual
Scale estimate: 11.69856
Log-likelihood: -175.0778 on 4 degrees of freedom
```

We plot the observed vs fitted values obtained using this model :-

```
plot(1:nrow(X),X$Y,type = "o",pch = 20,ylab = "observed & predicted values",xlab = "Index")
lines(1:nrow(X),rmodel_1$fitted.values,type = "o",pch = 20,col = "red",lty = 2)
abline(v = 1:nrow(X),lty = 2,col = rgb(0,1,0,alpha = 0.3))
legend("topleft",legend = c("Obs","Pred"),fill = c("black","red"))
```



The plot shows here the predicted values are more or less accurate for all the observations.

Least Median Square

Here we minimize the median of the squared residuals $\hat{\beta}_{LMS} = \underset{\mathbf{b}}{\operatorname{argmin}} \operatorname{med}_i e_i^2(\mathbf{b})$.


- Using R, we get the estimated value of $\hat{\beta}_{LMS}$ as :-

```
rmodel_1 <- lqs(Y~A1+A2+A3+A4,data = X,method = "lms")
rmodel_1$coefficients

(Intercept)          A1          A2          A3          A4
100.67646579   1.43011898  -3.48160404   3.15085397  -0.04186281
```

We plot the observed vs fitted values obtained using this model :-

```
par(mfrow = c(1,1))
plot(1:nrow(X),X$Y,type = "o",pch = 20,ylab = "observed & predicted values",xlab = "Index")
lines(1:nrow(X),rmodel_1$fitted.values,type = "o",pch = 20,col = "red",lty = 2)
abline(v = 1:nrow(X),lty = 2,col = rgb(0,1,0,alpha = 0.3))
legend("topleft",legend = c("Obs","Pred"),fill = c("black","red"))
```



We can see the prediction is quite accurate at some places where as it is bad at some others possibly due to the presence of outliers.

- Using R, we get the estimated value of $\hat{\beta}_{LMS}$ as for A1 & A3 :-

```
rmodel_l <- lqs(Y~A1+A3,data = X,method = "lms")
rmodel_l$coefficients
```


```
(Intercept)      A1      A3
 5.7138767  0.6726872  1.8555066
```

```
summary(rmodel_l)
```

	Length	Class	Mode
crit	1	-none-	numeric
sing	1	-none-	character
coefficients	3	-none-	numeric
bestone	3	-none-	numeric
fitted.values	46	-none-	numeric
residuals	46	-none-	numeric
scale	2	-none-	numeric
terms	3	terms	call
call	4	-none-	call
xlevels	0	-none-	list
model	3	data.frame	list

We plot the observed vs fitted values obtained using this model :-

```
par(mfrow = c(1,1))
plot(1:nrow(X),X$Y,type = "o",pch = 20,ylab = "observed & predicted values",xlab = "Index")
lines(1:nrow(X),rmodel_l$fitted.values,type = "o",pch = 20,col = "red",lty = 2)
abline(v = 1:nrow(X),lty = 2,col = rgb(0,1,0,alpha = 0.3))
legend("topleft",legend = c("Obs","Pred"),fill = c("black","red"))
```



We can see the prediction is quite accurate at some places where as it is bad at some others possibly due to the presence of outliers.

Least Trimmed Squares Estimate


Lastly, we compute the LTS estimates of β where we minimize the trimmed mean of the squared residuals $\hat{\beta}_{LMS} = \underset{\mathbf{b}}{\operatorname{argmin}} \frac{1}{h} \sum_{i=1}^h e_i^2(\mathbf{b})$ for some appropriate choice of h . (Here we choose $h = \lceil n/2 \rceil + 1$)

```
rmodel_l <- lqs(Y ~ A1 + A2 + A3 + A4, data = X, method = "lts")
rmodel_l$coefficients
```

(Intercept)	A1	A2	A3	A4
-21.7304500	0.2861684	1.2215938	2.4980685	-0.1567162

We plot the observed vs fitted values obtained using this model :-

```
plot(1:nrow(X), X$Y, type = "o", pch = 20, ylab = "observed & predicted values", xlab = "Index")
lines(1:nrow(X), rmodel_l$fitted.values, type = "o", pch = 20, col = "red", lty = 2)
abline(v = 1:nrow(X), lty = 2, col = rgb(0, 1, 0, alpha = 0.3))
legend("topleft", legend = c("Obs", "Pred"), fill = c("black", "red"))
```



We can see that this method performs better in terms of prediction compared to LMS estimates.

Lastly, we compute the LTS estimates of β where we minimize the trimmed mean of the squared residuals $\hat{\beta}_{LMS} = \underset{\mathbf{b}}{\operatorname{argmin}} \frac{1}{h} \sum_{i=1}^h e_i^2(\mathbf{b})$ for some appropriate choice of h . (Here we choose $h = \lceil n/2 \rceil + 1$) for A1 & A3:

```
rmodel_l <- lqs(Y ~ A1 + A3, data = X, method = "lts")
rmodel_l$coefficients
```


```
(Intercept)      A1      A3
 2.0536872  0.6331754  2.4071090
```

```
summary(rmodel_l)
```

	Length	Class	Mode
crit	1	-none-	numeric
sing	1	-none-	character
coefficients	3	-none-	numeric
bestone	3	-none-	numeric
fitted.values	46	-none-	numeric
residuals	46	-none-	numeric
scale	2	-none-	numeric
terms	3	terms	call
call	4	-none-	call
xlevels	0	-none-	list
model	3	data.frame	list

We plot the observed vs fitted values obtained using this model :-

```
plot(1:nrow(X), X$Y, type = "o", pch = 20, ylab = "observed & predicted values", xlab = "Index")
lines(1:nrow(X), rmodel_l$fitted.values, type = "o", pch = 20, col = "red", lty = 2)
abline(v = 1:nrow(X), lty = 2, col = rgb(0,1,0,alpha = 0.3))
legend("topleft", legend = c("Obs", "Pred"), fill = c("black", "red"))
```



We can see that this method performs better in terms of prediction compared to LMS estimates.

Comparative Study Of All Models

Finally, as a measure of comparison of different models, we use the root mean square error $RMSE = \sqrt{\frac{1}{n} \sum_{i=1}^n (\hat{y}_i - y_i)^2}$ where \hat{y}_i denotes the fitted values using different regression models. We calculate this measure for all the “good” models we have found so far :-

Methods	$RMSE$ values
OLS model with “A1” & “A3”	8.313178
Ridge Model	8.195655
Lasso Model	9.889319
LAD Model	10.59494
LAD Model with “A1” & “A3”	
LMS Model	11.6656
LMS Model with “A1” & “A3”	
LTS Model	11.0904
LTS Model with “A1” & “A3”	

Hence, we can conclude the Ridge and the OLS model with influential points removed with covariates “A1” & “A3” performs more or less better than the others in terms of prediction accuracy. Also it can be seen that LASSO and Ridge model brings coefficients of “A2” and “A4” to 0 and close to 0 respectively.