

Dynamics on the Moduli Space of Non-Orientable Surfaces

Sayantani Khan

University of Michigan

Tuesday, April 2nd 2024

Orientable surfaces



Orientable surfaces



Non-orientable surfaces




Goal

Understand the collection of geometric structures we can put on a topological surface.

Goal

Understand the collection of geometric structures we can put on a topological surface.

In particular, understand the set of metrics on a surface with curvature -1 .



$$\mathcal{M}(\mathcal{S}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{S}_g\}$$

$$\mathcal{M}(\mathcal{S}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{S}_g\}$$

- ▶ $\mathcal{M}(\mathcal{S}_g)$ is $6g - 6$ dimensional.

$$\mathcal{M}(\mathcal{S}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{S}_g\}$$

- ▶ $\mathcal{M}(\mathcal{S}_g)$ is $6g - 6$ dimensional.
- ▶ $\mathcal{M}(\mathcal{S}_g)$ is a metric space: x and y in $\mathcal{M}(\mathcal{S}_g)$ are close if the derivative of some map $x \rightarrow y$ maps circles to “almost” circles.

$$\mathcal{M}(\mathcal{S}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{S}_g\}$$

- ▶ $\mathcal{M}(\mathcal{S}_g)$ is $6g - 6$ dimensional.
- ▶ $\mathcal{M}(\mathcal{S}_g)$ is a metric space: x and y in $\mathcal{M}(\mathcal{S}_g)$ are close if the derivative of some map $x \rightarrow y$ maps circles to “almost” circles.
- ▶ Unit (co)tangent bundle of $\mathcal{M}(\mathcal{S}_g)$ is non-compact, but finite volume, and admits a geodesic flow (and an $SL(2, \mathbb{R})$ action) with good dynamical properties.


$$\mathcal{M}(\mathcal{S}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{S}_g\}$$

- ▶ $\mathcal{M}(\mathcal{S}_g)$ is $6g - 6$ dimensional.
- ▶ $\mathcal{M}(\mathcal{S}_g)$ is a metric space: x and y in $\mathcal{M}(\mathcal{S}_g)$ are close if the derivative of some map $x \rightarrow y$ maps circles to “almost” circles.
- ▶ Unit (co)tangent bundle of $\mathcal{M}(\mathcal{S}_g)$ is non-compact, but finite volume, and admits a geodesic flow (and an $SL(2, \mathbb{R})$ action) with good dynamical properties.


Guiding principle: Dynamics on (co)tangent bundle should have analogies with dynamics of the $SL(2, \mathbb{R})$ action on $SL(2, \mathbb{R})/SL(2, \mathbb{Z})$.

$$S^1\mathcal{M}(\mathcal{S}_g) = \{\text{Area 1 half-translation surface structures on } \mathcal{S}_g\}$$

$$S^1\mathcal{M}(\mathcal{S}_g) = \{\text{Area 1 half-translation surface structures on } \mathcal{S}_g\}$$



$$S^1\mathcal{M}(\mathcal{S}_g) = \{\text{Area 1 half-translation surface structures on } \mathcal{S}_g\}$$



Theorem (Masur's criterion)

If the vertical flow on translation surface is not uniquely ergodic, then the geodesic ray in $\mathcal{M}(\mathcal{S}_g)$ escapes to infinity.

$$\mathcal{M}(\mathcal{N}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{N}_g\}$$

$$\mathcal{M}(\mathcal{N}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{N}_g\}$$

- ▶ $\mathcal{M}(\mathcal{N}_g)$ is $3g - 6$ dimensional.

$$\mathcal{M}(\mathcal{N}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{N}_g\}$$

- ▶ $\mathcal{M}(\mathcal{N}_g)$ is $3g - 6$ dimensional.
- ▶ $\mathcal{M}(\mathcal{N}_g)$ is a metric space, with a similarly defined metric.

$$\mathcal{M}(\mathcal{N}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{N}_g\}$$

- ▶ $\mathcal{M}(\mathcal{N}_g)$ is $3g - 6$ dimensional.
- ▶ $\mathcal{M}(\mathcal{N}_g)$ is a metric space, with a similarly defined metric.
- ▶ Unit (co)tangent bundle of $\mathcal{M}(\mathcal{N}_g)$ is non-compact, and also infinite volume (with respect to the canonical volume form).

$$\mathcal{M}(\mathcal{N}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{N}_g\}$$

- ▶ $\mathcal{M}(\mathcal{N}_g)$ is $3g - 6$ dimensional.
- ▶ $\mathcal{M}(\mathcal{N}_g)$ is a metric space, with a similarly defined metric.
- ▶ Unit (co)tangent bundle of $\mathcal{M}(\mathcal{N}_g)$ is non-compact, and also infinite volume (with respect to the canonical volume form).
- ▶ $S^1\mathcal{M}(\mathcal{N}_g)$ admits a geodesic flow but not an $SL(2, \mathbb{R})$ action: however, the dynamical properties are not great.

$$\mathcal{M}(\mathcal{N}_g) = \{\text{Curvature } -1 \text{ metrics on } \mathcal{N}_g\}$$

- ▶ $\mathcal{M}(\mathcal{N}_g)$ is $3g - 6$ dimensional.
- ▶ $\mathcal{M}(\mathcal{N}_g)$ is a metric space, with a similarly defined metric.
- ▶ Unit (co)tangent bundle of $\mathcal{M}(\mathcal{N}_g)$ is non-compact, and also infinite volume (with respect to the canonical volume form).
- ▶ $S^1\mathcal{M}(\mathcal{N}_g)$ admits a geodesic flow but not an $SL(2, \mathbb{R})$ action: however, the dynamical properties are not great.

Guiding principle?

Goal

Analyze generic tangent vector in $S^1\mathcal{M}(\mathcal{N}_g)$.

Goal

Analyze generic tangent vector in $S^1\mathcal{M}(\mathcal{N}_g)$.

- Analyze vertical flow on translation surface.

Goal


Analyze generic tangent vector in $S^1\mathcal{M}(\mathcal{N}_g)$.

- ▶ Analyze vertical flow on translation surface.
- ▶ Will suffice to look at first return map to a horizontal arc.

Goal

Analyze generic tangent vector in $S^1\mathcal{M}(\mathcal{N}_g)$.


- ▶ Analyze vertical flow on translation surface.
- ▶ Will suffice to look at first return map to a horizontal arc.



Goal

Analyze generic tangent vector in $S^1\mathcal{M}(\mathcal{N}_g)$.


- ▶ Analyze vertical flow on translation surface.
- ▶ Will suffice to look at first return map to a horizontal arc.



Goal

Analyze generic tangent vector in $S^1\mathcal{M}(\mathcal{N}_g)$.

- ▶ Analyze vertical flow on translation surface.
- ▶ Will suffice to look at first return map to a horizontal arc.



Theorem

Almost every geodesic in $S^1\mathcal{M}(\mathcal{N}_g)$ escapes to infinity.

$\mathrm{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

$\mathrm{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\mathrm{MCG}(\mathcal{N}_g)$ is finitely generated.

$\mathrm{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\mathrm{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.

$\text{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\text{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.

$\text{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\text{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.

Geometrically finite Γ action on \mathbb{H}

$\text{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\text{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.

Geometrically finite Γ action on \mathbb{H}

- ▶ Γ is finitely generated.

$\text{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\text{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.

Geometrically finite Γ action on \mathbb{H}

- ▶ Γ is finitely generated.
- ▶ Γ action is infinite covolume (with respect to the Liouville measure).

$\text{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\text{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.

Geometrically finite Γ action on \mathbb{H}

- ▶ Γ is finitely generated.
- ▶ Γ action is infinite covolume (with respect to the Liouville measure).
- ▶ Almost no geodesics recur.

$\text{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\text{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.

Geometrically finite Γ action on \mathbb{H}

- ▶ Γ is finitely generated.
- ▶ Γ action is infinite covolume (with respect to the Liouville measure).
- ▶ Almost no geodesics recur.
- ▶ Limit set of Γ is a subset of the boundary with smaller Hausdorff dimension.

$MCG(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $MCG(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.
- ▶ **Question:** What is the limit set for the $MCG(\mathcal{N}_g)$ action?

Geometrically finite Γ action on \mathbb{H}

- ▶ Γ is finitely generated.
- ▶ Γ action is infinite covolume (with respect to the Liouville measure).
- ▶ Almost no geodesics recur.
- ▶ Limit set of Γ is a subset of the boundary with smaller Hausdorff dimension.

$\text{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\text{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.
- ▶ **Question:** What is the limit set for the $\text{MCG}(\mathcal{N}_g)$ action?

Geometrically finite Γ action on \mathbb{H}

- ▶ Γ is finitely generated.
- ▶ Γ action is infinite covolume (with respect to the Liouville measure).
- ▶ Almost no geodesics recur.
- ▶ Limit set of Γ is a subset of the boundary with smaller Hausdorff dimension.
- ▶ Have a Patterson-Sullivan measure on the limit set with “good” dynamical properties.

$\text{MCG}(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $\text{MCG}(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.
- ▶ **Question:** What is the limit set for the $\text{MCG}(\mathcal{N}_g)$ action?
- ▶ **Question:** Can we construct a Patterson-Sullivan measure on the limit set with “good” dynamical properties?

Geometrically finite Γ action on \mathbb{H}

- ▶ Γ is finitely generated.
- ▶ Γ action is infinite covolume (with respect to the Liouville measure).
- ▶ Almost no geodesics recur.
- ▶ Limit set of Γ is a subset of the boundary with smaller Hausdorff dimension.
- ▶ Have a Patterson-Sullivan measure on the limit set with “good” dynamical properties.

$MCG(\mathcal{N}_g)$ -action on $\mathcal{T}(\mathcal{N}_g)$

- ▶ $MCG(\mathcal{N}_g)$ is finitely generated.
- ▶ The action is infinite covolume.
- ▶ Almost no geodesics recur.
- ▶ **Question:** What is the limit set for the $MCG(\mathcal{N}_g)$ action?
- ▶ **Question:** Can we construct a Patterson-Sullivan measure on the limit set with “good” dynamical properties?

Geometrically finite Γ action on \mathbb{H}

- ▶ Γ is finitely generated.
- ▶ Γ action is infinite covolume (with respect to the Liouville measure).
- ▶ Almost no geodesics recur.
- ▶ Limit set of Γ is a subset of the boundary with smaller Hausdorff dimension.
- ▶ Have a Patterson-Sullivan measure on the limit set with “good” dynamical properties.

Theorem (K., Erlandsson-Gendulph-Pasquinelli-Souto)

The limit set of the $MCG(\mathcal{N}_g)$ action on $\mathcal{T}(\mathcal{N}_g)$ is $\mathbb{P}\mathcal{MF}^+(\mathcal{N}_g)$.

What do we need for “good” Patterson-Sullivan measures?

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $\text{CAT}(k)$ for $k \leq 0$, etc.

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $CAT(k)$ for $k \leq 0$, etc.
- ▶ Teichmüller spaces are not hyperbolic on the nose, but somewhat hyperbolic.

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $\text{CAT}(k)$ for $k \leq 0$, etc.
- ▶ Teichmüller spaces are not hyperbolic on the nose, but somewhat hyperbolic.

Thick part

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $\text{CAT}(k)$ for $k \leq 0$, etc.
- ▶ Teichmüller spaces are not hyperbolic on the nose, but somewhat hyperbolic.

Thick part

- ▶ Region where all curves on underlying surface are longer than ε .

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $CAT(k)$ for $k \leq 0$, etc.
- ▶ Teichmüller spaces are not hyperbolic on the nose, but somewhat hyperbolic.

Thick part

- ▶ Region where all curves on underlying surface are longer than ε .
- ▶ Geometry of thick region mostly governed by curve complex of surface, which is hyperbolic.

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $CAT(k)$ for $k \leq 0$, etc.
- ▶ Teichmüller spaces are not hyperbolic on the nose, but somewhat hyperbolic.

Thick part

- ▶ Region where all curves on underlying surface are longer than ε .
- ▶ Geometry of thick region mostly governed by curve complex of surface, which is hyperbolic.

Thin part

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $CAT(k)$ for $k \leq 0$, etc.
- ▶ Teichmüller spaces are not hyperbolic on the nose, but somewhat hyperbolic.

Thick part

- ▶ Region where all curves on underlying surface are longer than ε .
- ▶ Geometry of thick region mostly governed by curve complex of surface, which is hyperbolic.

Thin part

- ▶ Region where some curve γ on underlying surface is shorter than ε .

What do we need for “good” Patterson-Sullivan measures?

- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $CAT(k)$ for $k \leq 0$, etc.
- ▶ Teichmüller spaces are not hyperbolic on the nose, but somewhat hyperbolic.

Thick part

- ▶ Region where all curves on underlying surface are longer than ε .
- ▶ Geometry of thick region mostly governed by curve complex of surface, which is hyperbolic.

Thin part

- ▶ Region where some curve γ on underlying surface is shorter than ε .
- ▶ Metric in this region is L^∞ product of metrics on subsurfaces (Minsky's product region theorem).

What do we need for “good” Patterson-Sullivan measures?


- ▶ Classically, Patterson-Sullivan measures were constructed for hyperbolic spaces.
- ▶ Then variable negative curvature, Gromov hyperbolic, $CAT(k)$ for $k \leq 0$, etc.
- ▶ Teichmüller spaces are not hyperbolic on the nose, but somewhat hyperbolic.

Thick part

- ▶ Region where all curves on underlying surface are longer than ε .
- ▶ Geometry of thick region mostly governed by curve complex of surface, which is hyperbolic.

Thin part

- ▶ Region where some curve γ on underlying surface is shorter than ε .
- ▶ Metric in this region is L^∞ product of metrics on subsurfaces (Minsky's product region theorem).



Statistical convex-cocompactness

Statistical convex-cocompactness

- Suffices to show geodesics enter the thin part with exponentially low probabilities. This is the notion of *statistical convex-cocompactness*, and Coulon and Yang showed that this is enough for “good” Patterson-Sullivan theory.

Statistical convex-cocompactness

- ▶ Suffices to show geodesics enter the thin part with exponentially low probabilities. This is the notion of *statistical convex-cocompactness*, and Coulon and Yang showed that this is enough for “good” Patterson-Sullivan theory.
- ▶ For Teichmüller spaces of orientable surfaces, Eskin and Mirzakhani proved statistical convex-cocompactness.

Statistical convex-cocompactness

- ▶ Suffices to show geodesics enter the thin part with exponentially low probabilities. This is the notion of *statistical convex-cocompactness*, and Coulon and Yang showed that this is enough for “good” Patterson-Sullivan theory.
- ▶ For Teichmüller spaces of orientable surfaces, Eskin and Mirzakhani proved statistical convex-cocompactness.
- ▶ They did this by studying a random walk on Teichmüller space instead, and show good recurrence properties for the random walk.

Statistical convex-cocompactness

- ▶ Suffices to show geodesics enter the thin part with exponentially low probabilities. This is the notion of *statistical convex-cocompactness*, and Coulon and Yang showed that this is enough for “good” Patterson-Sullivan theory.
- ▶ For Teichmüller spaces of orientable surfaces, Eskin and Mirzakhani proved statistical convex-cocompactness.
- ▶ They did this by studying a random walk on Teichmüller space instead, and show good recurrence properties for the random walk.
- ▶ There are two obstructions in adapting their random walk based techniques to the non-orientable setting.

Statistical convex-cocompactness

- ▶ Suffices to show geodesics enter the thin part with exponentially low probabilities. This is the notion of *statistical convex-cocompactness*, and Coulon and Yang showed that this is enough for “good” Patterson-Sullivan theory.
- ▶ For Teichmüller spaces of orientable surfaces, Eskin and Mirzakhani proved statistical convex-cocompactness.
- ▶ They did this by studying a random walk on Teichmüller space instead, and show good recurrence properties for the random walk.
- ▶ There are two obstructions in adapting their random walk based techniques to the non-orientable setting.
- ▶ Obstruction 1: Thin region where some *one-sided curve* gets short does not have good recurrence properties for the random walk.

Statistical convex-cocompactness

- ▶ Suffices to show geodesics enter the thin part with exponentially low probabilities. This is the notion of *statistical convex-cocompactness*, and Coulon and Yang showed that this is enough for “good” Patterson-Sullivan theory.
- ▶ For Teichmüller spaces of orientable surfaces, Eskin and Mirzakhani proved statistical convex-cocompactness.
- ▶ They did this by studying a random walk on Teichmüller space instead, and show good recurrence properties for the random walk.
- ▶ There are two obstructions in adapting their random walk based techniques to the non-orientable setting.
- ▶ Obstruction 1: Thin region where some *one-sided curve* gets short does not have good recurrence properties for the random walk.

Theorem (K.)

The map $(\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g), \text{induced path metric}) \rightarrow (\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g), \text{Teichmüller metric})$ is $(1 + \varepsilon_d)$ bi-Lipschitz.

Statistical convex-cocompactness

- ▶ Suffices to show geodesics enter the thin part with exponentially low probabilities. This is the notion of *statistical convex-cocompactness*, and Coulon and Yang showed that this is enough for “good” Patterson-Sullivan theory.
- ▶ For Teichmüller spaces of orientable surfaces, Eskin and Mirzakhani proved statistical convex-cocompactness.
- ▶ They did this by studying a random walk on Teichmüller space instead, and show good recurrence properties for the random walk.
- ▶ There are two obstructions in adapting their random walk based techniques to the non-orientable setting.
- ▶ Obstruction 1: Thin region where some *one-sided curve* gets short does not have good recurrence properties for the random walk.

Theorem (K.)

The map $(\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g), \text{induced path metric}) \rightarrow (\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g), \text{Teichmüller metric})$ is $(1 + \varepsilon_d)$ bi-Lipschitz.

- ▶ Obstruction 2: It's not obvious that the volume growth entropy of $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)$ is equal to the lattice point growth entropy for the $\text{MCG}(\mathcal{N}_g)$ action.

How to understand random walks on Teichmüller space

How to understand random walks on Teichmüller space


- ▶ Random walk is with respect to a net in $\mathcal{T}(\mathcal{N}_g)$: the Teichmüller metric controls the law of the random walk.

How to understand random walks on Teichmüller space

- ▶ Random walk is with respect to a net in $\mathcal{T}(\mathcal{N}_g)$: the Teichmüller metric controls the law of the random walk.
- ▶ In the thin part, we understand the Teichmüller metric really well, thanks to Minsky.


How to understand random walks on Teichmüller space

- ▶ Random walk is with respect to a net in $\mathcal{T}(\mathcal{N}_g)$: the Teichmüller metric controls the law of the random walk.
- ▶ In the thin part, we understand the Teichmüller metric really well, thanks to Minsky.



How to understand random walks on Teichmüller space


- ▶ Random walk is with respect to a net in $\mathcal{T}(\mathcal{N}_g)$: the Teichmüller metric controls the law of the random walk.
- ▶ In the thin part, we understand the Teichmüller metric really well, thanks to Minsky.



- ▶ Since the ball is approximately an L^∞ product, picking a point uniformly at random is equivalent to picking a points *independently* on the two coordinates.

How to understand random walks on Teichmüller space


- ▶ Random walk is with respect to a net in $\mathcal{T}(\mathcal{N}_g)$: the Teichmüller metric controls the law of the random walk.
- ▶ In the thin part, we understand the Teichmüller metric really well, thanks to Minsky.



- ▶ Since the ball is approximately an L^∞ product, picking a point uniformly at random is equivalent to picking a points *independently* on the two coordinates.
- ▶ We can show strong recurrence for random walks on a horoball in \mathbb{H} .

How to understand random walks on Teichmüller space

- ▶ Random walk is with respect to a net in $\mathcal{T}(\mathcal{N}_g)$: the Teichmüller metric controls the law of the random walk.
- ▶ In the thin part, we understand the Teichmüller metric really well, thanks to Minsky.



- ▶ Since the ball is approximately an L^∞ product, picking a point uniformly at random is equivalent to picking a points *independently* on the two coordinates.
- ▶ We can show strong recurrence for random walks on a horoball in \mathbb{H} .
- ▶ For one-sided thin regions, we get a symmetric random walk on \mathbb{Z} , which is not strongly recurrent.

Volume entropy vs lattice point entropy

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.
- ▶ Hope: the entropies might be equal since $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)/\text{MCG}(\mathcal{N}_g)$ has finite volume.

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.
- ▶ Hope: the entropies might be equal since $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)/\text{MCG}(\mathcal{N}_g)$ has finite volume.
- ▶ Technical tool to prove a statement like this was invented recently by Dowdall and Masur: *complexity length*.

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.
- ▶ Hope: the entropies might be equal since $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)/\text{MCG}(\mathcal{N}_g)$ has finite volume.
- ▶ Technical tool to prove a statement like this was invented recently by Dowdall and Masur: *complexity length*.
- ▶ Key idea is based off of Minsky's theorem again: we need to understand how volume grows in product region, and therefore, a horoball.

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.
- ▶ Hope: the entropies might be equal since $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)/\text{MCG}(\mathcal{N}_g)$ has finite volume.
- ▶ Technical tool to prove a statement like this was invented recently by Dowdall and Masur: *complexity length*.
- ▶ Key idea is based off of Minsky's theorem again: we need to understand how volume grows in product region, and therefore, a horoball.

Volume growth for ball of radius R in $\mathbb{H} \sim \exp(2R)$

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.
- ▶ Hope: the entropies might be equal since $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)/\text{MCG}(\mathcal{N}_g)$ has finite volume.
- ▶ Technical tool to prove a statement like this was invented recently by Dowdall and Masur: *complexity length*.
- ▶ Key idea is based off of Minsky's theorem again: we need to understand how volume grows in product region, and therefore, a horoball.

Volume growth for ball of radius R in $\mathbb{H} \sim \exp(2R)$

Volume growth for ball of radius R intersected with a horoball $\sim \exp(R)$

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.
- ▶ Hope: the entropies might be equal since $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)/\text{MCG}(\mathcal{N}_g)$ has finite volume.
- ▶ Technical tool to prove a statement like this was invented recently by Dowdall and Masur: *complexity length*.
- ▶ Key idea is based off of Minsky's theorem again: we need to understand how volume grows in product region, and therefore, a horoball.

Volume growth for ball of radius R in $\mathbb{H} \sim \exp(2R)$

Volume growth for ball of radius R intersected with a horoball $\sim \exp(R)$

- ▶ Gap in exponential growth rates tells us that the thin part does not contribute a significant fraction of total volume.

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.
- ▶ Hope: the entropies might be equal since $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)/\text{MCG}(\mathcal{N}_g)$ has finite volume.
- ▶ Technical tool to prove a statement like this was invented recently by Dowdall and Masur: *complexity length*.
- ▶ Key idea is based off of Minsky's theorem again: we need to understand how volume grows in product region, and therefore, a horoball.

Volume growth for ball of radius R in $\mathbb{H} \sim \exp(2R)$

Volume growth for ball of radius R intersected with a horoball $\sim \exp(R)$

- ▶ Gap in exponential growth rates tells us that the thin part does not contribute a significant fraction of total volume.

Theorem (K.)

For all surfaces S , and any $\varepsilon_t > 0$, we have the following equality of entropies.

$$h_V(\mathcal{T}_{\varepsilon_t}^-(S)) = h_{LP}(\mathcal{T}_{\varepsilon_t}^-(S))$$

Volume entropy vs lattice point entropy

- ▶ The random walk argument gives us an estimate in terms of the volume growth entropy. Statistical convex cocompactness requires an estimate in terms of lattice point entropy.
- ▶ Hope: the entropies might be equal since $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)/\text{MCG}(\mathcal{N}_g)$ has finite volume.
- ▶ Technical tool to prove a statement like this was invented recently by Dowdall and Masur: *complexity length*.
- ▶ Key idea is based off of Minsky's theorem again: we need to understand how volume grows in product region, and therefore, a horoball.

Volume growth for ball of radius R in $\mathbb{H} \sim \exp(2R)$

Volume growth for ball of radius R intersected with a horoball $\sim \exp(R)$

- ▶ Gap in exponential growth rates tells us that the thin part does not contribute a significant fraction of total volume.

Theorem (K.)

For all surfaces S , and any $\varepsilon_t > 0$, we have the following equality of entropies.

$$h_V(\mathcal{T}_{\varepsilon_t}^-(S)) = h_{LP}(\mathcal{T}_{\varepsilon_t}^-(S))$$

Theorem (K.)

The action of $\text{MCG}(\mathcal{N}_g)$ on $\mathcal{T}_{\varepsilon_t}^-(\mathcal{N}_g)$ is statistically convex-cocompact.

On complexity length

On complexity length

Goal

Want to count net points (which stay in thin part) distance R away from base point p , with good upper bounds.

On complexity length

Goal

Want to count net points (which stay in thin part) distance R away from base point p , with good upper bounds.

- ▶ Results of Athreya, Bufetov, Eskin, and Mirzakhani can be applied, but the upper bound in this setting is not tight enough.

On complexity length

Goal

Want to count net points (which stay in thin part) distance R away from base point p , with good upper bounds.


- ▶ Results of Athreya, Bufetov, Eskin, and Mirzakhani can be applied, but the upper bound in this setting is not tight enough.
- ▶ Reduction: Assume geodesic joining p to a net point n stays entirely within a product region where γ is short.

On complexity length

Goal

Want to count net points (which stay in thin part) distance R away from base point p , with good upper bounds.

- ▶ Results of Athreya, Bufetov, Eskin, and Mirzakhani can be applied, but the upper bound in this setting is not tight enough.
- ▶ Reduction: Assume geodesic joining p to a net point n stays entirely within a product region where γ is short.




On complexity length

Goal

Want to count net points (which stay in thin part) distance R away from base point p , with good upper bounds.

- ▶ Results of Athreya, Bufetov, Eskin, and Mirzakhani can be applied, but the upper bound in this setting is not tight enough.
- ▶ Reduction: Assume geodesic joining p to a net point n stays entirely within a product region where γ is short.




- ▶ Can count independently along both axes: count along $\mathcal{T}(S \setminus \gamma)$ grows like $\exp((6g-8)R)$, and count along $\mathcal{T}(\gamma)$ grows like $\exp(R)$.

On complexity length

Goal

Want to count net points (which stay in thin part) distance R away from base point p , with good upper bounds.

- ▶ Results of Athreya, Bufetov, Eskin, and Mirzakhani can be applied, but the upper bound in this setting is not tight enough.
- ▶ Reduction: Assume geodesic joining p to a net point n stays entirely within a product region where γ is short.



- ▶ Can count independently along both axes: count along $\mathcal{T}(S \setminus \gamma)$ grows like $\exp((6g - 8)R)$, and count along $\mathcal{T}(\gamma)$ grows like $\exp(R)$.
- ▶ Complexity length is based on this idea, but accounts for the fact that the geodesic joining p and n could travel through several product regions while staying in the thin part.

What next?

What next?

- ▶ Figure out what is wrong with the growth of simple closed curves on non-orientable surfaces.

What next?

- ▶ Figure out what is wrong with the growth of simple closed curves on non-orientable surfaces.
- ▶ Show that the action of $\text{MCG}(\mathcal{N}_g)$ on $\mathcal{T}(\mathcal{N}_g)$ is statistically convex cocompact.

What next?

- ▶ Figure out what is wrong with the growth of simple closed curves on non-orientable surfaces.
- ▶ Show that the action of $\text{MCG}(\mathcal{N}_g)$ on $\mathcal{T}(\mathcal{N}_g)$ is statistically convex cocompact.
- ▶ Figure out computational (or other) methods to actually estimate the entropies.

What next?

- ▶ Figure out what is wrong with the growth of simple closed curves on non-orientable surfaces.
- ▶ Show that the action of $\text{MCG}(\mathcal{N}_g)$ on $\mathcal{T}(\mathcal{N}_g)$ is statistically convex cocompact.
- ▶ Figure out computational (or other) methods to actually estimate the entropies.

