

**Effect of Dietary Patterns on Chronic Kidney Disease (CKD) Measures (ACR), and on
the Mortality of CKD Patients**

by

Sayed Ahmed

BSc Eng. in Computer Science and Engineering, BUET, Bangladesh, 2001

MSc in Computer Science, University of Manitoba, Canada, 2006

A Major Research Paper

Presented to Ryerson University

in partial fulfillment of the requirements for the degree of

Master of Science

in the Program of

Data Science and Analytics

Toronto, Ontario, Canada, 2019

© Sayed Ahmed 2019

Supervisor

Dr. Youcef Derbal

Information Technology Department

Ted Rogers School of ITM

Ryerson University

AUTHOR'S DECLARATION FOR ELECTRONIC SUBMISSION OF A MAJOR RESEARCH PROJECT (MRP)

I hereby declare that I am the sole author of this Major Research Paper. This is a true copy of the MRP, including any required final revisions.

I authorize Ryerson University to lend this MRP to other institutions or individuals for the purpose of scholarly research.

I further authorize Ryerson University to reproduce this MRP by photocopying or by other means, in total or in part, at the request of other institutions or individuals for the purpose of scholarly research.

I understand that my MRP may be made electronically available to the public.

Sayed Ahmed

Effect of Dietary Patterns on Chronic Kidney Disease (CKD) Measures (ACR), and on the Mortality of CKD Patients

Sayed Ahmed

Master of Science 2019

Data Science and Analytics

Ryerson University

Abstract

Chronic Kidney Disease (CKD) leading to End-Stage Renal Disease (ESRD) is very prevalent today. Over 37 millions of Americans have CKD. CKD/ESRD and interrelated diseases cause a majority of the early deaths. Many research studies have investigated the effects of drugs on CKD. However, less attention has been given to the study of the dietary patterns on CKD. This research study has uncovered significant correlations between dietary patterns and CKD mortality as well as identified diagnostic markers for CKD such as the Albumin to Creatinine Ratio (ACR). In this project, Dietary surveys from NHANES, and CKD Mortality dataset from USRDS were utilized to study the correlation between dietary patterns and morbidity of CKD patients. Principal Component Analysis and Regression were utilized to find the effect. Machine Learning Approaches including Regression, and Bayesian were applied to predict ACR values. Vegetables such as Other Vegetables, Starchy Vegetables, and Red and Orange Vegetables showed moderate positive correlations with mortality whereas Grains, Fruits, Alcohol, Sugar, and Nuts showed negative correlations. ACR values were not found strongly correlated with dietary patterns. For ACR value prediction, 10 Fold Cross Validations with Polynomial Regression showed 95% accuracy.

Keywords:

CKD, ESRD, Dietary Patterns, Mortality, ACR, Polynomial Regression, Bayesian

Acknowledgements

I am very grateful to Professor Dr. Youcef Derbal to supervise this research project towards my MSc. Dr. Derbal has been a great support throughout the work. Dr. Derbal has helped to formalize the problem, referred to valuable ideas, articles, and research papers, and provided valuable feedback. I am grateful for the guide and direction that I have received from Dr. Derbal.

Thank you, Professor Dr. Derbal.

Contents

Author's Declaration	2
Acknowledgement	4
1 Introduction	8
1.1 How CKD is Identified and Measured	8
2 Literature Review	9
3 Exploratory Study	11
4 Methodology	11
4.1 Methodology Overview	11
4.2 Study Selection	14
4.3 Data Synthesis	14
5 Experiment Design	15
6 Results	17
6.1 Food Groups, Food Subgroups, Food Nutrients and CKD Mortality	17
6.2 Food Subgroups and Albumin to Creatinine Ratio (ACR) Association	21
6.3 Predictability of ACR values based on Dietary patterns	21
7 Conclusions	22
8 Appendix	23
9 References	23

List of Figures

1	Exploratory Analysis and Representative Output	12
2	Methodology in a Diagram	13
3	Food Groups (Grains) and Mortality - Negative Correlations	18
4	Food Groups (Fruits) and Mortality - Negative Correlations	18
5	Food Groups and Mortality - Positive Correlations (Vegetables)	19
6	Food Subgroups and Mortality - Positive Correlations	20
7	Food Subgroups and Mortality - Negative Correlations	20

List of Tables

1	Regression Output from Exploratory Analysis using Excel (Data Analysis Module)	11
2	Set 1: Association of Food Groups with CKD Mortality	15
3	Set 2: Association of Food Subgroups with CKD Mortality	15
4	Set 3: Effect of Food Groups on ACR	16
5	Set 4: Effect of Nutrients on ACR	16
6	Set 5: Effect of Food Subgroups on ACR	16
7	Set 6: Use Regression and Bayesian to predict ACR using Food Subgroups intake	16
8	Set 7: Find effect on CKD Mortality using survey data with no data aggregation by Age Groups	17
9	Set 8: Utilize Regression and Bayesian to predict Mortality using survey data with no data aggregation by Age Groups	17
10	Set 9: Association between Food Groups/Subgroups and Remaining Life for CKD Patients: Use not aggregated dietary intake data	17
11	Outcome when ACR Values are used as the Target	22
12	Outcome when ACR Category is used as the Target	22

1 Introduction

Chronic kidney disease (CKD) is very prevalent in today's world and CKD incidents are continually increasing whereby 10 to 13% of the US population are affected by Chronic Kidney Disease. CKD/ESRD and other interrelated diseases such as Hypertension, Heart Diseases, and Diabetes cause the majority of early deaths [31]. In addition to kidney failure, CKD is also a major cause of death from stroke, and heart diseases. On the other hand, hypertension and diabetes are also major causes of CKD. CKD is not reversible, progressive and gradually reduces kidney function.

Primary causes of CKD include High Blood Pressure, and Diabetes. Other causes include infection, kidney stones, genetics, genetical polycystic kidney disease, certain foods and food habits, pain killers, and drug usage/abuse. As CKDs are neither curable nor reversible controlling diabetes and blood pressure with or without medication can slow the progress of CKDs in most cases. As Kidneys filter waste products and our diet produces those waste products controlling diet have an effect on how much work kidney has to perform, and how well kidneys will function. Studies show that drugs as well as lifestyle choices (food, diet, exercise) can prevent CKD, slow the progression of CKD [29], delay dialysis and kidney transplantation; consequently can prevent early deaths. Although there are many studies on the effect of drugs to control CKD and related complications, there are few studies on the effect of diets, dietary patterns, and lifestyles [29]. There are studies on how controlling nutrients/chemicals in food items can help to prevent or to slow the progression of CKD. CKD patients are also provided recommendations on certain chemicals or food items. However, adhering to the recommended amount of nutrients and/or food item is challenging. Hence, there is an emerging trend where the effect is studied utilizing dietary patterns with food groups and food subgroups rather than nutrients/chemicals in food or individual food item. This research analyzes the effect of dietary patterns, using food groups and food subgroups, on the mortality and survival of CKD patients. Also studied in this research is the correlation between dietary patterns and ACR. Additionally, ACR values are predicted based on a dataset utilizing machine learning approaches.

1.1 How CKD is Identified and Measured

CKD is identified with one of two measures such as a blood test named Glomerular Filtration Rate (GFR) or a urine test named Albumin to Creatinine Ratio (ACR). ACR values less than 30 indicates no CKD or mild CKD. ACR values between 30 and 300 indicate moderate CKD. ACR values >300 indicates severe CKD. Patients are diagnosed with a CKD disease if the ACR values persist within the above ranges for three months. GFR is measured in $\text{ml/min}/1.73 \text{ m}^2$. CKDs as measured with GFRs are described in stages such as Stage 1 with normal or high GFR (GFR greater than or equal to 90 mL/min), Stage 2 with Mild CKD (GFR = $60\text{-}89 \text{ mL/min}$), Stage 3A with Moderate CKD (GFR = $45\text{-}59 \text{ mL/min}$), Stage 3B with Moderate CKD (GFR = $30\text{-}44 \text{ mL/min}$), Stage 4 with Severe CKD (GFR = $15\text{-}29 \text{ mL/min}$), Stage 5 with End Stage CKD (GFR $<15 \text{ mL/min}$) [5]. At stage 5, patients loose complete kidney function. At this point, patient survival will require either dialysis or organ transplantation.

2 Literature Review

Kidney patients are commonly given dietary advice about intake of individual nutrients, chemicals, food items, or about their whole eating patterns. However, dietary recommendations are challenging to adhere to for the majority of the patients [2]. Also, there is limited evidence that adherence to such advice prevents clinical complications [23]. Hence, studying the whole dietary patterns rather than single nutrient or food group restrictions is an emerging trend for CKD/ESRD patient diets [2] [24-26]. This is also easier to adhere to. There are several studies on analyzing the relation between dietary patterns and clinical outcomes for CKD patients [3, 4, 5, 6, 7, 8, 9, 26].

Chen et al [3] studied the association between plant protein intake and mortality in CKD. In the study higher plant protein ratio was found to cause lower mortality for CKD patients in stage 3 or higher ($\text{eGFR} < 60 \text{ ml/min/1.73 m}^2$) though not for others (stage 1 and 2) [3]. This study primarily used statistical methods and Regression Models such as Cox regression models to find the association [3]. Hao-Wen et al [26] studied the association between vegetarian diets and CKD. The study found that vegetarian diets including vegan and ovo-lacto vegetarian diets were possible protective factors. The study utilized The multivariable logistic regression analysis [26].

Gutiérrez et al [4] studied 5 empirically derived dietary patterns such as "convenience" (Chinese and Mexican foods, pizza, and other mixed dishes), "plant-based" (fruits and vegetables), "sweets/fats" (sugary foods), "Southern" (fried foods, organ meats, and sweetened beverages), and "alcohol/salads" (alcohol, green-leafy vegetables, and salad dressing) [4]. The study found that dietary pattern rich in processed and fried foods was associated with higher mortality in persons with CKD. On the other hand, a diet rich in fruits and vegetables was found to be protective [4].

Huang et al. [5] studied whether Mediterranean diet can preserve kidney function along with maintaining favorable cardiometabolic profile with reduced mortality risk for individuals with CKD. The study found that adhering to Mediterranean diet has a lower likelihood of having CKD in elderly men. The study also found that a greater adherence to this diet can improve survival for CKD patients [5]. Huang et al [5] in the above study, used unpaired t test, nonparametric Mann–Whitney test, or chi-square test as appropriate for Comparisons between CKD and non-CKD men. To evaluate the association of Mediterranean diet with the presence of CKD, Crude and multiple adjusted logistic regression models were fitted. All tests were two-tailed, and $P < 0.05$ was considered significant [5]. One aspect of Muntner et al [6] study was to find out how Life's Simple 7 factors (Smoke, Activity, BMI, Diet, Blood Pressure, Cholesterol, and Glucose) affect in getting ESRD. For each factor, participants were assigned to one of three levels of compliance to the recommendations. The three levels are ideal, intermediate, and poor. The study shows that people who have ideal scores in more of these factors have lower likelihood of getting ESRD. This study utilized Cox proportional hazards models. Adjustment were made for age, race, sex, stroke-based geographic region of residence, income, education, and history of stroke or coronary heart disease [6].

Ricardo et al [7] studied the association of death to healthy lifestyles especially in relation to CKD. The

study found that adherence to healthy lifestyles was associated with lower risk for mortality in CKD patients. In this study, to determine the association between a healthy lifestyle and survival among individuals with CKD, Cox proportional hazards models were used while also adjusting for important covariates. Stratified survival analyses by eGFR and UACR was performed for Sensitivity analyses [7]. Suruya et al. [8] studied dietary patterns in hemodialysis patients in Japan and researched associations between dietary patterns and clinical outcomes. The study found that patients with unbalanced diet were more likely to have adverse clinical outcomes. Hence, such patients when in addition to portion control, maintains a well-balanced diet especially for the food groups meat, fish, and vegetables will have less adverse clinical outcomes [8]. Suruya et al [8] utilized a principal components analysis (PCA) with Promax rotation to reduce to a smaller set of food groups for analysis. PCA was used to find food groups eaten with equal frequencies [8]. Cox regression model was used for the analysis with multiple models where each model had a different combination of covariants [8].

Another study by Ricardo et al [9] estimated the degree of adherence to a healthy lifestyle that decreases the risk of renal and cardiovascular events among adults with chronic kidney disease (CKD). The study found that adherence to a healthy lifestyle was associated with lower all-cause mortality risk in CKD. The greatest reduction in all-cause mortality was related to nonsmoking [9]. This study by Ricardo et al [9], to compare categorical and continuous variables used Chi-squared and analysis of variance tests respectively. To examine the association between healthy lifestyle and outcomes, Cox proportional hazards models were used. Death was treated as a censoring event. Three nested Cox proportional hazards models were fitted and were adjusted sequentially for potential explanatory variables [9]. G. Asghari et al [27] studied the association of population-based dietary pattern with the risk of incident CKD. The study concluded that high fat and high sugar diet pattern is associated with significantly increased (46%) odds of incident CKD where a lacto-vegetarian diet can be protective of CKD by 43%. The study utilized multivariable logistic regression to calculate odds ratio for the association.

Overall, most of the reviewed studies primarily used direct clinical data of patients for several years and applied statistical analysis primarily. This research primarily utilized public datasets from CDC, USRDS. One of the studies above utilized the dietary pattern data from CDC and NHANES like this study. However, this study will differ in the methodology, exploration, and analysis. This study is finding relations between datasets from multiple sources and is focused on finding patterns and relations in general population than specific/selected individuals. Most of the studies above utilized primarily statistical methods and sensitivity analysis where primarily regression models especially Cox regression models were used. In a couple of cases, Principal Component Analysis (PCA) was used. This research primarily utilized public datasets from CDC, USRDS, Health.gov. For association and prediction, PCA, Regression, and several machine learning approaches are utilized. For food groups and subgroups, we utilized the USDA categorization. Recommended amounts for food groups provided by CDC/Health.gov is used. Additionally, ACR values are predicted based on datasets by CDC such as NHANES dietary intake survey, and laboratory data on ACR. For the prediction, machine learning approaches are used. The machine learning approaches used for ACR value prediction are: Regression and Bayesian.

3 Exploratory Study

Initially Datasets from USRDS on CKD and ESRD patients were explored for Patient characteristics, prevalence of CKD, dietary intake, dietician care, mortality, and survival. Dietary intake data were missing in these datasets, although dietician care data, mortality and survival data by age groups were included. Furthermore, the dataset did not include any linking data for dietician care data, and target mortality/survival data. Hence, datasets from a study on dietary shift recommendation by CDC was explored. These datasets include recommended intake amounts for each age groups including average intake amount (of food groups and food subgroups) by age groups were provided. However, the dietary shift recommendation of the CDC datasets and USRDS mortality i.e. target variable data were not aligned, the dietary intake data from NHANES survey were explored. Hence, NHANES survey data were regrouped to reflect the USRDS age groups, while the USDA codes were used for Food groups/subgroups/intake food for NHANES survey. Finally the USRDS data, combined with the shift recommendation dataset along with additional data from CDC and other sources were explored to create food group recommended amounts for matching age groups. An initial exploratory analysis was performed on the combined datasets using univariate analysis, bivariate analysis, regression, and factor analysis. The results of these explorations are given in Figure 1.

Table 1: **Regression Output from Exploratory Analysis using Excel (Data Analysis Module)**

Metrics	Food Groups	Food Subgroups
Multiple R	0.880156954	0.999378722
R Square	0.774676264	0.99875783
Adjusted R Square	0.616949648	0.978883104
Standard Error	6.223447127	1.461167293

4 Methodology

The research methodology is summarized by the diagram given in Figure 2. The primary purpose of this research is to assess the effect of dietary patterns on the ACR levels, survival and morbidity for chronic kidney disease (CKD)/End Stage Renal Disease (ESRD) patients.

4.1 Methodology Overview


A dataset released by CDC/Health.gov with the dietary habits and ACR values for 10,000 individuals are studied. Also, age group based mortality and survival of CKD/ESRD patients provided by USRDS are studied. Afterwards, utilized Principal Component Analysis to identify the most important food groups and subgroups affecting the ACR value and the Mortality/Survival. Statistical regression and factor analysis was then applied to understand the correlation between ACR, and Mortality/Survival to dietary patterns. Machine learning approaches including Regression, Polynomial Regression, and

		Coefficients	Standard Error	t Stat	P-value	Lower 95%	Upper 95%	Lower 95.0%	Upper 95.0%
	Intercept	44.479	95.829	0.464	0.652	-169.041	257.999	-169.041	257.999
Actual Vegetable Intake	X Variable 1	0.191	0.165	1.155	0.275	-0.177	0.559	-0.177	0.559
Actual Fruit intakes	X Variable 5	0.031	0.169	0.183	0.858	-0.345	0.407	-0.345	0.407
Avg Fats oils and salad dressings take	X Variable 7	0.012	0.978	0.013	0.990	-2.168	2.192	-2.168	2.192
Actual Taken Sugars sweets and beve	X Variable 6	-0.015	0.013	-1.118	0.290	-0.045	0.015	-0.045	0.015
Actual Protein Intake	X Variable 2	-0.023	0.154	-0.152	0.882	-0.367	0.320	-0.367	0.320
Actual Dairy Intake	X Variable 4	-0.085	0.099	-0.852	0.414	-0.306	0.137	-0.306	0.137
Actual Grain Intake	X Variable 3	-0.086	0.066	-1.298	0.223	-0.233	0.062	-0.233	0.062


Regression on Actual Food Group Intake

		Coefficients	Standard Error	t Stat	P-value	Lower 95%	Upper 95%	Lower 95.0%	Upper 95.0%
	Intercept	1.372	54.715	0.025	0.984	-693.852	696.595	-693.852	696.595
Actual Starchy vegetables Intake	X Variable 3	0.310	0.135	2.299	0.261	-1.404	2.024	-1.404	2.024
Actual Other vegetables Intake	X Variable 4	0.159	0.069	2.287	0.262	-0.722	1.039	-0.722	1.039
Avg Meat, Poultry and Eggs subgroup taken	X Variable 7	0.144	0.111	1.304	0.416	-1.263	1.552	-1.263	1.552
Actual Dark-green vegetables Intake	X Variable 1	0.098	0.121	0.807	0.568	-1.443	1.639	-1.443	1.639
Avg Added Sugars/Sugars and sweets taken	X Variable 10	0.047	0.091	0.520	0.695	-1.110	1.205	-1.110	1.205
Actual Whole grains intakes	X Variable 5	0.036	0.114	0.312	0.807	-1.416	1.488	-1.416	1.488
Actual Taken Refined grains amount	X Variable 6	0.027	0.108	0.255	0.841	-1.339	1.394	-1.339	1.394
Avg Nonalcoholic beverages taken	X Variable 16	0.014	0.010	1.414	0.392	-0.115	0.143	-0.115	0.143
Avg Milks and milk drinks taken	X Variable 13	-0.001	0.079	-0.008	0.995	-1.009	1.008	-1.009	1.008
Avg Alcoholic beverages intake	X Variable 15	-0.016	0.008	-1.949	0.302	-0.118	0.087	-0.118	0.087
Avg Water, noncarbonated intake	X Variable 14	-0.032	0.014	-2.288	0.262	-0.211	0.147	-0.211	0.147
Avg Seafood taken	X Variable 8	-0.052	0.071	-0.742	0.594	-0.951	0.846	-0.951	0.846
Avg Nuts, Seeds, and Soy Products taken	X Variable 9	-0.153	0.068	-2.245	0.267	-1.016	0.711	-1.016	0.711
Actual Red and orange vegetables Intake	X Variable 2	-0.245	0.075	-3.248	0.190	-1.202	0.712	-1.202	0.712
Avg Solid Fats taken	X Variable 12	-0.583	0.389	-1.497	0.375	-5.528	4.363	-5.528	4.363
Avg Oils taken	X Variable 11	-0.594	0.668	-0.889	0.538	-9.087	7.899	-9.087	7.899

Regression on Actual Food Subgroup Intake



Line Plot for Regression outcome



Heatmap for exploration

Figure 1: Exploratory Analysis and Representative Output




Figure 2: Methodology in a Diagram

Bayesian with or without 10 fold cross validations are applied on the datasets to understand if dietary patterns can be used to predict ACR values and mortality.

In addition to studying the association of CKD mortality and ACR values with ratios from the recommended high, we have explored the potential impact of diet recommendation shift [11], on the incidence of CKD in the general population.

4.2 Study Selection

For dietary patterns, CKD measures (such as Albumin to Creatinine Ratio - ACR), and Kidney condition measures, a dataset from the National Health and Nutrition Examination Survey on dietary habits conducted by the CDC [10] was used. The survey has data from 1996 to 2016 [10]. This study primarily utilized data for 2015-2016. The survey recorded 24 hours individual food item intake amount. Two surveys were taken within 3 to 10 days apart. Each survey provided food item intake amounts in a day and recorded the diet style as well as diet-restrictions. Individual food items are represented using USDA food code. The survey also provided total nutrients data. CDC also released examination, laboratory, demographics, and other related data for those participants. This study explored and utilized examination data such as Kidney Condition data, laboratory data such as ACR data & Blood Pressure data, and demographics data for age.

For mortality and survival information, dataset from the United States Renal Data System (USRDS) on CKD and ESRD [16, 17] were utilized. “USRDS investigates the transition of care from CKD to ESRD and end-of-life care for those with advanced kidney disease” [19]. USRDS also releases data on the Incidence, Prevalence, Patient Characteristics, and Treatment Modalities on CKD, and ESRD patients. USRDS reports survival and mortality using metrics such as Mortality rates, Total Mortality Count, 90 day survival for dialysis and/or transplantation patients, 10 year survival for dialysis and/or transplantation patients, Average Expected remaining lifetime with or without pre-condition and treatment options used. The data are either aggregated or patient specific detail data. However, only aggregated data are public where patient specific data access requires special request and permission. This research utilized only the public dataset i.e. age-group based aggregated data. In couple of experiments, aggregated age-group based data from USRDS was mapped to specific age for NHANES data for each age and participants.

The dietary survey data (NHANES) represented the food items taken by the participants using USDA food codes [14, 12, 13]. Hence, USDA food codes [14, 12, 13] are used to assign food groups and subgroups to the NHANES [10] survey data to properly group/subgroup the dietary intake of the participants with some customizations.

4.3 Data Synthesis

NHANES survey data as provided for two days are averaged to get the intake amount for one day. Both individual food item data and nutrients intake data are averaged. USDA food codes are used to map food items to food groups and subgroups.

ACR and Kidney condition data for each individual are merged with the averaged food groups, subgroups, and nutrients data. This data was further complemented with the food group recommendations data from health.gov, and mortality rate by age from the USRDS. ACR values and mortality rate are used as the target variables.

PCA was applied to find out important food groups and subgroups while regression was applied to find potential associations with ACR and Mortality. Linear Regression, Polynomial Regression, Random Forest Regression, Bayesian prediction with or without 10 fold cross validations were applied to study the predictability of ACR Values and Mortality in the test dataset.

In another mortality experiment, the above synthesized datasets were aggregated for USRDS age groups to calculate average food group/subgroup intake by age groups.

5 Experiment Design

Experiments as provided in Tables 2 to 10 are designed to find associations between dietary patterns and CKD mortality as well as to predict ACR values. All of these experiments except set 8 and 9 are conducted.

Table 2: **Set 1: Association of Food Groups with CKD Mortality**

Primary Input Dataset:	NHANES dietary intake survey aggregated by USRDS age groups to calculate average food item intake by the participants
Target Variable:	ESRD: Avg. Annual Mortality rates
Experiment 1.1:	Identify contributing and important food groups in the dataset using PCA a) Using Actual Intake Amount b) Using ratios of intake and recommended high amounts
Experiment 1.2:	Find correlation (using Pearson’s correlation and regression) between CKD mortality and important food groups as found using PCA in experiment 1.1. a) Using Actual Intake Amount b) Using ratios of intake and recommended high amounts

Table 3: **Set 2: Association of Food Subgroups with CKD Mortality**

Primary Input Dataset:	NHANES survey aggregated by USRDS age groups to calculate average food item (food subgroups) intake by the participants
Target Variable:	ESRD: Avg. Annual Mortality rates
Experiment 2.1:	Identify important food sub groups in the dataset using PCA and Actual Average Intake Amount
Experiment 2.2:	Similar to experiment 1.2 (Regression); however, used food subgroups and actual average intake amount only

Table 4: **Set 3: Effect of Food Groups on ACR**

Primary Input Dataset:	NHANES survey data for each participant; intake amounts for food groups were averaged for two surveys. This data was merged with laboratory tests for ACR
Target Variable:	Albumin to Creatinine Ratio (ACR)
Experiment 3.1:	Identify contributing and important food groups in the input dataset using PCA. This is different than Experiment 1 because entire survey is being used here; not the aggregated data by age groups
Experiment 3.2:	Find out correlation (using Pearson's correlation, and regression) between ACR Values and important food groups as found using PCA in experiment 3.1.

Table 5: **Set 4: Effect of Nutrients on ACR**

Utilize the same experiments as done for ACR and Food Groups. However, use nutrients intake a) with or b) without combining with food groups	
Experiment 4.1:	PCA to identify contributing factors
Experiment 4.2:	Regression to find correlations among factors found in experiment 4.1

Table 6: **Set 5: Effect of Food Subgroups on ACR**

Primary Input Dataset:	NHANES survey data for each participant; intake amounts for food subgroups were averaged for two surveys. This data was merged with laboratory tests for ACR
Target Variable:	Albumin to Creatinine Ratio (ACR)
Similar experiments like set 3 and set 4. However, used food subgroups as the input/source variables	

Table 7: **Set 6: Use Regression and Bayesian to predict ACR using Food Subgroups intake**

Utilize Machine Learning Approaches for ACR prediction in test dataset.	
Primary Input Dataset:	Input dataset from Set 5 is used here as the input dataset
Target Variable:	ACR Values; Also, ACR categories (CKD or Not)
Experiment 6.1:	ACR value prediction using linear regression. (ACR category was also an option)
Experiment 6.2:	Conduct experiment 6.1; Use 10 folds cross validation where applicable.
Goal:	Find ACR predictability in the test dataset such as calculate R2 Score or generate Confusion Matrix
Experiment 6.3:	Conduct experiment 6.1; however use Polynomial Regression
Experiment 6.4:	Conduct experiment 6.3; With 10 Folds Cross Validations

Experiment 6.5:	Conduct experiment 6.1; however, use a) Random Forest Regression with or without 10 Folds Cross Validations. b) Utilize Polynomial Regression in the process.
Experiment 6.6:	Conduct experiment 6.1; however, use a) Bayesian prediction with or without 10 Folds Cross Validations. b) Use Polynomial Fit

Table 8: **Set 7: Find effect on CKD Mortality using survey data with no data aggregation by Age Groups**

Input Data:	Bring CKD mortality data to each participant using the corresponding ages
Use PCA (to find contributing food groups and subgroups) and then Regression to find correlation between mortality and Food Groups/Subgroups/ACR values	

Table 9: **Set 8: Utilize Regression and Bayesian to predict Mortality using survey data with no data aggregation by Age Groups**

Input Data:	Bring CKD mortality data to each participant using the corresponding age i.e. Input dataset from Set 7 can be used here as the Input dataset
And then utilize Machine Learning Approaches including Regression and Bayesian for Mortality Prediction on Test Dataset.	

Table 10: **Set 9: Association between Food Groups/Subgroups and Remaining Life for CKD Patients: Use not aggregated dietary intake data**

Input Data:	Bring remaining life data such as 5 years survival probabilities for each participant using the corresponding age and CKD status
Use PCA (to find contributing food groups and subgroups) and then utilize Regression to find correlation between remaining life probabilities and Food Groups/Subgroups	

6 Results

Associations of Food Groups, Food Subgroups, Food Nutrients with CKD mortality as discovered by the experiments above are provided in this section. Outcome of ACR value prediction in the test data are also provided.

6.1 Food Groups, Food Subgroups, Food Nutrients and CKD Mortality

Food Groups and Mortality

Experiments (Set 1, Table 2) with aggregated NHANES and USRDS data to find associations between food groups and CKD mortality using PCA and Regression show that Grains (-0.84) and Fruits (-0.43) have negative correlations with CKD mortality i.e. mortality is high for the patients who took significantly lower amount of Grains and Fruits than recommended amounts. Data exploration (Figures 3 and 4) also reflects the negative relation. As the correlation for fruits is -0.43 i.e. not very high, hence, Fruits can be thought of mildly/moderately associated.




Figure 3: **Food Groups (Grains) and Mortality - Negative Correlations**




Figure 4: **Food Groups (Fruits) and Mortality - Negative Correlations**

Vegetables show positive (0.58) correlation i.e. mortality is high for the patients who took more vegetables. The correlation of 0.58 does not provide a very strong conclusion. Data shows (Figure 5) such correlations in older adults. Even though ratios of food intake amount to high end of recommended amounts (actual intake amounts also show positive relation) were used; age might have biased the correlation. This does not show conformity with the general recommendation to take more vegetables for CKD patients. However, as experiments with food subgroups show that vegetable subgroups such as Other vegetables (0.68), Red and Orange vegetables (0.55), and Starchy vegetables (0.44) show positive




Figure 5: **Food Groups and Mortality - Positive Correlations (Vegetables)**

correlations that have an impact on the Vegetable group correlation. Although more vegetable intake is a general recommendation for CKD patients, certain vegetables with more carbohydrates (and sugars) such as Starchy Vegetables as well as Vegetables with more Potassium and Calcium such as Tomatoes, and Spinach are recommended to be taken at a lower amount. For diabetes induced CKD, starchy vegetables are not highly recommended in general. Considering the subgroups, the moderate positive correlation as this study found for vegetables food group is in conformity with the general recommendations for CKD patients.

Experiments (Set 2, Table 3) with aggregated NHANES and USRDS data to find associations between food subgroups and CKD mortality using PCA and Regression show that Other vegetables (0.68), Red and orange vegetables (0.55), and Starchy vegetables (0.44) have positive correlations with mortality i.e. mortality is low when the intake amounts are low, and mortality is high when intake amounts are high. Data Exploration plots (Figure 6) also show these positive relations.

Food subgroups such as Alcoholic Beverages (-0.79), Added Sugars/Sugars and Sweets (-0.64), Whole Grains (-0.61), and 'Nuts, Seeds, and Soy Products' (-0.55) show the most negative correlations with CKD mortality. Data exploration (Figure 7) also shows negative correlations as shown in the charts below. These outcomes are also consistent with current knowledge except for Sugars. Prevalence of Stage 3 CKD is lower in Alcohol Drinkers than non-drinkers [2, 49], Nuts being Phosphorus rich and Whole Grains being Potassium rich are detrimental to CKD patients and can cause higher mortality when taken in higher quantities.

Mortality Study with Non-aggregated Data

For experiments (Set 7, Table 8) where mortality rates based on ages were used for each NHANES survey row (i.e. not aggregated), the following food subgroups show more positive correlations than others: Fats, Eggs, Other vegetables, Nonalcoholic beverages, White potatoes and Puerto Rican starchy vegetables, Tomatoes and tomato mixtures, Oils, Deep-yellow vegetables respectively. In the study, the following subgroups showed more negative correlations than others: Grain mixtures, Frozen plate meals, Soups, Crackers and Salty snacks from grain products, Milks and milk drinks, Sandwiches with Meat, Poultry, and fish, Poultry. Although the correlation numbers in this study are very low, the positive and




Figure 6: Food Subgroups and Mortality - Positive Correlations




Figure 7: Food Subgroups and Mortality - Negative Correlations

negative correlations are consistent with current knowledge [48, 49, 50, 51, 52, 53, 54, 55, 56, 57].

Food Groups, Food Nutrients, and Albumin to Creatinine Ratio (ACR) Association

The experiments (Set 3, 4, and 5 i.e. Table 4, 5, 6) using PCA and Regression showed negligible correlation between ACR and food groups and subgroups intake. However, some food groups and/or nutrients such as Dairy, and ‘Sugars, Sweets, and Beverages’ have higher and positive though negligible (0.02) effect than the others where Fruits (-0.01) showed negative effect. For nutrients, Polyunsaturated fatty acids (-0.02), and Iron (-0.02) have negative correlations where Choline (0.02) showed more positive correlation than others. Findings for Choline matches with medical knowledge [48]. As the correlations are not significant further analysis can be done on the data especially for food groups and nutrients that are found important (using PCA) in the data as provided below:

Dairy, Fats, oils, and salad dressings’, Fruits, Grains, Protein, Sugars, sweets, and beverages’, Vegetables, Avg energy kcal, avg protein gm, avg carbohydrate gm, avg total fat gm, avg total saturated fatty acids gm, avg total monounsaturated fatty acids gm, avg total polyunsaturated fatty acids gm, avg lutein

zeaxanthin mcg, avg thiamin vitamin B1 mg, avg riboflavin Vitamin B2 mg, avg Niacin mg, avg Calcium mg, avg Phosphorus mg, avg Magnesium mg, avg Iron mg, avg Zinc mg, avg Copper mg, avg Sodium mg, avg Potassium mg, avg Selenium mcg, Hexadecenoic gm, Octadecenoic gm

6.2 Food Subgroups and Albumin to Creatinine Ratio (ACR) Association

The experiments showed ‘Milk desserts, Sauces, Gravies’ (0.22), and Alcoholic Beverages (0.087) have more positive correlations with ACR than the other food subgroups i.e. taking more of these food subgroups results higher ACR values. Research by Uehara et al. [55] also shows that excessive Alcohol consumption can cause Proteinuria/Albuminuria (high ACR). Nettleton et al. [56] found that high fat dairy can be linked to high ACR values where low-fat dairy is not strongly linked to high ACR values. However, the correlation as this research found is very low. Low values might still explain a correlation where ACR values might depend on other factors in together than only these food subgroups. Fruits and juicy baby foods show negative correlation (-0.04) though not significant i.e. taking high amount does not increase ACR values that matched with current knowledge [57].

6.3 Predictability of ACR values based on Dietary patterns

Experiments (Set 6, Table 7) were conducted to predict ACR values from the dietary intake patterns data. Machine Learning (ML) Approaches of Regression, Polynomial Regression, Random Forest Regression, Bayesian prediction with or without 10 fold cross validations were applied on Food Subgroups intake dataset. Only the food subgroups that were found to be important using PCA were used for the ML approaches.

Target Variables

Absolute ACR values and ACR Categories were used as the target variables. For ACR category, ACR <30 is assigned to class 0 (i.e. no ckd), and ACR >30 is assigned to class 1 (CKD). ACR values less than 30 indicate no CKD, where ACR values between 30 and 300 indicate moderate CKD. ACR >300 is considered severe CKD.

Outcome when ACR Values are used as the Target

The best test set accuracies were found using approaches such as: 10 Fold Cross Validations with Polynomial Regression (95%), Polynomial Bayesian with 10 fold Cross Validations (68%), Polynomial Regression (57%), Bayesian on Polynomial fit (41%), Cross Validation with Polynomial Random Forest Regression (21%). A list of the best performing approaches and the outcome are provided in Table 11 below.

Outcome when ACR Category is Used as the Target

After regression, $y > 0.5$ is assigned to category 1 (CKD), others were assigned to category 0. Test accuracies are 88%. Format for confusion matrix used in the Table 12 below is: (Total, % Correct : [TP, FN, FP, TN]). The high prediction accuracies might relate to the fact that only 10 to 13% of the population has ACR >30. However, as cross validations also show high accuracies, it can be concluded

Table 11: **Outcome when ACR Values are used as the Target**

Data Normalized	Target	Approach	MSE Train	MSE Test	RMSE Train	RMSE Test	R2 Score Train	Accuracy: Test R2 Score if not mentioned
No	ACR Value	10 Fold Cross Validation Polynomial Regression			1	2	3	-0.957 cross val score
No	ACR Value	Polynomial Bayesian with Cross Validation			1	2	3	-0.682 cross val score
No	ACR Value	Polynomial Regression	90965	52946	301	301	0.359	-0.579 r2 score on test data
No	ACR Value	Bayesian on Polynomial fit	93047	47431	305	305	0.344	-0.414 r2 score on test data

Table 12: **Outcome when ACR Category is used as the Target**

Data Normalized	Target	Approach	Train Confusion Matrix	Test Confusion Matrix
No	Category	Linear Regression	6927, 87%: [6032,0, 895,0]	770, 88% (692, 0, 78, 0)
Yes	Category	Linear Regression	6927, 87%: [6032,0, 895,0]	770, 88% (692, 0, 78, 0)

that ACR values can be well predicted using Machine Learning approaches especially with 10 Fold Cross Validation Polynomial Regression having accuracy 95%.

7 Conclusions

Chronic Kidney Disease (CKD) leading to End Stage Renal Disease (ESRD) is very prevalent today; treatment facilities for dialysis, and donors for organ transplantation are limited. Consequently, many patients die waiting for proper treatment (recent news on USA, Bloomberg). Majority of the studies focused on drugs to control CKD progression where some studies focused on diets, Nutrients, and Food Items. Controlling CKD using changes to dietary patterns can be beneficial to both the patients and the economy. Hence, this study focused on the effect of dietary patterns on CKD mortality as well as on a

CKD measure named Albumin to Creatinine Ratio (ACR). PCA and Regression are used to study the association between ACR and CKD mortality/survival. Additionally, regression models are trained to predict ACR values from dietary patterns. Vegetables such as Other Vegetables, Starchy Vegetables, and Red and Orange Vegetables showed moderate positive correlations with mortality whereas Grains, Fruits, Alcohol, Sugar, and Nuts showed negative correlations. ACR values were not found strongly correlated with dietary patterns. Overall, the results of this research matched the findings of other studies and current knowledge with few exceptions. Machine Learning approaches showed that ACR values could be predicted in the test dataset with high accuracy; the best performing approach was 10 Fold Cross Validation for Polynomial Regression (95%).

8 Appendix

[Github Link Root](#)

[Github link for Python and SQL Code](#)

9 References

- [1] The National Institute of Diabetes and Digestive and Kidney Diseases. What Is Chronic Kidney Disease? <https://www.niddk.nih.gov/health-information/kidney-disease/chronic-kidney-disease-ckd/what-is-chronic-kidney-disease>
- [2] Jaimon T. K., Suetonia C. P., Shu N. W., Marinella R., Juan-Jesus C., Katrina L. C., and Giovanni F. M. S. Healthy Dietary Patterns and Risk of Mortality and ESRD in CKD: A Meta-Analysis of Cohort Studies
- [3] Chen X, Wei G, Jalili T, Metos J, Giri A, Cho ME, Boucher R, Greene T, Beddhu S: The associations of plant protein intake with all-cause mortality in CKD. *Am J Kidney Dis* 67: 423–430, 2016 (26)
- [4] Gutierrez O.M., Muntner P, Rizk D.V., McClellan W.M., Warnock D.G., Newby P.K., Judd S.E.: Dietary patterns and risk of death and progression to ESRD in individuals with CKD: A cohort study. *Am J Kidney Dis* 64: 204–213, 2014 (27)
- [5] Huang X., Jime nez-Moleo J. J., Lindholm B., Cederholm T., Arnoldv J., Rise rus U., Sjogren P., Carrero J. J.: Mediterranean diet, kidney function, and mortality in men with CKD. *Clin J Am Soc Nephrol* 8: 1548–1555, 2013 (28)
- [6] Muntner P, Judd S.E., Gao L, Gutierrez O.M., Rizk D.V., McClellanW., Cushman M., Warnock D.G.: Cardiovascular risk factors in CKD associated with both ESRD and mortality. *J Am Soc Nephrol* 24: 1159–1165, 2013 (29)
- [7] Ricardo A.C., Madero M., Yang W., Anderson C., Menezes M., Fischer M.J., Turyk M., Daviglus M.L., Lash J.P.: Adherence to a healthy lifestyle and all-cause mortality in CKD. *Clin J Am Soc Nephrol* 8: 602–609, 2013 (30)
- [8] Tsuruya K., Fukuma S., Wakita T., Ninomiya T., Nagata M., Yoshida H., Fujimi S., Kiyohara Y.,

- Kitazono T., Uchida K., Shirota T., Akizawa T., Akiba T., Saito A., Fukuhara S.: Dietary patterns and clinical outcomes in hemodialysis patients in Japan: A cohort study. PLoS One 10: e0116677, 2015 (31)
- [9] Ricardo A.C., Anderson C.A., Yang W., Zhang X., Fischer M. J., Dember L. M., Fink J. C., Frydrych A., Jensvold N. G., Lustigova E., Nessel L. C., Porter A. C., Rahman M., Wright Nunes J. A., Daviglius M. L., Lash J. P.; CRIC Study Investigators: Healthy lifestyle and risk of kidney disease progression, atherosclerotic events, and death in CKD: Findings from the Chronic Renal Insufficiency Cohort (CRIC) Study. Am J Kidney Dis 65: 412–424, 2015 (17)
- [10] Centers for Disease Control and Prevention. National Health and Nutrition Examination Survey. <https://wwwn.cdc.gov/nchs/nhanes/search/datapage.aspx?Component=Dietary>.
- [11] Health.gov. Shifts Needed to Align With Healthy Eating Patterns. <https://health.gov/dietaryguidelines/2015/guidelines/chapter-2/a-closer-look-at-current-intakes-and-recommended-shifts/>
- [12] Agricultural Research Service (ARS), USDA. Food Code Numbers and the Food Coding Scheme. https://readdir.arsnet.usda.gov/codesearchwebapp/gcp3kq55ssdyc445ry2k2rus/coding_scheme.pdf
- [13] The U.S. Department of Agriculture's (USDA). Vegetable Subgroups. <http://www.cn.nysed.gov/common/cn/files/Vegetable%20Subgroups.pdf>
- [14] Centers for Disease Control and Prevention. Key Concepts About the USDA Food Coding Scheme. <https://www.cdc.gov/nchs/tutorials/Dietary/SurveyOrientation/ResourceDietaryAnalysis/Info2.htm>
- [15] Health.gov. Appendix 3. USDA Food Patterns: Healthy U.S.-Style Eating Pattern. <https://health.gov/dietaryguidelines/2015/guidelines/appendix-3/>
- [16] United states Renal Data System (USRDS). The 2018 USRDS Annual Data Report Reference Tables : <https://www.usrds.org/reference.aspx>
- [17] United states Renal Data System (USRDS). 2018 ADR Chapters. <https://www.usrds.org/2018/view/Default.aspx>
- [18] ARS, USDA. Food Surveys Research Group: Beltsville, MD . Documentation and Dataset: <https://www.ars.usda.gov/northeast-area/beltsville-md-bhnrc/beltsville-human-nutrition-research-center/food-surveys-research-group/docs/wweia-documentation-and-data-setsY>
- [19] United states Renal Data System (USRDS). 2018 USRDS Annual Data Report: Executive Summary. https://www.usrds.org/2018/download/v1_00_ExecSummary_18.pdf
- [20] Health.gov, CDC. Dietary Guidelines for Americans 2015-2020. https://health.gov/dietaryguidelines/2015/resources/2015-2020_dietary_guidelines.pdf
- [21] Health.gov, CDC. Dietary Guidelines for Americans. <https://health.gov/dietaryguidelines/dga95/9DIETGUI.HTM>
- [22] Mortality and Causes of Death. https://www.usrds.org/2018/ref/ESRD_Ref_H_Mortality_2018.xlsx
- [23] Tong A, Chando S, Crowe S, Manns B, Winkelmayer WC, Hemmelgarn B, Craig JC: Research priority setting in kidney disease: A systematic review. Am J Kidney Dis 65: 674–683, 2015
- [24] Lin J, Fung TT, Hu FB, Curhan GC: Association of dietary patterns with albuminuria and kidney function decline in older white women: A subgroup analysis from the Nurses' Health Study. Am J Kidney Dis 57: 245–254, 2011

- [25] Taylor EN, Fung TT, Curhan GC: DASH-style diet associates with reduced risk for kidney stones. *J Am Soc Nephrol* 20: 2253–2259, 2009
- [26] Liu, Hao-Wen; Tsai, Wen-Hsin; Liu, Jia-Sin; Kuo, Ko-Lin. 2019. "Association of Vegetarian Diet with Chronic Kidney Disease." *Nutrients* 11, no. 2: 279.
- [27] Golaleh Asghari, Mehrnaz Momenan, Emad Yuzbashian, Parvin Mirmiran Email author and Fereidoun Azizi. Dietary pattern and incidence of chronic kidney disease among adults: a population-based study
- [28] Tanushree Banerjee¹, Deidra C. Crews², Delphine S. Tuot³, Meda E. Pavkov⁴, Nilka Rios Burrows⁴, Austin G. Stack⁵, Rajiv Saran^{6,7}, Jennifer Bragg-Gresham⁶ and Neil R. Powe^{1,8}; for the Centers for Disease Control and Prevention Chronic Kidney Disease Surveillance Team⁹ Poor accordance to a DASH dietary pattern is associated with higher risk of ESRD among adults with moderate chronic kidney disease and hypertension
- [29] Jacek R., Beata F., Aleksandra C., Anna G. The Effect of Diet on the Survival of Patients with Chronic Kidney Disease. *Nutrients* 2017, 9(5), 495; <https://doi.org/10.3390/nu9050495>
- [30] National Kidney Foundation. One in Seven American Adults Estimated to Have Chronic Kidney Disease. <https://www.kidney.org/news/one-seven-american-adults-estimated-to-have-chronic-kidney-disease>
- [31] Hannah N. What are the leading causes of death in the US?. <https://www.medicalnewstoday.com/articles/282929.php>
- [32] Davita. Five Stages of CKD. <https://www.davita.com/education/kidney-disease/stages>
- [33] Goal: How to Identify the Most Important Predictor Variables in Regression Models <https://blog.minitab.com/blog/adventures-in-statistics-2/how-to-identify-the-most-important-predictor-variables-in-regression-models>
- [34] How to Interpret Regression Analysis Results: P-values and Coefficients <https://blog.minitab.com/blog/adventures-in-statistics-2/how-to-interpret-regression-analysis-results-p-values-and-coefficients>
- [35] Regression Analysis: How to Interpret the Constant (Y Intercept) <https://blog.minitab.com/blog/adventures-in-statistics-2/regression-analysis-how-to-interpret-the-constant-y-intercept>
- [36] How to Compare Regression Slopes: How to statistically test the difference between regression slopes and constants <https://blog.minitab.com/blog/adventures-in-statistics-2/how-to-compare-regression-lines-between-different-models>
- [37] How Do I Interpret R-squared and Assess the Goodness-of-Fit? <https://blog.minitab.com/blog/adventures-in-statistics-2/regression-analysis-how-do-i-interpret-r-squared-and-assess-the-goodness-of-fit>
- [38] How High Should R-squared Be in Regression Analysis? <https://blog.minitab.com/blog/adventures-in-statistics-2/how-high-should-r-squared-be-in-regression-analysis>
- [39] How to Interpret a Regression Model with Low R-squared and Low P values <https://blog.minitab.com/blog/adventures-in-statistics-2/how-to-interpret-a-regression-model-with-low-r-squared-and-low-p-values>

- [40] Use Adjusted R-Squared and Predicted R-Squared to Include the Correct Number of Variables <https://blog.minitab.com/blog/adventures-in-statistics-2/multiple-regression-analysis-use-adjusted-r-squared-and-predicted-r-squared-to-include-the-correct-number-of-variables>
- [41] How to Interpret S, the Standard Error of the Regression <https://blog.minitab.com/blog/adventures-in-statistics-2/regression-analysis-how-to-interpret-s-the-standard-error-of-the-regression>
- [42] What Is the F-test of Overall Significance in Regression Analysis?. <https://blog.minitab.com/blog/adventures-in-statistics-2/what-is-the-f-test-of-overall-significance-in-regression-analysis>
- [43] Understanding Analysis of Variance (ANOVA) and the F-test. <https://blog.minitab.com/blog/adventures-in-statistics-2/understanding-analysis-of-variance-anova-and-the-f-test>
- [44] How to Compare Regression Slopes. <https://blog.minitab.com/blog/adventures-in-statistics-2/how-to-compare-regression-lines-between-different-models>
- [45] How to Present and Use the Results to Avoid Costly Mistakes, part 1. <https://blog.minitab.com/blog/adventures-in-statistics-2/applied-regression-analysis-how-to-present-and-use-the-results-to-avoid-costly-mistakes-part-1>
- [46] How to Identify the Most Important Predictor Variables in Regression Models. <https://blog.minitab.com/blog/adventures-in-statistics-2/how-to-identify-the-most-important-predictor-variables-in-regression-models>
- [47] How to Interpret your Regression Results <http://sitestree.com/how-to-interpret-your-regression-results>
- [48] Fernandez-Prado R., Esteras R., Perez-Gomez M.V, Gracia-Iguacel C., Gonzalez-Parra E., Sanz A.B., Ortiz A., Sanchez-Nino M.D. Nutrients Turned into Toxins: Microbiota Modulation of Nutrient Properties in Chronic Kidney Disease. *Nutrients*. 2017 May; 9(5): 489. Published online 2017 May 12. doi: 10.3390/nu9050489 PMID: PMC5452219 <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5452219/>
- [49] Hsu, Y. H., Pai, H. C., Chang, Y. M., Liu, W. H., Hsu, C. C. (2013). Alcohol consumption is inversely associated with stage 3 chronic kidney disease in middle-aged Taiwanese men. *BMC nephrology*, 14, 254. doi:10.1186/1471-2369-14-254
- [50] Rippe, J. M., Angelopoulos, T. J. (2016). Relationship between Added Sugars Consumption and Chronic Disease Risk Factors: Current Understanding. *Nutrients*, 8(11), 697. doi:10.3390/nu8110697
- [51] Karalius V.P., Shoham D.A. Dietary sugar and artificial sweetener intake and chronic kidney disease: a review. *Adv Chronic Kidney Dis*. 2013; 20:157–164. doi: 10.1053/j.ackd.2012.12.005
- [52] Jaimon T. Kelly, Suetonia C. Palmer, Shu Ning Wai, Marinella Ruospo, Juan-Jesus Carrero, Katrina L. Campbell and Giovanni F. M. Strippoli Healthy Dietary Patterns and Risk of Mortality and ESRD in CKD: A Meta-Analysis of Cohort Studies. *CJASN* February 2017, 12 (2) 272-279; DOI: <https://doi.org/10.2215/CJN.06190616>
- [53] Jilian k. The 20 Best Foods for People With Kidney Problems. <https://www.healthline.com/nutrition/best-foods-for-kidneys>
- [54] National Kidney Foundation (NKF), USA. Drinking Alcohol Affects Your Kidneys <https://www.kidney.org/news/kidneyCare/winter10/AlcoholAffects>

- [55] Uehara, S., Hayashi, T., Kogawa Sato, K., Kinuhata, S., Shibata, M., Oue, K., Hashimoto, K. (2016). Relationship Between Alcohol Drinking Pattern and Risk of Proteinuria: The Kansai Healthcare Study. *Journal of epidemiology*, 26(9), 464–470. doi:10.2188/jea.JE20150158
- [56] Nettleton, J. A., Steffen, L. M., Palmas, W., Burke, G. L., & Jacobs, D. R., Jr (2008). Associations between micro albuminuria and animal foods, plant foods, and dietary patterns in the Multiethnic Study of Atherosclerosis. *The American journal of clinical nutrition*, 87(6), 1825–1836. doi:10.1093/ajcn/87.6.1825
- [57] Jacobs, D. R., Jr, Gross, M. D., Steffen, L., Steffes, M. W., Yu, X., Svetkey, L. P., Sacks, F. (2009). The effects of dietary patterns on urinary albumin excretion: results of the Dietary Approaches to Stop Hypertension (DASH) Trial. *American journal of kidney diseases : the official journal of the National Kidney Foundation*, 53(4), 638–646. doi:10.1053/j.ajkd.2008.10. 048. <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2676223/>
- [58] National Kidney Foundation. 37 Million American Adults Now Estimated to Have Chronic Kidney Disease. <https://www.kidney.org/news/37-million-american-adults-now-estimated-to-have-chronic-kidney-disease>