

The Allometric Trap: Scaling Laws Predict When Growing Organisms Cannot Afford Their Own Shape

Dr. Sayuj Krishnan S, MBBS, DNB (Neurosurgery)
Consultant Neurosurgeon and Spine Surgeon, Yashoda Hospitals, Malakpet, Hyderabad, India
dr.sayujkrishnan@gmail.com

March 6, 2026

Abstract

Living organisms maintain complex shapes against gravity, but the physical limits of this maintenance are unknown. Here we derive a dimensionless scaling parameter — the Bio-Gravitational Number $\mathcal{B}_g = EI/(MgL^2)$ — that predicts when passive structure becomes insufficient and active metabolic maintenance is required. Cross-species analysis confirms that humans uniquely occupy an “Allometric Trap” ($\mathcal{B}_g \approx 0.01$) where shape maintenance depends entirely on metabolic energy. Using an Information–Elasticity Coupling framework that connects HOX-patterned developmental fields to Cosserat rod mechanics, we show that the spinal S-curve emerges as the energetic ground state, reproducing clinical lordosis ($\sim 42^\circ$) and kyphosis ($\sim 35^\circ$). AlphaFold structural analysis of 23 mechanotransduction proteins reveals a 72% demand–supply anisotropy gap ($p = 0.011$): the proteins required for gravity sensing are thermodynamically expensive, while their metabolic regulators are intrinsically disordered and fragile. This molecular asymmetry creates an Energy Deficit Window during adolescent growth ($L > 0.35$ m) that our model predicts produces scoliosis-like instabilities. These results reclassify Adolescent Idiopathic Scoliosis as a thermodynamic scaling failure and identify molecular targets for metabolic intervention. The human spine maintains a complex S-shaped sagittal profile against gravity, yet its physical principle and adolescent vulnerability remain unexplained. We introduce the Information–Elasticity Coupling (IEC) framework, integrating a developmental information field representing HOX patterning with Cosserat rod mechanics, implemented via PyElastica simulations. The IEC framework selects the S-shaped spinal profile as the energetic ground state, reproducing normal lordosis and kyphosis. Phase diagram analysis reveals gravity-dominated, cooperative, and information-dominated morphological regimes. During simulated adolescent growth, small information-field asymmetries are amplified into scoliotic deformities through a supercritical bifurcation at a critical length, explaining adolescent scoliosis onset. Furthermore, AlphaFold structural analysis of mechanotransduction proteins reveals a demand-supply anisotropy gap quantifying this structural vulnerability. We conclude that spinal geometry emerges as a thermodynamic standing wave maintained by metabolic expenditure. The IEC framework unifies developmental patterning and biomechanics, generating specific testable predictions for scoliosis prevention.

Impact Statement: We identify a universal scaling law governing the metabolic cost of maintaining body shape against gravity, explaining why human spines are uniquely vulnerable to scoliosis during adolescent growth and reclassifying a common disease as a predictable thermodynamic scaling failure.

Model Summary: The Allometric Trap

The Question: What physical law governs how organisms maintain shape against gravity, and why does this maintenance fail during adolescent growth?

The Scaling Law: The Bio-Gravitational Number $\mathcal{B}_g = EI/(MgL^2)$ predicts that humans ($\mathcal{B}_g \approx 0.01$) cannot passively resist gravity and must actively maintain spinal geometry at metabolic cost.

The Mechanism:

- **Metabolic Scaling Mismatch:** The cost of anti-gravitational form maintenance scales as L^4 , while biological supply scales as L^2 , creating a critical Energy Deficit Window at $L > 0.35$ m.
- **Molecular Asymmetry:** AlphaFold analysis reveals that mechanosensory proteins (mean anisotropy 3.08) are structurally expensive, while their metabolic regulators are intrinsically disordered and fragile — a 72% gap ($p = 0.011$).
- **Scoliosis as Scaling Failure:** During adolescent growth, this deficit forces collapse from the active S-curve into a lower-energy scoliotic mode — a predictable thermodynamic transition, not a random genetic error.

1 Introduction

1.1 Background: Gravity, Mechanobiology, and Spinal Morphogenesis

Living organisms routinely maintain complex, non-equilibrium shapes against the relentless pull of gravity. A tree grows vertically despite wind loads; a flamingo stands on one leg; the human spine sustains an intricate S-shaped profile (lordosis and kyphosis) rather than sagging into the passive C-curve predicted by classical beam mechanics [1–3]. The physical principle that governs this “anti-gravitational form maintenance” — and the conditions under which it fails — remains poorly understood.

Spinal morphogenesis is fundamentally driven by mechanobiology, integrating genetic instructions with physical forces to shape tissue architecture [4–6]. From embryonic somitogenesis, which establishes the segmental architecture of the spine via the segmentation clock [7, 8], to the rapid phases of adolescent growth, the spine is continuously shaped by the interplay between intrinsic genetic patterning (e.g., HOX genes [9, 10]) and extrinsic mechanical forces. Tissues actively remodel in response to gravitational loading and cellular tension [11, 12], a process governed by cellular mechanotransducers such as PIEZO channels, integrins, and the YAP/TAZ signaling axis [13–16].

In the intervertebral discs and paraspinal muscles, these mechanosensors continuously integrate mechanical signals to regulate extracellular matrix production and cellular metabolism [17–19]. However, maintaining an upright posture in a gravitational field is energetically expensive. It requires continuous muscular actuation, proprioceptive feedback, and cellular mechanotransduction to oppose gravity and prevent structural collapse [20–23].

1.2 The Bridge: Counter-Curvature as an Organizing Hypothesis

To understand how the spine maintains its shape and why it is uniquely vulnerable during growth, we introduce the concept of “Biological Counter-Curvature” as an organizing hypothesis. For passive structures, the mechanics are straightforward: a beam clamped at one end sags monotonically under its own weight into a gravitational geodesic [2]. The vertebrate spine, however, actively opposes this trajectory, functioning as a **Thermodynamic Standing Wave** [24]. This counter-curved geometry is a dissipative structure sustained by the continuous investment of metabolic free energy [25, 26].

Framing spinal alignment as an active counter-curvature provides a physical basis for its developmental vulnerability. A fundamental scaling crisis emerges during growth. As the spine elongates, the gravitational bending moment scales as $M_g \propto L^4$ [3, 27], while the nutrient supply to the avascular intervertebral disc — constrained by vertebral endplate surface area and diffusion limits — scales only as L^2 [28, 29].

We formalize this divergence through the **Bio-Gravitational Number**, $\mathcal{B}_g = EI/(MgL^2)$, a dimensionless ratio of passive flexural stiffness to gravitational moment. Cross-species analysis reveals that most vertebrates maintain $\mathcal{B}_g > 0.1$ (passively stable), but humans occupy a unique “Allometric Trap” ($\mathcal{B}_g \approx 0.01$) where passive stiffness is entirely insufficient and active metabolic maintenance is required [30].

This scaling mismatch creates a transient **Energy Deficit Window** during the adolescent growth spurt, when the metabolic cost of maintaining the S-shaped counter-curvature ($\propto L^2$) outpaces the biological supply capacity ($\propto L^{0.5}$). Beyond a critical spinal length $L_{\text{crit}} \approx 0.35$ m, the system becomes energetically insolvent. We propose that Adolescent Idiopathic Scoliosis (AIS) — a 3D spinal deformity affecting 3% of children worldwide [31, 32] — is the predictable consequence of this thermodynamic insolvency: a “metabolic buckling” event in which the spine sheds its expensive counter-curvature and collapses into a lower-energy, asymmetric mode, representing a predictable thermodynamic transition rather than a random genetic error.

1.3 Predictions

By modeling the spine through an Information–Elasticity Coupling (IEC) framework that connects developmental patterning to Cosserat rod mechanics, we demonstrate that the spinal S-curve emerges as the energetic ground state. This framework moves beyond correlative observations to a first-principles causal mechanism, generating specific, measurable predictions:

- **Metabolic Rescue of Curve Progression:** Because scoliosis represents an energy deficit, supplementation with mitochondrial cofactors (e.g., Nicotinamide Riboside to boost NAD+) during peak height velocity should measurably delay curve progression or reduce Cobb angle severity in susceptible populations.
- **Anisotropy-Driven Protein Degradation:** Under simulated metabolic stress (e.g., hypoxia), high-anisotropy mechanosensory proteins (such as PIEZO2) will exhibit significantly higher degradation rates compared to low-anisotropy metabolic regulators, measurable via mass spectrometry, precipitating the sensory mismatch that precedes structural deformity.
- **Microgravity-Induced Counter-Curvature Adaptation:** In prolonged microgravity ($g \rightarrow 0$), the relief of the gravitational bending moment will cause the spine to relax toward its information reference state. We predict a measurable reduction in the specific mechanical strain metric $g_{\text{eff}}(s)$, leading to the “Convective Shutdown” of intervertebral discs and subsequent scoliotic adaptations observable in astronauts [29, 33].

2 Theory

We propose that the robust S-shaped geometry of the spine arises from an active counter-curvature mechanism driven by developmental information. We formalize this using an Information–Elasticity Coupling (IEC) framework, where a scalar information field $I(s)$ modifies the effective geometry and energetics of a Cosserat rod. The full mathematical derivations, including Cosserat kinematics, boundary conditions, and the information-geometric formalism, are provided in the Supplementary Information.

2.1 The Bio-Gravitational Number and the Allometric Trap

For a vertebral column of length L , flexural stiffness EI , mass M , and gravitational acceleration g , we define the **Bio-Gravitational Number**:

$$\mathcal{B}_g = \frac{EI}{MgL^2}, \quad (1)$$

a dimensionless ratio of passive flexural resistance to gravitational bending moment. When $\mathcal{B}_g > 0.1$, the organism can resist gravitational collapse through passive stiffness alone. When $\mathcal{B}_g < 0.1$, the organism enters the “Allometric Trap”: it must actively maintain its shape at continuous metabolic cost. Cross-species analysis reveals that small quadrupeds operate well above this threshold, while humans ($\mathcal{B}_g \approx 0.01$) are deep within the trap, relying almost entirely on active metabolic maintenance to prevent spinal collapse. The central problem of vertebrate morphogenesis is the translation of a one-dimensional genetic code into a robust three-dimensional geometry that can withstand environmental loads. In the case of the human spine, this challenge is acute: the column must support the head and trunk against the relentless downward pull of gravity ($g \approx 9.81 \text{ m/s}^2$) while maintaining flexibility for locomotion. Classical structural mechanics predicts that a flexible column clamped at the base and subjected to its own weight will buckle or sag into a monotonic C-shape [2]. Yet, the healthy human spine exhibits a complex, alternating S-shape (lordosis and kyphosis).

This anomaly suggests that the S-shape is not a passive equilibrium but an active *Counter-Curvature*—a geometric standing wave maintained by the continuous expenditure of metabolic energy against the gravitational field. This hypothesis bridges two foundational concepts in biology: the *Morphogenetic Field* of developmental biology [34, 35] and the *Mechanostat* of tissue physiology [11].

The Information Field as a Target Metric. Developmental patterning establishes a “Positional Information” field along the anterior-posterior axis, encoded by the nested expression of HOX genes [9]. We posit that this information field $I(s)$ does not merely specify cell identity but encodes a *target metric*—a preferred intrinsic curvature $\kappa_{rest}(s)$ that the tissue strives to achieve. This is analogous to the “growth field” concepts in plant morphogenesis [36] but applied to a load-bearing animal structure. The spine is thus a “smart beam” that continuously computes the error between its actual geometry (sensed via proprioception) and its genetic target.

Active Maintenance and the Hydrostatic Skeleton. Achieving this target shape against gravity requires work. The spine functions partly as a *hydrostatic skeleton* [37], where the nucleus pulposus of the intervertebral disc acts as a pressurized hydraulic element. The maintenance of this pressure—and thus the maintenance of the S-shape—depends on the balance between osmotic swelling pressure (driven by proteoglycan synthesis) and the mechanical stress of gravity. This active maintenance constitutes a biological implementation of *optimal feedback control* [38], where the system minimizes a cost function comprising both “geometric error” (deviation from the HOX-specified shape) and “metabolic effort” (ATP consumption).

Predictions. This unifying hypothesis generates three testable predictions that distinguish it from purely mechanical buckling models:

1. **Frequency-Hypoxia Coupling:** We predict that the maintenance of the S-shape requires dynamic loading frequencies $> 0.5 \text{ Hz}$ to drive convective transport. Reducing frequency (e.g., microgravity) will stabilize HIF-1 α in the nucleus pulposus before any structural deformity appears [39].

2. **The "Small-Molecule Rescue"**: If scoliotic buckling is a metabolic failure (Energy Deficit), then enhancing the amplitude of the circadian clock (e.g., via Nobiletin) or buffering local pH should restore spinal alignment even without mechanical correction.
3. **The Spinal Phase Shift**: The severity of spinal curvature should correlate with the phase lag between the central (SCN) and peripheral (IVD) circadian clocks, providing a quantifiable biomarker for "Counter-Curvature" failure.

2.2 Rod Geometry and Kinematic Parameterization

The human vertebral column is structurally modeled as an active, continuous Cosserat rod embedded in $\mathbb{R}^3$, an idealization rigorously justified by its high aspect ratio ($L/d \approx 13$) and the dense, segmented distribution of intervertebral joints which act as quasi-continuous elastic hinges [2, 40]. Let the centerline curve be denoted by $\mathbf{r}(s, t) \in \mathbb{R}^3$, where $s \in [0, L]$ is the reference arc length from the sacral base ($s = 0$) to the cranium ($s = L$). To fully capture the macroscopic 3D kinematics, including flexural bending, shear, and torsional deformations, an orthonormal material frame $\{\mathbf{d}_1(s, t), \mathbf{d}_2(s, t), \mathbf{d}_3(s, t)\}$ is attached to each cross-section.

By convention, $\mathbf{d}_3(s, t)$ defines the local tangent vector in the absence of shear strains, while $\mathbf{d}_1(s, t)$ and $\mathbf{d}_2(s, t)$ align with the principal anatomic axes of the vertebral body (e.g., the anteroposterior and mediolateral axes). The spatial evolution of the rod's configuration is thus strictly governed by the kinematic equations:

$$\partial_s \mathbf{r}(s, t) = \mathbf{v}(s, t) = \mathbf{d}_3(s, t) + \boldsymbol{\varepsilon}(s, t), \quad (2)$$

$$\partial_s \mathbf{d}_i(s, t) = \mathbf{u}(s, t) \times \mathbf{d}_i(s, t), \quad (3)$$

where $\boldsymbol{\varepsilon}(s, t)$ captures the shear and axial extension strains. The Darboux vector $\mathbf{u}(s, t) = \kappa_1 \mathbf{d}_1 + \kappa_2 \mathbf{d}_2 + \tau \mathbf{d}_3$ encodes the bending curvatures κ_1, κ_2 and the torsional twist τ . We explicitly impose a clamped boundary condition at the sacrum ($\mathbf{r}(0) = \mathbf{0}, \mathbf{d}_i(0) = \mathbf{e}_i$) reflecting pelvic anchoring, and a free-end condition at the cranium, subject solely to the localized mass of the skull.

2.3 Developmental Information Fields from Morphogenetic Patterning

Rather than treating the spine as a passive beam reacting to gravitational loading, our framework strictly formalizes developmental inputs as a time-dependent, macroscopic scalar information field $I(s, t) \in [0, 1]$. This field acts as a coarse-grained, physical representation of anterior-posterior patterning morphogens (such as the highly conserved HOX gene clusters) and segmentation clock outputs that prescribe regional identity.

Empirically, we map $I(s)$ via a bimodal Gaussian parameterization (Eq. 16) exhibiting distinct maxima at the cervical ($A_c = 0.5, s_c = 0.80$) and lumbar ($A_l = 0.7, s_l = 0.25$) enlargements. These peaks physically correspond to zones of heightened morphogenetic specification required for lordotic posture. The gradient of this morphogenetic field, ∇I , acts as an active "curvature generator." It induces a non-zero rest curvature through a direct geometric coupling parameter χ_κ :

$$\boldsymbol{\kappa}_{\text{rest}}(s) = \boldsymbol{\kappa}_{\text{gen}} + \chi_\kappa \nabla I(s). \quad (4)$$

Furthermore, to quantify the patterning precision, we evaluate the local Shannon entropy $\mathcal{H}[I]$:

$$\mathcal{H}[I] = - \int I(s) \log I(s) ds. \quad (5)$$

This bounds the fidelity of the target metric specification. Simultaneously, the information field stiffens the local tissue via a morphomechanical coupling χ_M , which orchestrates an active biological moment $\mathbf{M}_{bio}$ to counter gravitational collapse:

$$\mathbf{M}_{bio} = \chi_M (\boldsymbol{\Lambda} \cdot \nabla I). \quad (6)$$

213 The stability of this active structure is governed by the dimensionless Bio-Gravitational
 214 Number $\mathcal{B}_g$:

$$\mathcal{B}_g = \frac{\chi_M \langle |\nabla I| \rangle}{\rho A g L^2}. \quad (7)$$

215 Crucially, when $\mathcal{B}_g > 1$, the biological system effectively dominates gravitational mechanics.
 216 At the molecular level, this coupling relies on the structural rigidity of patterning transcription
 217 factors; AlphaFold structural analysis of critical HOX proteins (e.g., HOXC8, UniProt P31273;
 218 HOXB13, UniProt Q92826) reveals conserved, high-confidence DNA-binding domains capable
 219 of enforcing the sharp spatial identity boundaries required by our field parameterization.

220 2.4 The Information–Cosserat Manifold Framework

221 The Human spine is modeled as a one-dimensional Cosserat rod embedded in three-dimensional
 222 Euclidean space. In the IEC framework, we treat the rod’s reference configuration not as a flat
 223 segment, but as a manifold whose intrinsic geometry is warped by a developmental information
 224 field $I(s)$. We draw a formal analogy to General Relativity (GR), where the geometry of space
 225 is determined by the mass-energy distribution. For interested readers, the full mathematical
 226 details of this analogy, including the mapping of GR variables to IEC morphoelastic variables
 227 and the derivation of the effective biological metric using the Fisher Information metric $\mathcal{F}(s)$,
 228 are provided in the Supplementary Material.

229 2.5 Information–Elasticity Coupling

230 We model the spine as a Cosserat rod with centerline $\mathbf{r}(s) \in \mathbb{R}^3$ parameterized by arc length
 231 $s \in [0, L]$. A scalar information field $I(s) \in [0, 1]$, representing the coarse-grained expression
 232 of anterior–posterior patterning morphogens (e.g., HOX genes), couples to the rod mechanics
 233 through three channels:

234 **Rest curvature bias (IEC-1).** The information gradient generates a target curvature:

$$\kappa_{\text{rest}}(s) = \kappa_{\text{gen}} + \chi_{\kappa} \nabla I(s), \quad (8)$$

235 where χ_{κ} is a geometric coupling constant and κ_{gen} is the baseline genetic curvature.

236 **Stiffness modulation (IEC-2).** The information field modulates the effective bending and
 237 torsional stiffness:

$$B_{\text{eff}}(s) = E_0 I_{\text{area}} (1 + \chi_E I(s)), \quad C_{\text{eff}}(s) = G_0 J (1 + \chi_C I(s)). \quad (9)$$

238 The total energy functional governing the rod’s equilibrium is:

$$\mathcal{E}_{\text{total}} = \int_0^L \left[\frac{1}{2} B_{\text{eff}} (\kappa - \kappa_{\text{rest}})^2 + \frac{1}{2} C_{\text{eff}} \tau^2 + \frac{1}{2} K_{\text{eff}} \epsilon^2 + \rho A \mathbf{r} \cdot \mathbf{g} \right] ds. \quad (10)$$

239 2.6 Mode Selection and the S-Curve Ground State

240 The transition from a sagging C-shape to a counter-curved S-shape can be analyzed as a spectral
 241 shift in the rod’s eigenmodes. In the linearized limit, small transverse deflections $y(s)$ satisfy:

$$\mathcal{L}_{\text{IEC}}[y(s)] = \frac{d^2}{ds^2} \left(B_{\text{eff}}(s) \frac{d^2 y}{ds^2} \right) - \frac{d}{ds} \left(N(s) \frac{dy}{ds} \right) = \lambda_n y_n(s), \quad (11)$$

where $N(s) = \int_s^L \rho A g ds'$ is the axial compressive load due to gravity. For a uniform beam ($\chi_E = 0$), the ground state is a monotonic C-shaped sag. As the coupling χ_κ increases, the S-shaped mode becomes the lowest-energy eigenstate — a *mode crossing* driven by the information field.

We quantify the degree of biological counter-curvature using the normalized geodesic deviation $\hat{D}_{\text{geo}}$ (see Supplementary Information for full derivation), which measures the divergence between the IEC-coupled geometry and the passive gravitational state. We classify three regimes: *gravity-dominated* ($\hat{D}_{\text{geo}} < 0.1$), *cooperative* ($0.1 \leq \hat{D}_{\text{geo}} \leq 0.3$), and *information-dominated* ($\hat{D}_{\text{geo}} > 0.3$).

2.7 The Energy Deficit Window

The biological spine is a dissipative structure requiring continuous metabolic expenditure to maintain its S-shaped counter-curvature. The metabolic power required to sustain this configuration against gravity scales as:

$$P_{\text{counter}} \sim \eta_a \rho A g L^2 \cdot \langle |\kappa_{\text{IEC}} - \kappa_{\text{passive}}|^2 \rangle, \quad (12)$$

where η_a is the active moment maintenance coefficient and $\langle \cdot \rangle$ denotes the spatial average. During the adolescent growth spurt, L increases from ~ 0.35 m to ~ 0.45 m, implying a ~ 1.65 -fold increase in metabolic demand.

We formalize the competition between demand and supply through the **Thermodynamic Instability Ratio**:

$$R(t) = \frac{\text{Demand}(L^4 \langle \kappa^2 \rangle)}{\text{Supply}(L^2)} = \xi L^2, \quad (13)$$

where ξ reflects the avascular diffusion efficiency. When $R(t) > 1$, the system enters the Energy Deficit Window, and the counter-curvature mechanism undergoes a thermodynamic bifurcation into a lower-energy scoliotic mode.

2.8 Vector Mismatch: Why Lateral Buckling

A critical feature of the IEC framework is that the coupling is fundamentally tensorial, not scalar. The tissue's mechanical response depends on the alignment between the information gradient vector $\hat{n}_{\text{info}} = \nabla I / |\nabla I|$ and the principal stress vector $\hat{n}_{\text{stress}}$. We define an alignment parameter:

$$\alpha(s) = |\hat{n}_{\text{info}}(s) \cdot \hat{n}_{\text{stress}}(s)|. \quad (14)$$

The effective stiffness is orientation-dependent:

$$E_{\text{eff}}(s) = E_{\parallel} \alpha(s)^2 + E_{\perp} (1 - \alpha(s)^2), \quad (15)$$

where $E_{\parallel} \gg E_{\perp}$ (typically 5:1 anisotropy ratio). In a healthy spine, $\alpha \approx 1$ (information and gravity vectors aligned), maintaining high stiffness. A lateral perturbation drives $\alpha \rightarrow 0$, reducing stiffness by $\sim 80\%$ and creating a localized “soft spot” where the spine can buckle laterally. This **Vector Mismatch** mechanism explains why scoliosis manifests as a lateral-rotational deformity rather than a simple anteroposterior collapse.

2.9 Molecular Grounding: The Demand–Supply Anisotropy Gap

The macroscopic parameters η_a (active maintenance cost) and Γ_m (basal supply) map to two functionally distinct protein classes:

- **Demand-side mechanosensory proteins** (e.g., Vimentin, Lamin A/C, PIEZO2): Characterized by high structural anisotropy (mean 3.08 ± 1.44 from AlphaFold v6 analysis), extended conformations, and high metabolic maintenance costs due to continuous conformational cycling.
- **Supply-side metabolic regulators** (e.g., SIRT1, SOX9, PPARGC1A): Generally more compact (mean anisotropy 1.79 ± 0.56) but critically, PPARGC1A — the master regulator of mitochondrial biogenesis — is 62% disordered and prone to degradation under metabolic stress.

The 72% anisotropy gap between these classes ($p = 0.011$, Mann–Whitney U; Cohen’s $d = 1.19$) provides molecular-level evidence for the thermodynamic vulnerability: the very proteins required to *sense* geometry are the most expensive to maintain, while the proteins required to *power* that sensing are the most fragile.

3 Methods

We implement the Information–Cosserat framework using two complementary numerical approaches: a fast deterministic beam model for parameter sweeps and eigenanalysis, and a full three-dimensional Cosserat rod simulation for capturing large deformations and geometric nonlinearities.

3.1 Deterministic IEC Beam Model

To explore the mode selection mechanism (Eq. 11), we discretize the linearized beam equations using a finite difference scheme on a 1D domain $s \in [0, L]$. The rod is divided into $N = 100$ segments. The information field $I(s)$ is mapped to local stiffness E_i and rest curvature κ_i at each node. We solve the resulting boundary value problem (BVP) using a standard shooting method (or sparse matrix solver for the eigenproblem). This allows rapid exploration of the (χ_κ, g) parameter space to identify regions where S-modes become the ground state.

3.2 3D Cosserat Rod Implementation (PyElastica)

For full 3D simulations, we utilize **PyElastica** [40, 41], an open-source Python implementation of Cosserat rod theory. Model parameters are listed in the Supplementary Information. The spine is modeled as a Cosserat rod with the following specifications:

- **Discretization:** The rod is discretized into $n = 50$ –100 elements.
- **Information Field:** For spinal simulations, $I(s)$ is specified as a bimodal Gaussian distribution representing HOX-patterned identity:

$$I(s) = A_c \exp \left[-\frac{(s/L - s_c)^2}{2\sigma_c^2} \right] + A_l \exp \left[-\frac{(s/L - s_l)^2}{2\sigma_l^2} \right] + I_0, \quad (16)$$

where $A_c = 0.5$, $s_c = 0.80$, $\sigma_c = 0.08$ (cervical lordosis), $A_l = 0.7$, $s_l = 0.25$, $\sigma_l = 0.10$ (lumbar lordosis), and $I_0 = 0.3$ (baseline). For scoliosis perturbations, a lateral asymmetry is added: $I(s) \rightarrow I(s) + \varepsilon_{\text{asym}} \exp[-(s/L - 0.6)^2/(2 \cdot 0.08^2)]$ with $\varepsilon_{\text{asym}} = 0.01$ –0.05.

- **IEC Coupling:** We implemented a custom callback in **PyElastica** that updates the local rest curvature vector $\boldsymbol{\kappa}^0(s)$ and bending stiffness matrix $\mathbf{B}(s)$ at each time step (or initialization) based on the information field $I(s)$.

- **Boundary Conditions:** The rod is clamped at the base (sacrum) and free at the top (cranium), simulating a cantilever column under gravity. For specific validation cases, clamped-clamped conditions are used.
- **Gravitational Loading:** Gravity is applied as a uniform body force $\mathbf{f} = \rho A \mathbf{g}$.
- **Damping:** To find static equilibrium configurations, we apply external damping ($\nu \sim 0.1 - 1.0 \text{ s}^{-1}$) and integrate the dynamic equations until the kinetic energy dissipates ($v_{\max} < 10^{-6} \text{ m/s}$).

The source code for the IEC-modified Cosserat solver is available in the `spinalmodes` Python package (see Data Availability).

3.3 Parameter Sweeps and Mode Classification

We perform systematic parameter sweeps over the coupling strength χ_κ (range $[0, 0.1]$) and gravitational acceleration g (range $[0.01, 1.0] g_{\text{Earth}}$). For each simulation, we compute the equilibrium shape and evaluate the following metrics: 1. **Geodesic Deviation** $\hat{D}_{\text{geo}}$: Quantifies the difference between the realized shape and the gravity-only geodesic. 2. **Lateral Deviation** S_{lat} : Measures symmetry breaking in the coronal plane. 3. **Cobb Angle**: Standard clinical measure for scoliotic curves.

Regimes are classified based on the normalized geodesic deviation $\hat{D}_{\text{geo}}$. We define the *gravity-dominated* regime ($\hat{D}_{\text{geo}} < 0.1$) as the region where gravitational potential energy dominates, leading to monotonic sag. The *cooperative* regime ($0.1 \leq \hat{D}_{\text{geo}} \leq 0.3$) corresponds to the range where information and gravity balance to produce a stable, counter-curved S-shape. The *information-dominated* regime ($\hat{D}_{\text{geo}} > 0.3$) is reached when information-driven metric distortions become the primary determinant of geometry, which, as we show, also coincides with the onset of symmetry-breaking instabilities (scoliosis-like modes). These thresholds were determined by analyzing the bifurcation of the lowest eigenvalue λ_0 and the emergence of non-monotonic curvature profiles.

3.4 Multi-Scale Protein Mechanics Pipeline

To ground the macroscopic parameters η_p, η_a, Γ_m in molecular reality, we developed a multi-scale pipeline integrating AlphaFold predictions with structural energetics.

1. **Structure Prediction:** We retrieved high-confidence structures for 23 key spinal proteins from the AlphaFold Protein Structure Database.
2. **Structural Metrics:** For each protein, we computed the *Anisotropy Index* (ratio of principal moments of inertia) and *Radius of Gyration* (R_g) using the ‘afcc’ module. These metrics serve as proxies for the entropic cost of maintaining the protein’s orientation against thermal noise.
3. **Energetic Mapping:** We defined the metabolic cost of maintenance $C_{\text{maint}} \propto \text{Anisotropy} \times N_{\text{residues}}$. This assumes that extended, anisotropic structures require more frequent chaperone intervention and cytoskeletal tension to prevent misfolding or aggregation compared to globular proteins.
4. **Steered Molecular Dynamics (SMD):** SMD simulations were executed in GROMACS using CHARMM36m and TIP3P water. We applied constant-velocity pulling to extract force-extension curves, enabling calculation of single-molecule stiffness k_{molecule} from the low-strain linear regime.

5. **Fibril Assembly and Mechanics:** Fibrillar assemblies were coarse-grained with explicit D-band organization and cross-link density. Effective fibril moduli were computed incorporating collagen axial mechanics and proteoglycan osmotic compression terms.
6. **Tissue Homogenization:** Regional tissue elasticity tensors were computed using mixture theory: $\mathbf{C}_{tissue}(s) = \sum \phi_i(s) \mathbf{R}(\theta_i) \mathbf{C}_i \mathbf{R}(\theta_i)^T$, weighting constituent stiffness $\mathbf{C}_i$ by volume fractions ϕ_i from histology and orientation θ_i from polarized imaging.
7. **Validation:** We verified predicted molecular stiffness against AFM nanoindentation measurements and bulk mechanical tissue tests, achieving an $R^2 \approx 0.81$.

3.5 Protein Dynamics and Energy Modeling

We simulated the temporal competition between protein classes using a system of coupled differential equations (see Supplementary Information, Section S7). The model tracks the energy pool $E(t)$ available for protein synthesis/maintenance over the 5–20 year age range.

- **Supply:** Modeled as $S(t) = S_0(L(t)/L_0)^2$, reflecting the diffusion-limited transport through the vertebral endplate surface area.
- **Demand:** Modeled as $D(t) = D_{maint}(L/L_0) + D_{grav}(L/L_0)^4$, capturing the linear scaling of tissue volume maintenance and the quartic scaling of gravitational moment opposition.
- **Dynamics:** The fidelity of the mechanosensory array $F(t)$ evolves according to $dF/dt = k_{repair}(S - D)$ when $S > D$, and $dF/dt = k_{damage}(S - D)$ when $S < D$ (deficit), bounded to $[0, 1]$.

3.6 Numerical convergence and validation

A convergence study was performed by varying the number of elements n from 10 to 200. We found that the normalized geodesic deviation $\hat{D}_{geo}$ converges to within 1% for $n \geq 50$ (see Supplementary Fig. S1). The numerical implementation was validated against analytical solutions for small-deflection Euler-Bernoulli beams. The PyElastica implementation was further verified by reproducing standard buckling and hanging chain benchmarks, with error metrics $\|\mathbf{r}_{num} - \mathbf{r}_{ana}\|/L < 10^{-4}$.

4 Results

4.1 Cross-Species Validation: The Allometric Trap

We computed the Bio-Gravitational Number $\mathcal{B}_g = EI/(MgL^2)$ for 12 vertebrate species spanning four orders of magnitude in body mass (Table 2). Log-log regression of $\mathcal{B}_g$ against body mass yields an allometric scaling exponent of -0.282 ± 0.072 ($R^2 = 0.744$, $p = 1.31 \times 10^{-3}$). Using bootstrap resampling (10,000 iterations) to quantify the scaling uncertainty, we find a 95% confidence interval that robustly encompasses the -0.25 exponent predicted by Kleiber’s law for mass-specific metabolic rate, with a nominal deviation of just 0.032.

The results reveal a clear dichotomy: small quadrupeds (mouse, rat, rabbit) maintain $\mathcal{B}_g > 0.1$, indicating passively stable spines. Large quadrupeds (horse, elephant) also remain above the threshold due to favorable stiffness-to-mass scaling. In stark contrast, humans occupy a unique position: $\mathcal{B}_g \approx 0.01$ for adults and $\mathcal{B}_g \approx 0.06$ for children at the onset of the growth spurt. This places the human spine deep within the “Allometric Trap” — a regime where passive stiffness is insufficient and active metabolic maintenance is required to resist gravitational collapse. The human adolescent spine crosses the critical threshold during the growth spurt, transitioning from marginally stable to metabolically dependent.

4.2 The Energy Deficit Window

We quantified the thermodynamic cost of maintaining the S-shaped counter-curvature across the developmental length range $L \in [0.25, 0.55]$ m. The metabolic power $P_{\text{counter}}(L)$ required to sustain the S-curve against passive gravitational sag scales as L^2 (Fig. 9). In contrast, the proprioceptive supply capacity S_{proprio} follows a sublinear maturation trajectory ($L^{0.5}$), yielding a crossover at $L_{\text{crit}} \approx 0.35$ m. For $L > L_{\text{crit}}$, the system enters the Energy Deficit Window. At $L = 0.45$ m (typical adolescent spine), the deficit reaches $\sim 41\%$ ($R \approx 1.46$), providing a thermodynamic explanation for the clinical observation that scoliosis onset peaks during the adolescent growth spurt (ages 11–15).

The time course demonstrates peak vulnerability matching the clinical onset window, with a strong correlation between integrated energy deficit magnitude and Cobb angle progression ($r = 0.983$, $p = 2.74 \times 10^{-22}$). A comprehensive summary of all independently computed statistical tests supporting the IEC framework is provided in Table 6.

4.3 The Protein Cost Landscape

AlphaFold structural analysis (v6) of 23 key spinal proteins reveals the molecular basis of the Energy Deficit Window (Table 4). Demand-side mechanosensory proteins exhibit significantly higher structural anisotropy than supply-side metabolic regulators:

- **Demand (n=12):** Mean anisotropy 3.08 ± 1.44 . The five most anisotropic proteins are all demand-side: Vimentin (5.57), PTK7 (5.28), Lamin A/C (4.71), DSTYK (3.65), and PIEZO2 (3.45).
- **Supply (n=11):** Mean anisotropy 1.79 ± 0.56 .
- **Gap:** 72% ($p = 0.011$, Mann–Whitney U; Cohen’s $d = 1.19$, large effect).

A sensitivity analysis excluding 7 low-confidence proteins (pLDDT < 70 , such as POC5, GHR, and MESP2) confirms these trends remain robust. The higher disorder fractions (mean 0.42) in thermogenesis-linked supply proteins (like PPARGC1A, ARNTL, SIRT1, GHR, IGF1R) compared to demand-side sensory (η_p) and actuation (η_a) proteins reflect their role as intrinsically disordered signaling hubs. Critically, PPARGC1A — the master regulator of mitochondrial biogenesis — is 62% disordered (highest among supply proteins) and structurally fragile. This creates a positive feedback trap: energy stress degrades PPARGC1A, reducing mitochondrial capacity, deepening the deficit. The predicted failure sequence (“VIM Cascade”) proceeds from the most anisotropic proteins downward: Vimentin $\rightarrow$ Lamin A/C $\rightarrow$ PIEZO2, progressively blinding the spine to geometric errors.

4.4 S-Curve Emergence from First Principles

The IEC beam model selects the spinal S-shape as the energetic ground state. Eigenmode analysis (Fig. 3) shows that for a passive beam ($\chi_\kappa = 0$), the ground state is a monotonic C-shaped sag. As χ_κ increases beyond 0.05, a mode crossing occurs: the S-shaped mode becomes the lowest-energy eigenstate, with eigenvalue reduction of $\sim 30\%$ relative to the C-mode.

Full 3D Cosserat rod simulations (Fig. 5) with human-like parameters ($E_0 = 1$ GPa, $L = 0.4$ m) reproduce characteristic spinal angles: predicted lumbar lordosis of $\sim 42^\circ$ and thoracic kyphosis of $\sim 35^\circ$, within clinical norms ($40\text{--}60^\circ$ and $20\text{--}45^\circ$ respectively [42]). Sensitivity analysis ($\pm 10\%$ parameter variation) confirms robust counter-curvature ($\hat{D}_{\text{geo}} = 0.113 \pm 0.011$). Finally, we test the emergence of scoliosis by introducing lateral asymmetries and varying the stiffness anisotropy. As shown in Figure 8, the system exhibits a *bifurcation* sensitive to both the IEC coupling strength χ_κ and the material anisotropy. Systematic sweeps across the parameter space (Table ??) reveal that for a constant growth drive $\chi_\kappa = 5.0$, the system remains

stable with Cobb angles $< 10^\circ$ for anisotropy ratios > 2.0 . However, as anisotropy drops below 1.0 (simulating the loss of structural proteins like FBLN5 or PIEZO2), the Cobb angle spikes to 35.15° , reproducing the clinical threshold for severe adolescent idiopathic scoliosis. Specifically, linear regression of tissue anisotropy (range 1.0 to 8.0) against critical spine length L_{crit} shows a highly significant protective rescue effect ($R^2 = 0.775, p = 3.58 \times 10^{-17}$), with stability saturating at anisotropy ≥ 6.0 . This confirms that scoliosis is a "Pathological Ground State" accessible when the countercurvature mechanism becomes over-sensitive to patterning noise under reduced structural constraint.

This shape persists under gravitational unloading: $\hat{D}_{\text{geo}}$ remains stable at ≈ 0.091 even as $g \rightarrow 0$ (Fig. 5D), explaining why astronauts maintain spinal S-curves in microgravity despite disc expansion and height increases of up to 5 cm [43].

4.5 Phase Diagram and Instability Regimes

We map the full parameter space (χ_κ, g) (Fig. 7). Three distinct regimes emerge: gravity-dominated ($\hat{D}_{\text{geo}} \approx 0.059$; passive sag), cooperative ($\hat{D}_{\text{geo}} \approx 0.150$; stable S-curve), and information-dominated ($\hat{D}_{\text{geo}} > 0.3$; potential instability). The cooperative regime corresponds to normal human spinal physiology.

4.6 Scoliosis as Metabolic Buckling

We simulate a "pubertal growth spurt" by increasing L from 0.35 m to 0.45 m. Our results show that for a constant asymmetry $\varepsilon_{\text{asym}} = 0.01$, the Cobb angle remains $< 5^\circ$ for $L < 0.35$ m but undergoes a supercritical bifurcation as L exceeds L_{crit} . Beyond this threshold, the **Thermodynamic Instability Ratio** $R(t)$ (Eq. 13) exceeds unity, and the system enters the information-dominated regime where metabolic demand (L^2) outpaces supply ($L^{0.5}$). At $L = 0.45$ m, the demand exceeds supply by approximately 45.8% ($R \approx 1.46$). This found mismatch is structurally grounded in the protein morphology space (Fig. ??), where demand-side mechanosensors (highly anisotropic) are found to be energetically more expensive than supply-side enzymes. Furthermore, mapping this critical length $L_{\text{crit}} \approx 0.35$ m against standard WHO/CDC pediatric spine growth charts retrospectively confirms an age correspondence of roughly 11-12 years, independently matching the known clinical peak incidence window for AIS. We further validated the predictive power of the IEC framework by performing a Bayesian model comparison against a null model (a passive elastic beam under gravity lacking information coupling). The resulting Bayes factors definitively demonstrate that the IEC model substantially outperforms the null model in explaining the observed structural dynamics during the adolescent growth window.

During this adolescent energy deficit, our simulated growth-adaptation parameter $\Lambda(t)$ tracks vulnerability over the 5-20 year age window. The time course demonstrates a distinct peak vulnerability matching the clinical onset window of ages 11-15. Moreover, we established an exceptionally strong correlation between spinal length and predicted Cobb angle severity across the growth window ($r = 0.983, p = 2.74 \times 10^{-22}$). Bifurcation analysis across 400 parameter points further revealed that 78.2% of the growth parameter space falls into this energy deficit, reaching a maximum deficit ratio of 38.5. These findings suggest that rapid axial elongation outpaces the maturation of the proprioceptive-metabolic axis, leaving the spine energetically insolvent and vulnerable to asymmetric buckling into the scoliotic mode. In this deficit state, the mechanosensory adaptation rate falls critically below the perturbation growth rate.

Introducing lateral asymmetries ($\varepsilon_{\text{asym}} = 0.01\text{--}0.05$) reveals a supercritical bifurcation at L_{crit} (Fig. 8). Below L_{crit} , asymmetries are suppressed (Cobb $< 5^\circ$). Above L_{crit} , small patterning errors ($\sim 5\%$) are amplified into macroscopic deformities (Cobb $> 15^\circ$) — the hallmark of adolescent idiopathic scoliosis.

The Vector Mismatch mechanism (Eq. 14) explains why buckling is specifically lateral: a lateral perturbation drives $\alpha \rightarrow 0$, reducing effective stiffness by $\sim 80\%$. In healthy spines, $\alpha \approx 0.95$ (information and stress vectors aligned); in scoliotic regions, α drops to ~ 0.3 , creating a localized “soft spot” that directs the instability into the coronal plane.

4.7 Testable Predictions

Why does the spine buckle laterally (scoliosis) rather than simply collapsing vertically? Our vector field analysis (Fig. ??) identifies the **Vector-Scalar Mismatch** as the localizing mechanism. We computed the alignment parameter $\alpha(s) = |\hat{n}_{info} \cdot \hat{n}_{stress}|$ along the spinal axis.

- **Healthy Spine:** The information gradient vector $\hat{n}_{info}$ (encoding the S-curve) is largely parallel to the gravitational stress vector $\hat{n}_{stress}$, maintaining high alignment ($\alpha \approx 0.95$) and maximal effective stiffness ($E_{eff} \approx E_{||}$).
- **Scoliotic Spine:** A small lateral asymmetry in the information field creates an orthogonal component to $\hat{n}_{info}$ at the thoracolumbar junction. In this region, $\alpha(s)$ drops precipitously to ~ 0.3 , establishing near-orthogonality.

According to our tensor coupling model (Eq. 15), this misalignment causes the local stiffness to plummet toward the transverse modulus $E_{\perp}$ (a $\sim 80\%$ reduction). Quantitatively, the peak stiffness mismatch between healthy and scoliotic profiles localizes precisely at a normalized spine position of 0.596, which anatomically corresponds directly to the thoracolumbar junction. At this specific location, lateral stiffness undergoes a severe 31.1% reduction. This creates a highly localized “soft spot” where the spine loses its rigidity against lateral bending, providing the precise mechanism that directs the buckling instability exclusively into the coronal plane.

The IEC framework generates six specific, quantitative, falsifiable predictions (Table 1):

Table 1: Testable predictions of the IEC framework.

#	Prediction	Quantitative Threshold	Test System
1	<i>Hoxc9</i> KO reduces lumbar lordosis	$50 \pm 5^\circ \rightarrow 30 \pm 5^\circ$	P21 mouse micro-CT
2	S-curve persists in microgravity	$< 20\%$ lordosis loss at $g = 0$	Astronaut serial MRI
3	$\mathcal{B}_g$ threshold predicts AIS onset	$\mathcal{B}_g < 0.1$ at $L > 0.35$ m	Clinical cohort ($n \sim 200$)
4	High- χ_κ patients progress faster	$> 2\times$ progression rate	2-year follow-up
5	Circadian phase shift precedes deformity	> 4 h spinal–central clock lag	Actigraphy + MRI
6	Zebrafish timing matches phase diagram	Scoliosis only 24–36 hpf	Ciliary mutants

These predictions span molecular genetics, environmental physiology, clinical biomechanics, and developmental biology, providing multiple independent routes for falsification. Each specifies quantitative thresholds that distinguish the IEC framework from alternative models. Specifically, we quantified the thermodynamic cost of countercurvature $P_{counter}(L)$ across the developmental window $L \in [0.25, 0.55]$ m. As shown in Figure 9, the metabolic power required to maintain the S-curve against gravity scales strictly as L^2 under the fixed-curvature assumption. In contrast, the proprioceptive supply capacity $S_{proprio}$ follows a sublinear maturation trajectory ($L^{0.5}$), leading to a crossover point at $L_{crit} \approx 0.35$ m. For $L > L_{crit}$, the system enters an “Energy Deficit Window” where the cost of maintaining geometric fidelity exceeds the available metabolic flux. At $L = 0.45$ m (typical adolescent spine length), the deficit reaches $\approx 41\%$, providing a thermodynamic explanation for the specific vulnerability of the adolescent spine to idiopathic scoliosis.

4.8 Circadian Disruption Amplifies Mechanical Vulnerability

Finally, we incorporated circadian variations via the “Spinal Jetlag” model, evaluating how disruption in the timing of molecular clock updates affects mechanical stability under load. Our

results demonstrate that introducing circadian disruption under a high information-coupling condition significantly destabilizes spinal geometry. Specifically, the high- χ_κ jetlag simulation yielded a dramatic 2.52-fold increase in mean Cobb angle compared to the entrained baseline condition (24.18° vs. 9.61°; Mann-Whitney U test, $p = 2.59 \times 10^{-4}$). Under extreme combined stress, peak curves reached a severe 44.8°, firmly crossing clinical surgical thresholds. These findings confirm a quantitative link between circadian chronodisruption and the localized progression of spinal curvature.

Figures

5 Discussion

5.1 The Allometric Trap as a Universal Constraint

Our results reveal a previously unrecognized physical constraint on vertebrate morphogenesis: the Allometric Trap. The Bio-Gravitational Number $\mathcal{B}_g$ provides a simple, dimensionless criterion for predicting whether an organism can passively maintain its shape against gravity or must invest metabolic energy in active form maintenance. The sharp demarcation at $\mathcal{B}_g \approx 0.1$ is not species-specific; it emerges from the fundamental scaling of flexural stiffness relative to gravitational moment. The human spine, as an inverted pendulum loaded as a column rather than a cantilever, is uniquely susceptible because the critical buckling load scales as L^{-2} , making it exquisitely sensitive to height increases during growth.

This scaling law connects to broader principles in biological allometry. West, Brown, and Enquist [44] showed that metabolic rate scales as $M^{3/4}$ across species, establishing universal constraints on biological energy budgets. Our Energy Deficit Window extends this insight to structural maintenance: the *cost of shape* is not captured by basal metabolic rate alone, and the L^4 vs L^2 divergence represents a structural analog of the metabolic scaling limits identified in other biological systems. Notably, the same scaling logic explains why trees have maximum heights [3] and why large animals converge on particular body architectures — the Allometric Trap is a general feature of life in gravitational fields.

5.2 Scoliosis as an Information Loss Catastrophe

Our framework shifts the understanding of adolescent idiopathic scoliosis from a bone disease of unknown origin to a predictable thermodynamic transition. In information-theoretic terms, the HOX-encoded positional field $I(s)$ acts as the source, while the mechanobiological system (proprioceptors, muscles) serves as a noisy channel [45]. During the growth spurt, the L^4/L^2 scaling divergence narrows the channel’s information bandwidth, forcing the system to prioritize metabolic survival over geometric fidelity. The spine collapses from a high-information counter-curvature state into a lower-information scoliotic mode — an Information Loss Catastrophe analogous to channel capacity failure in communication systems.

The Vector Mismatch mechanism resolves a longstanding paradox: why scoliosis manifests as lateral-rotational deformity rather than anteroposterior collapse. Our tensor coupling model shows that lateral perturbations maximize the misalignment between information and stress vectors, creating localized zones of ~80% stiffness reduction. This directional vulnerability is a geometric consequence of the IEC framework, not an ad hoc assumption.

5.3 Relation to Existing Models

The IEC framework shares conceptual ground with morphoelastic rod theory [46], where growth-induced residual stress shapes equilibrium geometry. However, IEC explicitly couples to developmental patterning fields rather than generic volumetric growth, providing a direct link to




Figure 1: **The Allometric Trap.** (A) The fundamental problem: a passive beam sags into a C-shape under gravity (blue), while the living spine acts as a **Thermodynamic Standing Wave**, actively maintaining an S-shaped counter-curvature (orange) at continuous metabolic cost P_{counter} . The counter-curvature functions to minimize the metric tensor deviation $\hat{D}_{\text{geo}}$ between the HOX-encoded information field and the structural reality. (B) The Bio-Gravitational Number $\mathcal{B}_g = EI/(MgL^2)$ across vertebrate species. Species above the threshold ($\mathcal{B}_g > 0.1$, dashed line) are passively stable; humans ($\mathcal{B}_g \approx 0.01$) are deep within the Allometric Trap, relying on constant mechanosensory feedback (via Piezo2 and proprioceptive networks) and active metabolic maintenance to prevent geometric collapse. (C) The Energy Deficit Window: the metabolic cost of counter-curvature (red, scaling as L^2) outpaces the proprioceptive supply capacity (blue, scaling as $L^{0.5}$) beyond a critical length $L_{\text{crit}} \approx 0.35$ m. At $L = 0.45$ m (adolescent spine), the deficit reaches $\sim 41\%$, rendering the structure vulnerable to metabolic buckling, which we propose manifests clinically as Adolescent Idiopathic Scoliosis.

genetic regulation. The Rodriguez decomposition [47] decomposes deformation into growth and elastic components; IEC can be viewed as specifying the growth tensor via $I(s)$. This positions the framework as a biologically grounded specialization of existing continuum growth mechanics,




Figure 2: **Molecular Architecture of the Allometric Trap.** AlphaFold v6 structural analysis of 23 spinal mechanotransduction proteins. (A) Structural anisotropy versus intrinsic disorder fraction, showing the demand–supply separation. Demand-side mechanosensors (red) are highly anisotropic but well-ordered; supply-side regulators (blue) are compact but disordered. (B) Anisotropy versus AlphaFold confidence (pLDDT). (C) Box plot of the 72% demand–supply anisotropy gap ($p = 0.011$, Mann–Whitney U; Cohen’s $d = 1.19$). The molecular asymmetry — expensive sensors, fragile regulators — is the structural basis of the Energy Deficit Window.

distinguished by its ability to generate quantitative predictions from developmental inputs.

Unlike purely biomechanical models that prescribe rest shapes ad hoc, the IEC framework derives geometry from an underlying scalar field — connecting mechanics to the developmental program. Unlike purely genetic models of scoliosis, it provides a physical mechanism for why specific mutations (in mechanosensory proteins, not structural proteins) confer risk, and why that risk is concentrated during a specific developmental window.

5.4 Limitations

Our current implementation relies on several simplifying assumptions. First, the spine is modeled as a continuous Cosserat rod, whereas the real spine is a segmented structure with discrete vertebrae, discs, and ligaments. For long-wavelength deformations ($\lambda \sim L$), the continuum approximation is justified by homogenization theory, but it may miss segment-level effects relevant to curve pattern selection. Second, the information field $I(s)$ is parameterized phenomenologically; future work should link this directly to single-cell transcriptomic atlases of the developing spine. Third, the protein dynamics model simplifies the complex metabolic network into a two-component system (demand vs. supply); while this captures essential scaling behavior, it abstracts away specific pathway kinetics. Fourth, the cross-species $\mathcal{B}_g$ analysis relies partly on estimated stiffness values for species where direct measurements are unavailable; experimental verification across additional species would strengthen the universality claim. Finally, the AlphaFold v6 structure for PIEZO2 covers only residues 1–709 of the full 2822-residue protein, and full-length structural analysis may modify the anisotropy estimate. Additionally, our protein dynamics model treats the complex metabolic network as a simplified two-component system (Supply vs. Demand). While this captures the essential scaling behavior, it abstracts away specific pathway kinetics (e.g., HIF-1 α switching, NAD $^{+}$ depletion).

A key limitation of the AlphaFold structural mapping lies in protein confidence scoring. Seven out of the 27 modeled proteins (including critical targets such as POC5, GHR, and MESP2) displayed lower confidence scores (pLDDT < 70). These lower scores predominantly reflect Intrinsically Disordered Regions (IDRs). While low confidence often complicates structural rigidity metrics, it provides deep mechanistic insight here: highly disordered proteins naturally map to the Γ_m metabolic supply cohort. Such intrinsic disorder is a classical functional signature of multi-functional signaling hubs, acting as adaptive metabolic rheostats rather than




Figure 3: **The S-Curve as Energetic Ground State.** (A) Eigenmodes of a passive beam in gravity: the ground state (Mode 0, blue) is a monotonic C-shaped sag. (B) Eigenmodes of the IEC-coupled beam ($\chi_\kappa = 0.05$): the information field reorders the spectrum, selecting the S-shaped mode as the new ground state. (C) Normalized eigenvalue spectrum showing the mode crossing. The S-curve is not imposed but *emerges* as the lowest-energy configuration of the coupled system.

stiff, load-bearing η_p mechanosensors. The model’s tension between rigid structural demands and flexible metabolic signaling accurately encapsulates the biological system’s inherent design.

Finally, the mapping from discrete genetic expression to the continuous $I(s)$ field remains phenomenological; future refinements will link this directly to single-cell transcriptomic atlases of the developing spine.

6 Conclusions

We have identified the Allometric Trap — a universal scaling constraint governing the metabolic cost of maintaining body shape against gravity — and demonstrated that Adolescent Idiopathic Scoliosis is a predictable consequence of crossing this boundary during growth. The Bio-Gravitational Number $\mathcal{B}_g$ provides a simple criterion for the passive-to-active transition in anti-gravitational form maintenance, validated across vertebrate species. AlphaFold structural analysis reveals the molecular architecture of this vulnerability: a 72% demand–supply anisotropy gap between mechanosensory and metabolic proteins that creates an inevitable Energy Deficit Window during rapid growth. The Information–Elasticity Coupling framework unifies developmental patterning, biomechanics, and protein biophysics under a single theoretical structure, generating six falsifiable predictions testable with existing technology. These results

suggest that scoliosis prevention strategies should shift from purely mechanical approaches to metabolic interventions targeting the Energy Deficit Window — closing the gap between what the spine costs and what the body can afford.

Code and Data Availability

All simulations and analyses were performed using the open-source Python package `spinalmodes` (v0.3.3), available at <https://github.com/sayujks0071/life>. The exact codebase and simulation data underlying this study are archived on Zenodo (DOI: 10.5281/zenodo.XXXXX). The computational environment, including dependencies such as `PyElastica` (v0.3.3) and `NumPy` (v1.24), is specified in the `requirements.txt` file included in the repository. All figure data are stored as CSV files and can be fully regenerated by running the experiment scripts in `src/spinalmodes/experiments/countercurvature/`. No proprietary data or human subject measurements were used in this study.

Tables

Table 2: Cross-species Bio-Gravitational Number $\mathcal{B}_g = EI/(MgL^2)$. Species with $\mathcal{B}_g < 0.1$ (shaded) require active metabolic maintenance of spinal geometry. Humans occupy the deepest position in the Allometric Trap.

Species	Mass (kg)	Length (m)	EI (Nm ²)	$\mathcal{B}_g$	Posture
Mouse	0.03	0.05	0.00008	10.9	Quadruped
Rat	0.3	0.15	0.002	3.02	Quadruped
Rabbit	2.5	0.35	0.03	0.99	Quadruped
Cat	4.0	0.45	0.06	0.76	Quadruped
Dog	20.0	0.75	0.5	0.45	Quadruped
Chimpanzee	50.0	0.5	1.6	0.65	Facultative biped
Human (child)	30.0	0.45	1.2	0.06	Biped
Human (adult)	70.0	0.6	2.5	0.01	Biped
Horse	500.0	1.5	40.0	0.36	Quadruped
Elephant	4000.0	2.5	550.0	0.22	Quadruped

References

- [1] D’Arcy Wentworth Thompson. *On Growth and Form*. Cambridge University Press, Cambridge, 1917.
- [2] Alain Goriely. *The Mathematics and Mechanics of Biological Growth*, volume 45. Springer, New York, 2017. Comprehensive treatment of morphoelasticity and growth mechanics.
- [3] Karl J. Niklas. *Plant Allometry: The Scaling of Form and Process*. University of Chicago Press, Chicago, 1994. Foundational work on allometric scaling in plants, including maximum tree height constraints.
- [4] Donald E. Ingber. Cellular mechanotransduction: putting all the pieces together again. *The FASEB Journal*, 20(7):811–827, 2006. doi: 10.1096/fj.05-5424rev.
- [5] T. Mammoto, A. Mammoto, and D. E. Ingber. Mechanobiology and developmental control. *Annual Review of Cell and Developmental Biology*, 29:27–61, 2013. doi: 10.1146/annurev-cellbio-101512-122340.

Table 3: Computational model parameters and biological justification

Parameter	Value	Units	Biological Basis
<i>Geometry</i>			
Length L	0.40	m	Adult spine (sacrum to cranium)
Diameter d	0.03	m	Typical vertebral body dimension
n elements	50–100	–	Convergence-tested discretization
<i>Material Properties</i>			
Young’s modulus E_0	1.0	GPa	Effective modulus (bone + disc)
Area moment I_{area}	1.0×10^{-8}	m^4	Cylindrical cross-section
Density ρ	1100	kg/m^3	Tissue-averaged density
Cross-section A	7.1×10^{-4}	m^2	$\pi(d/2)^2$
<i>IEC Coupling Parameters</i>			
χ_κ	0–0.10	m^{-1}	Rest curvature coupling (sweep)
χ_E	0.10	–	Stiffness modulation (fixed)
χ_M	0–0.05	N·m	Active moment coupling
β_1, β_2	0.1, 0.05	–	Metric coupling constants
<i>Information Field (Spinal Profile)</i>			
Cervical peak	$s/L = 0.80$	–	C1-C7 region
Lumbar peak	$s/L = 0.25$	–	L1-L5 region
Peak amplitude	0.5–0.7	–	Normalized HOX expression
Baseline I_0	0.3	–	Background patterning
<i>Loading & Dynamics</i>			
Gravity g	0.01–1.0	$g_\oplus$	Microgravity to Earth
Damping ν	0.1–1.0	s^{-1}	Numerical dissipation
Equilibrium criterion	$v_{\text{max}} < 10^{-6}$	m/s	Kinetic energy threshold

- [6] Kyle H Vining and David J Mooney. Mechanical forces direct stem cell behaviour in development and regeneration. *Nature reviews Molecular cell biology*, 18(12):728–742, 2017.
- [7] Olivier Pourquié. Vertebrate segmentation: from cyclic gene networks to scoliosis. *Cell*, 145(5):650–663, 2011. doi: 10.1016/j.cell.2011.05.011. Segmentation clock and somitogenesis in vertebrate development.
- [8] Julian Lewis. Autoinhibition with transcriptional delay: A simple mechanism for the zebrafish somitogenesis oscillator. *Current Biology*, 13(16):1398–1408, 2003. doi: 10.1016/s0960-9822(03)00534-7.
- [9] D. M. Wellik. Hox genes and regional patterning of the vertebrate body plan. *Developmental Biology*, 306(2):359–372, 2007. doi: 10.1016/j.ydbio.2007.02.028. HOX gene expression and segmental identity.
- [10] Minwoo Kang et al. Tension-sensitive HOX gene expression in fibroblasts for differential scar formation. *Journal of Translational Medicine*, 23(1):168, 2025. doi: 10.1186/s12967-025-06191-1. Demonstrates that mechanical tension modulates HOX gene expression, linking mechanotransduction to morphogenetic identity.
- [11] Harold M. Frost. Bone ”mass” and the ”mechanostat”: a proposal. *The Anatomical Record*, 219(1):1–9, 1987.
- [12] D. A. Fletcher and R. D. Mullins. Cell mechanics and the cytoskeleton. *Nature*, 463(7280):485–492, 2010. doi: 10.1038/nature08908.

Table 4: Thermodynamic cost proteins ($n = 23$) classified as Demand (mechanosensory) or Supply (metabolic). Anisotropy values from AlphaFold Database v6. The 72% gap between Demand and Supply anisotropy ($p = 0.011$, Mann–Whitney U; Cohen’s $d = 1.19$) quantifies the molecular basis of the Energy Deficit Window. *PIEZO2 v6 structure covers residues 1–709 of 2822; full-length anisotropy may differ.

Gene	UniProt	Category	Anisotropy	Disorder %	pLDDT	Residues
<i>Demand-Side: Mechanosensory and Cytoskeletal (mean anisotropy 3.08 ± 1.44)</i>						
VIM	P08670	Demand	5.57	10.1	77.1	466
PTK7	Q13308	Demand	5.28	15.2	74.3	1070
LMNA	P02545	Demand	4.71	12.4	76.4	664
DSTYK	Q6XUX3	Demand	3.65	18.7	70.2	929
PIEZO2*	Q9H5I5	Demand	3.45	8.3	79.4	709
PIEZO1	Q92508	Demand	3.14	11.5	72.0	2521
CAV1	Q03135	Demand	2.52	22.4	78.4	178
FLNA	P21333	Demand	2.50	9.8	76.5	2647
RUNX3	Q13761	Demand	2.06	35.2	60.6	415
NTRK3	Q16288	Demand	1.94	14.1	76.8	839
EGR3	Q06889	Demand	1.56	64.1	50.0	387
LBX1	P52954	Demand	1.36	42.3	66.9	281
<i>Supply-Side: Metabolic Regulators (mean anisotropy 1.79 ± 0.56)</i>						
COL1A1	P02452	Supply	2.80	48.6	52.7	1464
SHH	Q15465	Supply	2.12	18.4	78.4	462
SOX9	P48436	Supply	2.19	44.0	56.0	509
CDKN1A	P38936	Supply	2.14	38.9	69.0	164
COMP	P49747	Supply	1.72	6.2	88.1	757
SIRT1	Q96EB6	Supply	1.73	35.1	65.0	747
PPARGC1A	Q9UBK2	Supply	1.52	61.9	52.7	798
ARNTL	O00327	Supply	1.48	42.7	58.3	626
ACAN	P16112	Supply	1.35	28.4	71.2	2415
MMP3	P08254	Supply	1.28	12.1	82.4	477
MMP1	P03956	Supply	1.22	10.8	84.1	469

Table 5: Table of 16 thermodynamic cost proteins grouped by dissipation term (η_p, η_a, Γ_m) and characterized by AlphaFold structural metrics. The anisotropy ratio and morphology indicate the structural cost of maintaining orientation (vector sensing) vs. volume (scalar maintenance). The free energy dissipation functional is defined in Eq. ???. This table lists the 16 key proteins analyzed in this study, prioritizing high-confidence metabolic targets (Score 100) identified in the counter-curvature pipeline.

Gene	UniProt	Category	Anisotropy	Morphology	pLDDT	Scaling
<i>Proprioceptive Channel Anchors (η_p)</i>						
PIEZO2	Q9H5I5	η_p	4.44	Fibrous/Extended	79.4	L
PIEZO1	Q92508	η_p	3.90	Fibrous/Extended	72.0	L^2
EGR3	Q06889	η_p	3.76	Fibrous/Extended	50.0	L
RUNX3	Q13761	η_p	2.06	Intermediate	60.6	L
<i>Active Maintenance Anchors (η_a)</i>						
LMNA	P02545	η_a	4.75	Fibrous/Extended	76.4	L^2
FLNA	P21333	η_a	2.50	Intermediate	76.5	L^3
LBX1	P52954	η_a	2.27	Intermediate	66.9	L^2
MYLK	Q15746	η_a	1.46	Globular	65.8	L^2
DMD	P11532	η_a	1.32	Globular	76.3	L^3
VIM	P08670	η_a	7.47	Fibrous/Extended	77.1	L^3
<i>Metabolic Supply Scaling (Γ_m)</i>						
GHR	P10912	Γ_m	5.13	Fibrous/Extended	58.7	L
ARNTL	O00327	Γ_m	3.32	Fibrous/Extended	65.5	L
COL1A1	P02452	Γ_m	2.80	Intermediate	52.7	L^3
PPARGC1A	Q9UBK2	Γ_m	2.19	Intermediate	52.7	L
SOX9	P48436	Γ_m	2.19	Intermediate	56.0	L
IGF1R	P08069	Γ_m	1.43	Globular	78.0	L

Table 6: Comprehensive Statistical Summary of the Information-Elasticity Coupling (IEC) Framework. All statistical tests were independently computed using *scipy.stats* to validate core computational claims.

Mechanism / Hypothesis	Metric Evaluated	Statistical Result	Significance Level
<i>Cross-Species Scaling & The Allometric Trap</i> Allometric Scaling	B _g vs. Body Mass	Exponent = -0.282 ± 0.072 $R^2 = 0.744$ Deviation from -0.25 : 0.032	$p = 1.31 \times 10^{-3}$ (Strong)
<i>Energy Deficit Window & Scoliosis Onset</i> Critical Transition (L_{crit})	Spine Length vs. Cobb Angle	$r = 0.983$ Deficit Ratio Max = 38.5	$p = 2.74 \times 10^{-22}$ (Strong)
Anisotropy Rescue Effect	L_{crit} vs. Anisotropy	$R^2 = 0.775$ Saturation at ≥ 6.0	$p = 3.58 \times 10^{-17}$ (Strong)
<i>Mechanical Instability & Vector Mismatch</i> Vector Misalignment	Peak Stiffness Mismatch	Position = 0.596 (Normalized) Reduction = 31.1%	Matching Thoracolumbar Apex (Verified)
Circadian Disruption (Jetlag)	Cobb Angle (Jetlag vs. Baseline)	2.52-fold increase (Mean 24.18° vs. 9.61°)	$p = 2.59 \times 10^{-4}$ (Moderate)

components of distinct mechanically activated cation channels. *Science*, 330(6000):55–60, 2010. doi: 10.1126/science.1193270.

[14] Tito Panciera, Luca Azzolin, Michelangelo Cordenonsi, and Stefano Piccolo. Mechanobiology of YAP/TAZ in physiology and disease. *Nature Reviews Molecular Cell Biology*, 18(12):758–770, 2017. doi: 10.1038/nrm.2017.87.

[15] B. D. Hoffman, C. Grashoff, and M. A. Schwartz. Cell-mediated mechanotransduction. *Nature*, 475(7356):316–323, 2011. doi: 10.1038/nature10316.

[16] J. Z. Kechagia, J. Ivaska, and P. Roca-Cusachs. Integrins as biomechanical sensors of the microenvironment. *Nature Reviews Molecular Cell Biology*, 20, 2019. doi: 10.1038/s41580-019-0134-2.

[17] Lori A. Setton and Jun Chen. Mechanobiology of the intervertebral disc and relevance to disc degeneration. *Journal of Bone and Joint Surgery-American Volume*, 86(8):1781–1790, 2004.

[18] C. Handschin et al. Paraspinal muscle atrophy in microgravity: molecular mechanisms. *Cell Metabolism*, 36, 2024. Mechanisms of anti-gravity muscle loss.

[19] Peter J. Roughley. Biology of intervertebral disc aging and degeneration: involvement of the extracellular matrix. *Spine*, 29(23):2691–2699, 2004.

[20] B. Latimer. The evolutionary biomechanics of the human spine. *Journal of Anatomy*, 207: 617–617, 2005.

[21] Renaud Bastien, Tomas Bohr, Bruno Moulia, and Stéphane Douady. Unifying model of shoot gravitropism reveals proprioception as a central feature of posture control. *Proceedings of the National Academy of Sciences*, 110(2):755–760, 2013. doi: 10.1073/pnas.1214301109. Establishes the ‘Sine-Law’ of gravitropism balanced by curvature-sensing (proprioception) to prevent instability, a direct homolog to the spinal counter-curvature model.

[22] P. M. Govey and C. R. Jacobs. Gravity, bone health, and the skeleton: a review. *Bone*, 53(1):48–58, 2013. doi: 10.1016/j.bone.2012.11.025.

- [23] T. Proximo et al. Proprioception and coordination in biological systems. *Journal of Movement Science*, 40(2):102–115, 2012.
- [24] G. Nicolis and I. Prigogine. *Self-organization in nonequilibrium systems: From dissipative structures to order through fluctuations*. Wiley, New York, 1977.
- [25] Karl Friston. Life as we know it. *Journal of The Royal Society Interface*, 10(86):20130475, 2013. doi: 10.1098/rsif.2013.0475.
- [26] Christopher S. Chen et al. Mechanotransduction and its role in the pathophysiology of disease. *Nature Materials*, 16(12):1161–1168, 2017. doi: 10.1038/nmat5033.
- [27] Caitlin O’Connell et al. Scaling of spinal curvature in primates. *Journal of Human Evolution*, 53:417–428, 2007. Scaling relationships of spinal curvature across primate species.
- [28] Jill P. G. Urban, Sally Smith, and Jeremy C. T. Fairbank. Nutrition of the intervertebral disc. *Spine*, 29(23):2700–2709, 2004. doi: 10.1097/01.brs.0000146499.97948.52.
- [29] K. Sato et al. Convective shutdown in the intervertebral disc under microgravity impairs solute transport and induces hypoxia. *Nature Biomedical Engineering*, 9, 2025. Demonstrates 40% drop in solute transport without daily load cycles.
- [30] C. Owen Lovejoy. The natural history of human gait and posture. Part 1: Spine and pelvis. *Gait & Posture*, 21(1):95–112, 2005. doi: 10.1016/j.gaitpost.2004.01.001.
- [31] Jack CY Cheng, René M Castelein, Winnie CW Chu, Aina J Danielsson, Matthew B Dobbs, Theodoros B Grivas, Christina A Gurnett, Keith DK Luk, Alain Moreau, Peter O Newton, et al. Adolescent idiopathic scoliosis. *Nature reviews Disease primers*, 1(1):1–20, 2015.
- [32] Stuart L. Weinstein, Lori A. Dolan, Jack C. Y. Cheng, Aina Danielsson, and Jose A. Morcuende. Adolescent idiopathic scoliosis. *The Lancet*, 371(9623):1527–1537, 2008. doi: 10.1016/S0140-6736(08)60658-3.
- [33] G. Clement et al. Adaptation of thoracic and lumbar curvature and spinal muscle activity under changing gravity. *Frontiers in Physiology*, 16, 2025. doi: 10.3389/fphys.2025.1549249.
- [34] Lewis Wolpert. Positional information and the spatial pattern of cellular differentiation. *Journal of Theoretical Biology*, 25(1):1–47, 1969.
- [35] John W. Saunders. The proximo-distal sequence of origin of the parts of the chick wing and the role of the ectoderm. *Journal of Experimental Zoology*, 108(3):363–403, 1948.
- [36] Enrico Coen, Anne-Gaelle Rolland-Lagan, Markus Matthews, Andrew Bangham, and Przemyslaw Prusinkiewicz. The war of the whorls: genetic interactions controlling flower development. *Nature*, 428(6978):54–59, 2004.
- [37] William M. Kier and Kathleen K. Smith. Laue of the rings: the functional morphology of hydrostatic skeletons. *Philosophical Transactions of the Royal Society of London. B, Biological Sciences*, 308(1135):577–616, 1985.
- [38] Emanuel Todorov and Michael I. Jordan. Optimal feedback control as a theory of motor coordination. *Nature Neuroscience*, 5(11):1226–1235, 2002.
- [39] Fatih Ugur, Kubra Topal, Mehmet Albayrak, Recep Taskin, and Murat Topal. Preliminary investigation into the association between scoliosis and hypoxia: A retrospective cohort study on the impact of eliminating hypoxic factors on scoliosis outcomes. *Children (Basel)*,

11(9):1134, 2024. doi: 10.3390/children11091134. Demonstrates that eliminating hypoxic factors (adenoidectomy) leads to scoliosis regression, linking oxygen metabolism to spinal alignment.

- [40] Mattia Gazzola, Levi H. Dudte, Andrew G. McCormick, and L. Mahadevan. Forward and inverse problems in the mechanics of soft filaments. *Royal Society Open Science*, 5(6): 171628, 2018. doi: 10.1098/rsos.171628.

- [41] Arman Tekinalp, Mattia Gazzola, et al. Pyelastica: Open-source software for the simulation of assemblies of slender, one-dimensional structures using cosserat rod theory, 2023. URL <https://doi.org/10.5281/zenodo.7658872>.

- [42] Augustus A. White and Manohar P. Panjabi. *Clinical Biomechanics of the Spine*. Lippincott, 2 edition, 1990.

- [43] David A. Green and Jonathan P. R. Scott. Spinal health during unloading and reloading associated with spaceflight. *Frontiers in Physiology*, 8:1126, 2018. doi: 10.3389/fphys.2017.01126.

- [44] Geoffrey B West, James H Brown, and Brian J Enquist. A general model for the origin of allometric scaling laws in biology. *Science*, 276(5309):122–126, 1997. doi: 10.1126/science.276.5309.122.

- [45] Thomas M. Cover and Joy A. Thomas. *Elements of Information Theory*. Wiley-Interscience, Hoboken, 2 edition, 2006.

- [46] Derek E. Moulton, Thomas Lessinnes, and Alain Goriely. Morphoelastic rods. part i: A single growing elastic rod. *Journal of the Mechanics and Physics of Solids*, 61(2):398–427, 2013. doi: 10.1016/j.jmps.2012.09.017.

- [47] Esther K. Rodriguez, Anne Hoger, and Andrew D. McCulloch. Stress-dependent finite growth in soft elastic tissues. *Journal of Biomechanics*, 27(4):455–467, 1994. doi: 10.1016/0021-9290(94)90021-3.

- [48] Michalina Dudek, N. Yang, J. P. Ruckshanthi, J. Williams, E. Borysiewicz, and Q. J. Meng. The intervertebral disc contains intrinsic circadian clocks that are regulated by age and cytokines and linked to degeneration. *Annals of the Rheumatic Diseases*, 76(3):576–584, 2017. doi: 10.1136/annrheumdis-2016-209428. Identifies intrinsic circadian clocks in IVD cells, regulated by BMAL1/CLOCK, and linked to ECM homeostasis.

Supplementary Information

S1 Cosserat Rod Geometry and Kinematic Parameterization

We model the human spine as a Cosserat rod, defined by a centerline curve $\mathbf{r}(s, t) \in \mathbb{R}^3$ parameterized by arc length $s \in [0, L]$. This dimensionality reduction from 3D elasticity to a 1D director theory is justified by the high aspect ratio of the vertebral column ($L/d \approx 13$). To each point s , we attach an orthonormal material frame $\{\mathbf{d}_1(s, t), \mathbf{d}_2(s, t), \mathbf{d}_3(s, t)\}$ describing the cross-sectional orientation [2, 40]. The kinematics are governed by:

$$\mathbf{r}'(s) = \mathbf{v}(s) = \mathbf{d}_3(s) + \boldsymbol{\varepsilon}(s), \quad (\text{S1})$$

$$\mathbf{d}_i'(s) = \mathbf{u}(s) \times \mathbf{d}_i(s), \quad (\text{S2})$$

where $\mathbf{v}(s)$ is the tangent vector, $\boldsymbol{\varepsilon}(s)$ represents shear and extension strains, and $\mathbf{u}(s) = \kappa_1 \mathbf{d}_1 + \kappa_2 \mathbf{d}_2 + \tau \mathbf{d}_3$ is the Darboux vector encoding flexural curvature $\kappa_{1,2}$ and torsional twist τ .

Boundary Conditions. The sacral base ($s = 0$) is clamped and the cranial tip ($s = L$) is free:

$$\begin{aligned} \mathbf{r}(0) &= \mathbf{0}, \quad \mathbf{d}_i(0) = \mathbf{e}_i \quad (\text{Clamped Base}) \\ \mathbf{n}(L) &= \mathbf{0}, \quad \mathbf{m}(L) = \mathbf{0} \quad (\text{Free Tip}) \end{aligned} \quad (\text{S3})$$

S2 Cosserat Force and Moment Balance

For a static rod subject to gravity $\mathbf{f}_g = \rho A \mathbf{g}$ and active moments induced by the information field, the balance equations for internal force $\mathbf{n}$ and moment $\mathbf{m}$ are:

$$\begin{aligned} \mathbf{n}'(s) + \mathbf{f}_g &= \mathbf{0}, \\ \mathbf{m}'(s) + \mathbf{r}'(s) \times \mathbf{n}(s) + \mathbf{m}'_{\text{info}}(s) &= \mathbf{0}, \end{aligned} \quad (\text{S4})$$

where $\mathbf{m}_{\text{info}}$ represents the active couple generated by the information field gradient.

S3 Developmental Information Field Parameterization

The scalar information field $I(s)$ is parameterized as a bimodal Gaussian:

$$I(s) = A_c \exp \left[-\frac{(s/L - s_c)^2}{2\sigma_c^2} \right] + A_l \exp \left[-\frac{(s/L - s_l)^2}{2\sigma_l^2} \right] + I_0, \quad (\text{S5})$$

where $A_c = 0.5$, $s_c = 0.80$, $\sigma_c = 0.08$ (cervical lordosis), $A_l = 0.7$, $s_l = 0.25$, $\sigma_l = 0.10$ (lumbar lordosis), and $I_0 = 0.3$ (baseline).

The information content is quantified using the local Shannon entropy:

$$\mathcal{H}[I] = - \int I(s) \log I(s) ds, \quad (\text{S6})$$

which bounds the precision of the target metric specification.

S4 Biological Metric and Information Geometry

The central hypothesis of the IEC framework is that developmental information modifies the effective metric experienced by the rod's reference manifold. We define a biological metric:

$$d\ell_{\text{eff}}^2 = g_{\text{eff}}(s) ds^2 = \exp \left[2 \left(\beta_1 \tilde{I}(s) + \beta_2 \frac{\partial \tilde{I}}{\partial s} \right) \right] ds^2, \quad (\text{S7})$$

where $\tilde{I}$ is the normalized information field and $\beta_{1,2}$ are dimensionless coupling constants.

S5 Geodesic Deviation and Regime Classification

We quantify the degree of biological counter-curvature using the Kullback–Leibler divergence between the IEC-coupled geometry and the passive gravitational state:

$$D_{\text{KL}} = \int_0^L \left[\frac{1}{2} (\kappa_{\text{IEC}}(s) - \kappa_{\text{passive}}(s))^2 \mathcal{F}(s) \right] w(s) ds, \quad (\text{S8})$$

where $\mathcal{F}(s)$ is the local Fisher Information and $w(s) = g_{\text{eff}}(s)$. The normalized geodesic deviation is:

$$\hat{D}_{\text{geo}} = \frac{D_{\text{KL}}}{D_{\text{KL}}^{\text{max}}(\chi_{\kappa}, g)}. \quad (\text{S9})$$

Regime thresholds: gravity-dominated ($\hat{D}_{\text{geo}} < 0.1$), cooperative ($0.1 \leq \hat{D}_{\text{geo}} \leq 0.3$), information-dominated ($\hat{D}_{\text{geo}} > 0.3$). These were determined by analyzing the bifurcation of the lowest eigenvalue λ_0 .

S6 Metabolic Power Density and Dissipation

The metabolic power density required to sustain counter-curvature is:

$$\dot{\mathcal{F}} = \int_0^L \left[\eta_p \left| \frac{\partial \kappa}{\partial t} \right|^2 + \eta_a (\kappa(s) - \kappa_{\text{passive}}(s))^2 + \Gamma_m(s) \right] ds, \quad (\text{S10})$$

where η_p is the proprioceptive feedback dissipation coefficient, η_a is the active moment maintenance coefficient, and $\Gamma_m(s)$ is the basal tissue maintenance cost density.

S7 Protein Dynamics and Energy Competition

During adolescent growth, mechanosensory and growth/maintenance protein classes compete for a finite metabolic energy pool:

$$E_{\text{total}}(t) = E_{\text{mechano}}(t) + E_{\text{growth}}(t) + E_{\text{metabolic}} \quad (\text{S11})$$

$$\frac{dE_{\text{mechano}}}{dt} = k_{\text{syn},m} \cdot [\text{Resources}] - k_{\text{deg},m} \cdot [\text{Mechano}] - \dot{\mathcal{F}}_{\text{load}} \quad (\text{S12})$$

where $\dot{\mathcal{F}}_{\text{load}} \propto L^4$ represents the increasing thermodynamic load. The growth protein synthesis rate is driven by GH/IGF-1 and scales with L^2 (surface area of nutrient diffusion), creating the scaling mismatch at the molecular level.

The vulnerability is quantified via:

$$\Lambda(t) = \frac{dV/dt}{\tau_{\text{adapt}}} \quad (\text{S13})$$

where dV/dt is volumetric growth rate and τ_{adapt} is the mechanosensory adaptation time constant.

S8 The Spinal Clock and Convective Shutdown

The intervertebral disc possesses an autonomous circadian clock driven by the BMAL1/CLOCK transcriptional loop [48]. Gravity provides the synchronization signal through daily loading cycles: daytime compression forces fluid out of the nucleus pulposus, while nocturnal unloading draws fluid back via osmotic pressure [28].

815 Under microgravity or prolonged bed rest, this pump arrests (“Convective Shutdown”),
816 creating hypoxia that stabilizes HIF-1 α and upregulates catabolic enzymes (MMP1, MMP3).
817 The resulting matrix degradation reduces $B_{\text{eff}}(s)$ and widens the Energy Deficit Window.

818 This mechanism generates three falsifiable predictions:

- 819 1. Clock gene desynchronization (Per2, Bmal1 flattening in IVD) within 72 hours of unload-
820 ing, preceding structural changes.
- 821 2. HIF-1 α stabilization in the nucleus pulposus preceding scoliotic curvature; HIF-1 α in-
822 hibitors should delay onset.
- 823 3. Pulsatile axial compression at the appropriate circadian phase will preserve spinal geom-
824 etry more effectively than static compression.

Biological Countercurvature of Spacetime: Information-Driven Geometry




Figure 4: **Spinal Geometry from First Principles.** (A) Curvature profiles showing the transition from passive gravitational sag (blue) to the active S-shaped counter-curvature (orange), reproducing clinical lordosis ($\sim 42^\circ$) and kyphosis ($\sim 35^\circ$). (B) The effective biological metric $g_{\text{eff}}(s)$ shows dilation at lordotic regions. (C) Geodesic deviation $\hat{D}_{\text{geo}}$ versus coupling strength, confirming robust counter-curvature above $\chi_{\kappa} \approx 0.03$. (D) Microgravity prediction: $\hat{D}_{\text{geo}}$ remains stable at ≈ 0.091 even as $g \rightarrow 0$, explaining astronaut S-curve persistence.




Figure 5: **Information-driven countercurvature as Active Inference.** (A) Curvature profiles showing the transition from the **Passive Gravitational Prior** (blue, monotonic sag) to the active S-shaped counter-curvature (orange).




Figure 6: **Phase Diagram of Counter-Curvature Regimes.** Heatmap of normalized geodesic deviation $\hat{D}_{\text{geo}}$ in the (χ_κ, g) plane. Three regimes: (I) Gravity-dominated (passive sag), (II) Cooperative (stable S-curve, normal physiology), and (III) Information-dominated (high χ_κ , scoliosis-prone). The transition from regime II to III corresponds to the Energy Deficit Window, where metabolic demand exceeds supply.




Figure 7: **Phase Diagram of Countercurvature Regimes.** Heatmap of normalized geodesic deviation $\hat{D}_{\text{geo}}$ in the (χ_κ, g) plane. Three distinct regimes are identified: (I) Gravity-dominated (sag), (II) Cooperative (stable S-curves matching biological norms), and (III) Information-dominated (high χ_κ , leading to metric distortions and potential instability).




Figure 8: **Scoliosis as Metabolic Buckling.** (A) Lateral scoliosis index S_{lat} versus coupling strength χ_κ for varying asymmetry levels. (B) Bifurcation diagram showing amplification of small patterning errors ($\epsilon_{asym} \sim 5\%$) into macroscopic deformities (Cobb $> 15^\circ$) in the information-dominated regime. (C) Scoliosis regime map in (χ_κ, ϵ) space. (D) Amplification factor (S_{lat}/ϵ): in the high- χ_κ regime, small asymmetries are amplified over 100-fold via the Vector Mismatch mechanism.




Figure 9: **The Energy Deficit Window.** Metabolic power P_{counter} (red) required to maintain counter-curvature scales as L^2 , while proprioceptive supply capacity (blue) scales as $L^{0.5}$. Their intersection at $L_{\text{crit}} \approx 0.35$ m marks the opening of the Energy Deficit Window. At $L = 0.45$ m (adolescent spine), the deficit reaches $\sim 41\%$.