


age ~ minutes_played+bmi+injured
26.051 + (0)*minutes_played + (−0.04)*bmi + (1.485)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 4.8146 on 3 and 616 Degrees of Freedom
F p−value: 0.00254
R−Squared: 0.02291
Adjusted R−Squared: 0.018151996262596
Root MSE: 3.81999851714024
AIC: 3431.39382387347



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against bmi
age
bmi
20
25
30
35
20
22
24
26
28



age against injured
age
injured
35
30
25
20
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
363
185
424
Residuals
10
5
0
−5
−10
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Cook's distance
546
169
363
0.00
0.01
0.02
0.03
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
363
169
546
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
Cook's distance
Leverage h_{ii}
lm(formula)
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
0
0.005
0.01
0.015
0.02
0.025
3
2.5
2
1.5
1
0.5
0
546
169
363