


age ~ minutes_played+injured
25.106 + (0)*minutes_played + (1.489)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 7.15535 on 2 and 617 Degrees of Freedom
F p−value: 0.00085
R−Squared: 0.02267
Adjusted R−Squared: 0.0195002388253563
Root MSE: 3.82047212776428
AIC: 3429.54755188366



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against injured
age
injured
35
30
25
20
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
1863
424
10
5
0
−5
−10
Residuals
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
363
185
424
3
2
1
0
−1
−2
−3


Scale−Location
√|Standardized residuals|
0.0
0.5
1.0
1.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Cook's distance
546
169
156
0.04
0.03
0.02
0.01
0.00
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
156
169
546
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
546
169
156
Cook's distance
0.00
0.01
0.02
0.03
0.04
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)
2
1.5
1
0.5
0