

age ~ minutes_played+prev_injuries+bmi+injured


26.106 + (0)*minutes_played + (−0.089)*prev_injuries + (−0.039)*bmi + (1.544)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC


F−statistic: 3.70013 on 4 and 615 Degrees of Freedom
F p−value: 0.00549
R−Squared: 0.0235
Adjusted R−Squared: 0.0171491213660605
Root MSE: 3.81884543100475
AIC: 3433.01946701304



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against prev_injuries
age
prev_injuries
20
25
30
35
0
1
2
3
4



age against bmi
age
bmi
20
25
30
35
20
22
24
26
28



age against injured
age
injured
35
30
25
20
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
363
185
424
Residuals
10
5
0
−5
−10
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
169
363
546
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
363
169
546
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
546
169
363
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.5
1
1.5
2
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)