


age ~ minutes_played+prev_injuries+bmi
26.47 + (0)*minutes_played + (0.043)*prev_injuries + (−0.051)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.1673 on 3 and 616 Degrees of Freedom
F p−value: 0.91843
R−Squared: 0.00081
Adjusted R−Squared: −0.00405207412270547
Root MSE: 3.86295084950026
AIC: 3445.25866782399



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against prev_injuries
age
prev_injuries
20
25
30
35
0
1
2
3
4



age against bmi
age
bmi
20
25
30
35
20
22
24
26
28



Residuals vs Fitted
363
185
156
Residuals
10
5
0
−5
−10
24.8
24.9
25.0
25.1
25.2
25.3
25.4
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
363
185
156
−3
−2
−1
0
1
2
3


Scale−Location
363
185
156
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.8
24.9
25.0
25.1
25.2
25.3
25.4
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
185
256
363
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
363
185
256
3
2
1
0
−1
−2
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
256
363
185
0.020
0.015
0.010
0.005
0.000
Cook's distance
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)