


age ~ minutes_played+prev_injuries+injured
25.18 + (0)*minutes_played + (−0.09)*prev_injuries + (1.549)*injured


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 4.8914 on 3 and 616 Degrees of Freedom
F p−value: 0.00228
R−Squared: 0.02327
Adjusted R−Squared: 0.0185106825074713
Root MSE: 3.81930069727456
AIC: 3431.16728565012



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against prev_injuries
age
prev_injuries
20
25
30
35
0
1
2
3
4



age against injured
age
injured
35
30
25
20
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
363
185
424
10
5
0
−5
−10
Residuals
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Cook's distance
546
169
256
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
256
169
546
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
546
169
256
2
1.5
1
0.5
0
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)