


age ~ minutes_played+prev_injuries
25.254 + (0)*minutes_played + (0.042)*prev_injuries


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 0.12718 on 2 and 617 Degrees of Freedom
F p−value: 0.8806
R−Squared: 0.00041
Adjusted R−Squared: −0.00282807350150449
Root MSE: 3.86372787486262
AIC: 3443.50806643028



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against prev_injuries
age
prev_injuries
20
25
30
35
0
1
2
3
4



Residuals vs Fitted
185
363
381
10
5
0
−5
−10
Residuals
24.9
25.0
25.1
25.2
25.3
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
361
−3
−2
−1
0
1
2
3


Scale−Location
185
363
381
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.9
25.0
25.1
25.2
25.3
Fitted values
lm(formula)


Cook's distance
256
439
546
0.000
0.005
0.010
0.015
0.020
0.025
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
0.000
0.005
0.010
0.015
0.020
Leverage
lm(formula)
439
546
256
Cook's distance


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
256
546
439
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
Leverage h_{ii}
lm(formula)