


age ~ minutes_played+training_load+bmi
25.713 + ( 0)*minutes_played + ( 0.262)*training_load + ( −0.045)*bmi


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 1.05456 on 3 and 616 Degrees of Freedom
F p−value: 0.36788
R−Squared: 0.00511
Adjusted R−Squared: 0.000264355670647531
Root MSE: 3.85463847418529
AIC: 3442.58753596038



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against training_load
age
training_load
20
25
30
35
1
2
3
4
5



age against bmi
age
bmi
20
25
30
35
20
22
24
26
28



Residuals vs Fitted
363
185
424
10
5
0
−5
−10
Residuals
24.5
25.0
25.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
363
185
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.5
25.0
25.5
Fitted values
lm(formula)


Cook's distance
363
185
546
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
363
185
546
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
363
185
546
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0
0.5
1
1.5
0
0.005
0.01
0.015
0.02
0.025
0.03
Leverage $h_{ii}$
lm(formula)