age ~ minutes_played+training_load+injured

24.646 + (0)*minutes_played + (0.192)*training_load + (1.427)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 5.26664 on 3 and 616 Degrees of Freedom


F p−value: 0.00136


R−Squared: 0.02501


Adjusted R−Squared: 0.0202594407650436


Root MSE: 3.81589668092856


AIC: 3430.0616218828



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against training_load
age
training_load
20
25
30
35
1
2
3
4
5



age against injured
age
injured
35
30
25
20
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
363
185
424
Residuals
10
5
0
−5
−10
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Cook's distance
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0.035
546
169
156
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
156
169
546
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3.5
3
2.5
546
169
156
2
1.5
1
0.5
0
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
0.030
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)