


age ~ minutes_played+training_load
24.647 + (0)*minutes_played + (0.265)*training_load


A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05,
High R and Adj R,
Low AIC


F−statistic: 1.4893 on 2 and 617 Degrees of Freedom
F p−value: 0.22634
R−Squared: 0.0048
Adjusted R−Squared: 0.00157845446813809
Root MSE: 3.85522972364293
AIC: 3440.77772061986



age against minutes_played
age
minutes_played
20
25
30
35
0
1000
2000
3000
4000



age against training_load
age
training_load
20
25
30
35
1
2
3
4
5



Residuals vs Fitted
363
185
424
Residuals
10
5
0
−5
−10
24.4
24.6
24.8
25.0
25.2
25.4
25.6
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.4
24.6
24.8
25.0
25.2
25.4
25.6
Fitted values
lm(formula)


Cook's distance
353
413
546
0.020
0.015
0.010
0.005
0.000
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
3
2
1
0
−1
−2
413
546
353
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
Leverage
lm(formula)


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
353
546
413
1.5
1
0.5
0
0.015
0.010
0.005
0.000
Cook's distance
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)