age ~ prev_injuries+bmi+injured

25.749 + (−0.088)*prev_injuries + (−0.038)*bmi + (1.533)*injured

A good model will have: big F−stat (at least >3 depending on n, 10+ is nice), small p−value < 0.05, High R and Adj R,
Low AIC

F−statistic: 4.82209 on 3 and 616 Degrees of Freedom


F p−value: 0.00251


R−Squared: 0.02295


Adjusted R−Squared: 0.0181869580844743


Root MSE: 3.81993050493461


AIC: 3431.3717464079



age against prev_injuries
age
prev_injuries
20
25
30
35
0
1
2
3
4



age against bmi
age
bmi
20
25
30
35
20
22
24
26
28



age against injured
age
injured
35
30
25
20
0.0
0.2
0.4
0.6
0.8
1.0



Residuals vs Fitted
363
185
424
Residuals
10
5
0
−5
−10
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Q−Q Residuals
Standardized residuals
Theoretical Quantiles
lm(formula)
185
363
424
−3
−2
−1
0
1
2
3


Scale−Location
363
185
424
√|Standardized residuals|
0.0
0.5
1.0
1.5
24.5
25.0
25.5
26.0
26.5
Fitted values
lm(formula)


Cook's distance
169
363
546
0.000
0.005
0.010
0.015
0.020
0.025
Cook's distance
0
100
200
300
400
500
600
Obs. number
lm(formula)


Residuals vs Leverage
Standardized residuals
Leverage
lm(formula)
Cook's distance
363
546
169
3
2
1
0
−1
−2
−3
0.000
0.005
0.010
0.015
0.020
0.025


Cook's dist vs Leverage* $h_{ii}/(1-h_{ii})$
3
2.5
2
169
363
546
Cook's distance
0.000
0.005
0.010
0.015
0.020
0.025
1.5
1
0.5
0
0
0.005
0.01
0.015
0.02
0.025
Leverage h_{ii}
lm(formula)